

QUADRATIC VARIATION, p -VARIATION AND
INTEGRATION
WITH APPLICATIONS TO STOCK PRICE MODELLING

Rimas Norvaiša¹
Institute of Mathematics and Informatics, Vilnius, Lithuania

February 1, 2008

¹This research was partially supported by Lithuanian State Science and Studies Foundation Grant K014.

Contents

1	Introduction and results	1
2	The p-variation and integrals of Stieltjes type	12
2.1	Additive and multiplicative functions	12
2.2	The Wiener class	17
2.3	Extended Riemann-Stieltjes integrals	23
2.4	Chain rules for extended Riemann-Stieltjes integrals	30
2.5	The Left and Right Young integrals	33
3	Quadratic variation and restricted integrals of Stieltjes type	38
3.1	The quadratic λ -variation and λ -covariation	38
3.2	The Left and Right λ -integrals	51
3.3	Vector valued functions	64
3.4	The product λ -integral	73
3.5	Extended Doléans exponentials	83
3.6	Augmented linear λ -integral equations	90
3.7	The evolution representation	99
3.8	Non-existence of λ -integrals	108
4	Extension of the class of semimartingales	115
4.1	Stochastic processes and p -variation	115
4.2	Stochastic integral and the Left Young integral	121
4.3	$(2 - \epsilon)$ -semimartingales and the stochastic $\ominus$ -integral	123
4.4	Quadratic covariation of $(2 - \epsilon)$ -semimartingales	126
4.5	The Itô formula and the linear stochastic $\ominus$ -integral equation	129
4.6	Concluding remarks	134
5	Stock price modelling in continuous time	135
5.1	Evolutionary asset pricing model	135
5.2	Almost sure approximation by a discrete time model	140
5.3	Option pricing and hedging	144
5.4	Returns and the p -variation: examples	148
5.5	Estimating the p -variation index	152
A	Convergence of directed functions	159
	Bibliography	162

Chapter 1

Introduction and results

This long paper deals with several aspects of calculus for real-valued functions having a quadratic variation. Here the notion of a quadratic variation is a property of a “deterministic” function rather than the well-known property of a Brownian motion discovered by Lévy [63]. Let f be a *regulated* real-valued function on a closed interval $[a, b]$, that is, for $a \leq s < t \leq b$, there exist the limits $f(t-) := \lim_{u \uparrow t} f(u)$ and $f(s+) := \lim_{u \downarrow s} f(u)$. Let $\lambda = \{\lambda_m: m \geq 1\}$ be a nested sequence of partitions $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ of $[a, b]$ such that $\cup_m \lambda_m$ is dense in $[a, b]$; the class of all such λ is denoted by $\Lambda[a, b]$. We say that f has the *quadratic λ -variation* on $[a, b]$, if there is a regulated function V on $[a, b]$ such that $V(a) = 0$ and for each $a \leq s < t \leq b$, letting $x_i^m := (t_i^m \wedge t) \vee s$ for $i = 0, \dots, n(m)$,

$$V(t) - V(s) = \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} [f(x_i^m) - f(x_{i-1}^m)]^2, \quad (1.1)$$

$$V(t) - V(t-) = [f(t) - f(t-)]^2 \quad \text{and} \quad V(s+) - V(s) = [f(s+) - f(s)]^2. \quad (1.2)$$

The function $[f]_\lambda := V$, if exists, is nondecreasing and is called the *bracket function* of f . Actually, existence of V with the stated properties is equivalent to the definition of the quadratic λ -variation (see Definition 3.3 and Proposition 3.4). The class of all functions having the quadratic λ -variation on $[a, b]$ is denoted by $Q_\lambda[a, b]$. Föllmer [34] introduced a quadratic variation for a regulated and right-continuous function on $[0, \infty)$ to be a Radon measure on $[0, \infty)$ if exists as a limit of sums of point masses with respect to a sequence of partitions of a half line $[0, \infty)$ with vanishing mesh. The bracket function $[f]_\lambda$ being a function rather than a measure, allows us to use a Stieltjes type integration for all classes of functions appearing in this paper. We notice that the quadratic variation of a function defined by Wiener [109], if exists, does not depend on partitions, and is closely related to the local 2-variation of a function defined by Love and Young [68]. To recall its definition let f be a function on $[a, b]$, and let $0 < p < \infty$. For a partition $\kappa = \{t_i: i = 0, \dots, n\}$ of $[a, b]$, let

$$s_p(f; \kappa) := \sum_{i=1}^n |f(t_i) - f(t_{i-1})|^p. \quad (1.3)$$

We say that f has the *local p -variation*, $1 < p < \infty$, if the limit

$$\lim_{\kappa, \mathfrak{P}} s_p(f; \kappa) \quad (1.4)$$

exists in the sense of refinements of partitions (see Appendix A for details). Wiener's quadratic variation of f means that (1.4) holds for $p = 2$ and the limit in the sense of refinements is replaced by the limit as the mesh of partitions tends to zero. Notice that the limit (1.1) is taken along a fixed sequence of partitions. Both Wiener's quadratic variation and the local 2-variation do *not* exist for almost all sample functions of a Brownian motion.

Again let f be a function on $[a, b]$. For $0 < p < \infty$, the p -variation of f is defined by

$$v_p(f; [a, b]) := \sup \{s_p(f; \kappa) : \kappa \in \Xi[a, b]\}, \quad (1.5)$$

where $\Xi[a, b]$ denotes the class of all partitions of $[a, b]$. We say that f has *bounded p -variation* if $v_p(f; [a, b]) < \infty$, and denote the class of all such functions by $\mathcal{W}_p[a, b]$. Usefulness of the p -variation property hinges on its relation to the Stieltjes type integrability theory established around the late of thirties by L. C. Young in connection to applications in the theory of Fourier series. Let g and f be a pair of functions on $[a, b]$ such that

$$f \in \mathcal{W}_p[a, b], \quad g \in \mathcal{W}_q[a, b] \quad \text{and} \quad p^{-1} + q^{-1} > 1 \quad \text{for} \quad p, q > 0. \quad (1.6)$$

Then by the L. C. Young Theorem on Stieltjes integrability [113] the integral $\int_a^b g df$ exists (a) in the sense of Riemann-Stieltjes if f and g have no common discontinuities, (b) in the sense of refinement Riemann-Stieltjes if f and g have no common discontinuities on the right and no common discontinuities on the left, (c) always in the sense defined by L. C. Young. The functional calculus developed by many authors later on has a full-fledged application to the class of functions having bounded p -variation for some $0 < p < 2$ (see [25]). In particular, the theory applies path by path to Lévy processes without a Gaussian component, and to a fractional Brownian motion having the Hurst exponent $H \in (1/2, 1)$ because their almost all sample functions have bounded p -variation for some $0 < p < 2$. Let $B = \{B(t) : t \geq 0\}$ be a standard Brownian motion, and let $X = \{X(t) : t \geq 0\}$ be a stochastic process both defined on a complete probability space $(\Omega, \mathcal{F}, \Pr)$. Then by L. C. Young's Theorem on Stieltjes integrability, for almost all $\omega \in \Omega$, the Riemann-Stieltjes integral

$$(RS) \int_0^T X(t, \omega) dB(t, \omega) \quad (1.7)$$

exists provided for some $0 < p < 2$, X has almost all sample functions of bounded p -variation on $[0, T]$. This fact was used by P. Lévy to solve suitable Riemann-Stieltjes integral equations with respect to a Brownian motion (see e.g. [64]). In (1.7), X may be an α -stable Lévy process with $0 < \alpha < 2$, or X may be a fractional Brownian motion with the Hurst index $1/2 < H < 1$. However, one cannot replace X in (1.7) by B (see Section 3.8 for details).

A function f on $[a, b]$ having bounded p -variation for some $1 \leq p < 2$ also has the quadratic λ -variation for each sequence of partitions $\lambda = \{\lambda_m : m \geq 1\} \in \Lambda[a, b]$ such that

$$\{t \in (a, b) : [f(t+) - f(t)][f(t) - f(t-)] \neq 0\} \subset \cup_m \lambda_m. \quad (1.8)$$

In this case, the bracket function $[f]_\lambda$ at $t \in [a, b]$ is a sum of squared jumps over $[a, t]$ (see Corollary 3.15 below). The converse is not true: given $\lambda \in \Lambda[0, 1]$, almost all sample functions of a Brownian motion have the quadratic λ -variation on $[0, 1]$, and have the p -variation on $[0, 1]$ unbounded for each $1 \leq p \leq 2$. Another related example is provided by a subclass of self-affine functions introduced by Kôno [57] (see Example 3.6 below). An extension of the p -variation calculus to Brownian motion path like functions require a more refined constructions than the Riemann-Stieltjes integral. The quadratic λ -variation and related integral constructions introduced below make the core of our attempts towards building a desirable calculus.

Evolution representation problem. The results in this paper are motivated by and applied to an evolution representation problem. To sketch the problem, let $\mathbb{B} = (\mathbb{B}, \|\cdot\|)$ be a Banach algebra with unit $\mathbb{I}$. A family $U = \{U(t, s): a \leq s \leq t \leq b\} \subset \mathbb{B}$ is called an *evolution* in $\mathbb{B}$ if it is multiplicative, that is,

$$\begin{cases} U(r, t)U(t, s) = U(r, s) & \text{for } a \leq s \leq t \leq r \leq b, \\ U(t, t) = \mathbb{I} & \text{for } a \leq t \leq b. \end{cases} \quad (1.9)$$

The notion of an evolution generalizes the concept of a one-parameter semigroup of bounded linear operators on a Banach space. The classical Hille-Yosida theorem describes any strongly continuous, contractive semigroup in terms of its generator. Analogous pairing results have been established for evolutions under various conditions on the function U_a defined by $U_a(t) := U(t, a)$ for $t \in [a, b]$. For example, U_a is analytic, or almost differentiable, or of bounded variation is one set of conditions. Other type of conditions imply that the generators will be unbounded operators in some cases, bounded in other.

Without additional assumptions a representation of an evolution may not be unique. We are interested in an evolution U which satisfies the condition: there is a function A on the simplex $\{(s, t): a \leq s \leq t \leq b\}$ such that

$$A(t, s) = \lim_{\kappa, \mathfrak{P}} S(U; [s, t], \kappa), \quad (1.10)$$

where $S(U; [s, t], \kappa) := \sum_{i=1}^n [U(t_i, t_{i-1}) - \mathbb{I}]$ for a partition $\kappa = \{t_i: i = 0, \dots, n\}$ of $[s, t]$. Actually in this paper we are more concerned with the case when the limit under refinement of partitions in (1.10) is replaced by a limit along a sequence of partitions. In Chapter 5 below, an evolution with such a condition and $\mathbb{B}$ being the set of real numbers, is used to build a continuous time stock price model. According to Dobrushin [17], condition (1.10) appeared in earlier works of M. Fréchet and E. B. Dynkin, in connection of extending the Chapman-Kolmogorov equations to more general transition probabilities of a Markov process. The function A defined by (1.10) is additive, and so letting $h := A(\cdot, a)$, we have $A(t, s) = h(t) - h(s)$ for each $a \leq s \leq t \leq b$. In some cases an evolution U satisfying (1.10) can be represented by a product integral with respect to the function $h = A(\cdot, a)$. For a $\mathbb{B}$ -valued function h on $[a, b]$, and for a partition $\kappa = \{t_i: i = 0, \dots, n\}$ of $[a, b]$, let

$$P(h; \kappa) = P(h; [a, b], \kappa) := \prod_{i=1}^n [\mathbb{I} + h(t_i) - h(t_{i-1})]. \quad (1.11)$$

The *product integral* on $[a, b]$ with respect to h is defined to be the limit of products $P(h; [a, b], \kappa)$, if it exists in the sense of refinements of partitions κ , that is, using the notion of a limit of a directed function (see Appendix A),

$$\mathfrak{Y}_a^b(\mathbb{I} + dh) := \lim_{\kappa, \mathfrak{P}} P(h; [a, b], \kappa). \quad (1.12)$$

Then the *product integral representation* of an evolution U means the relation

$$U(t, s) = \mathfrak{Y}_s^t(\mathbb{I} + dh), \quad a \leq s \leq t \leq b,$$

where $h = A(\cdot, a)$ and A is defined by (1.10). Apparently the product integral representation of an evolution U was known for a long time, but in all such cases an evolution U satisfies the

condition:

$$\sup \left\{ \sum_{i=1}^n \|U(t_i, t_{i-1}) - \mathbb{I}\| : a = t_0 < t_1 < \cdots < t_n = b, \ n \geq 1 \right\} < \infty. \quad (1.13)$$

More often a product integral representation of an evolution was used under various additional smoothness conditions (see Masani [75] for a survey).

A most general product integral representation of an evolution U satisfying (1.13) was proved by Mac Nerney [71] in the form of a duality between multiplicative and additive functions. Let $\mathcal{U}_1 = \mathcal{U}_1([a, b]; \mathbb{B})$ be the class of all evolutions U such that (1.13) holds, and let $\mathcal{W}_1 = \mathcal{W}_1([a, b]; \mathbb{B})$ be the class of all $\mathbb{B}$ -valued functions on $[a, b]$ with bounded variation, that is, with bounded 1-variation. By Theorem 3.3 of Mac Nerney [71], there is a reversible mapping $\mathcal{P}$, from $\mathcal{W}_1$ onto $\mathcal{U}_1$, such that for $U \in \mathcal{U}_1$ and $h \in \mathcal{W}_1$, the following three statements are equivalent:

- (a) $U = \mathcal{P}(h)$;
- (b) $U(t, s) = \lim_{\kappa, \mathfrak{P}} P(h; [s, t], \kappa)$ for each $a \leq s \leq t \leq b$;
- (c) $h(t) - h(s) = \lim_{\kappa, \mathfrak{P}} S(U; [s, t], \kappa)$ for each $a \leq s \leq t \leq b$.

Thus (b) gives the product integral representation of the evolution U provided (1.13) and (1.10) hold.

We illustrate the preceding representation in the case when an evolution U on $\mathbb{B} = \mathbb{M}^d$, the Banach algebra of $d \times d$ matrices for some integer $d \geq 1$, is continuous and translation-invariant. That is, suppose that the function $U_a \equiv U(\cdot, a)$, is continuous and

$$U(t+u, s+u) = U(t, s) \quad \text{for each } a \leq s \leq t \leq t+u \leq b. \quad (1.14)$$

Then letting $T(t-s) := U(t, s)$ for $a \leq s \leq t \leq b$, by (1.9), it follows that the family $T = \{T(u) : u \geq 0\}$ is a continuous one-parameter semigroup:

$$\begin{cases} T(u)T(v) = T(u+v) & \text{for each } u, v \geq 0, \\ T(0) = \mathbb{I}. \end{cases}$$

In this case it is well-known that there exists the right-derivative $T'(0)$ and $T(u) = \exp\{uT'(0)\}$ for $u \geq 0$, where the exponential $\exp x := \sum_{k=0}^{\infty} x^k/k!$ for $x \in \mathbb{M}^d$ (see e.g. Theorem I.2.9 and Proposition I.2.8 in Engel and Nagel [30]). For any partition $\kappa = \{t_i : i = 0, \dots, n\}$ of $[a, b]$, we have

$$\sum_{i=1}^n \|U(t_i, t_{i-1}) - \mathbb{I}\| = \sum_{i=1}^n \|T(t_i - t_{i-1}) - \mathbb{I}\| \leq (b-a) \sup_{u>0} u^{-1} \|T(u) - \mathbb{I}\| < \infty,$$

and so (1.13) holds for U . Therefore the above stated Mac Nerney [71] evolution representation when applied to U gives the representation $U = \mathcal{P}(h)$ with the function h such that $h(t) - h(s) = (t-s)T'(0)$ and

$$\mathfrak{Y}_s^t(\mathbb{I} + dh) = U(t, s) = T(t-s) = \exp\{(t-s)T'(0)\} = \exp\{h(t) - h(s)\}$$

for all $a \leq s \leq t \leq b$.

As we have seen, an evolution U in a Banach algebra $\mathbb{B}$ has the product integral representation provided (1.13) holds. Also in this case, an evolution gives a solution of a linear integral equation

as follows. A $\mathbb{B}$ -valued function f on $[a, b]$ is invertible if $f(t)$ is invertible for each $t \in [a, b]$, and f^{inv} is the reciprocal function of f defined by $f^{\text{inv}}(t) := [f(t)]^{-1}$ for $t \in [a, b]$. If the function $U_a \equiv U(\cdot, a)$ has bounded reciprocal U_a^{inv} , then the reverse $\mathcal{P}^{-1}$ of Mac Nerney's mapping $\mathcal{P}$, has the Left Cauchy-Stieltjes integral representation. Let f, g be $\mathbb{B}$ -valued functions defined on $[a, b]$, and for any partition $\kappa = \{t_i: i = 0, \dots, n\}$ of $[a, b]$, let

$$S_{LC}(f, g; [a, b], \kappa) := \sum_{i=1}^n [f(t_i) - f(t_{i-1})]g(t_{i-1}).$$

The *Left Cauchy-Stieltjes* integral is defined by

$$(LCS) \int_a^b df g := \lim_{\kappa, \mathfrak{P}} S_{LC}(f, g; [a, b], \kappa)$$

provided the limit exists in the sense of refinement of partitions. Let the reciprocal U_a^{inv} of the function U_a be defined and bounded on $[a, b]$. By (1.9), for any partition $\kappa = \{t_i: i = 0, \dots, n\}$ of $[a, b]$, we have

$$S(U; [a, b], \kappa) = \sum_{i=1}^n [U(t_i, t_{i-1}) - \mathbb{I}] = \sum_{i=1}^n [U_a(t_i) - U_a(t_{i-1})]U_a^{\text{inv}}(t_{i-1}) = S_{LC}(U_a, U_a^{\text{inv}}; [a, b], \kappa).$$

Also (1.13) holds if U_a has bounded variation. Therefore the integral $(LCS) \int_a^b dU_a U_a^{\text{inv}}$ exists, and the representation $U = \mathcal{P}(h)$ holds with the function h defined by

$$h(t) = (LCS) \int_a^t dU_a U_a^{\text{inv}}, \quad a \leq t \leq b.$$

A representation of a solution of a linear Left Cauchy-Stieltjes integral equation in terms of a flow in $\mathbb{B}$ is another well-known characterization of the evolution representation. Suppose that $U = \mathcal{P}(h)$ for some $U \in \mathcal{U}_1([a, b]; \mathbb{B})$ and $h \in \mathcal{W}_1([a, b]; \mathbb{B})$. Then by Theorem 4.1 of Mac Nerney [71], in the class $\mathcal{W}_1$, the linear Left Cauchy-Stieltjes integral equation

$$f(t) = f(a) + (LCS) \int_a^t dh f, \quad a \leq t \leq b,$$

has the unique solution $f(t) = U(t, a)f(a)$, $t \in [a, b]$.

Replacing boundedness of the variation of the function U_a , which is condition (1.13) above, by a weaker condition provides an interesting challenge to the evolution representation problem. Suppose that the reciprocal function U_a^{inv} of the function $U_a \equiv U(\cdot, a)$ exists and is bounded. In this case, boundedness of the p -variation of U_a for some $1 \leq p < \infty$ is one such weaker condition. The main advantage of this condition is a well established integration theory for a class of functions with bounded p -variation. Freedman [37] applied L. C. Young's results and ideas to extend the product integral representation of an evolution U provided the function U_a is continuous and has bounded p -variation for some $1 \leq p < 2$. Using the Banach fixed point theorem Freedman [37, Theorem 5.1] proved the existence of the product integral as a unique solution of the linear Riemann-Stieltjes integral equation. A non-linear Riemann-Stieltjes integral equation with respect to a continuous function of bounded p -variation, $1 \leq p < 2$, was solved by Lyons [70].

Dudley and Norvaiša [23, Part II] started their investigation of the product integral (1.12) with respect to a Banach algebra valued function h from scratch. By their Theorem 4.4, for a real valued function f , the product integral with respect to f exists and is non-zero if and only if f is regulated function satisfying (a) and (b):

(a) f has local 2-variation, that is, (1.4) holds with $p = 2$;

(b) for any $a \leq s < t \leq b$,

$$\Delta^+ f(s) := f(s+) - f(s) \neq -1 \neq f(t) - f(t-) =: \Delta^- f(t). \quad (1.15)$$

Moreover, if the product integral exists then for any $a \leq s < t \leq b$,

$$\mathfrak{Y}_s^t(1 + df) = e^{f(t) - f(s)} \prod_{[s,t]} [(1 + \Delta^- f)(1 + \Delta^+ f)] e^{-\Delta^- f - \Delta^+ f},$$

where $\Delta^- f(s) := 0$ and $\Delta^+ f(t) := 0$. Assuming that f is a regulated function, a new proof of this fact is given below by Theorem 3.36. For a function h with values in a Banach algebra $\mathbb{B}$, let $\mathcal{P}_a(h) = \{\mathcal{P}_a(h; t) : a \leq t \leq b\}$ be the indefinite product integral defined by

$$\mathcal{P}_a(h; t) := \mathfrak{Y}_a^t(\mathbb{I} + dh), \quad a \leq t \leq b,$$

and let $\mathcal{W}_p([a, b]; \mathbb{B})$ be the Banach space of all $\mathbb{B}$ -valued functions with bounded p -variation on $[a, b]$. The main result in [23, Part II] is that the mapping $\mathcal{P}_a$ from $\mathcal{W}_p([a, b]; \mathbb{B})$ into itself, is Fréchet differentiable and $\mathcal{P}_a$ is analytic when restricted to right- or left-continuous functions h . To formulate the evolution representation in this case, for a $\mathbb{B}$ -valued function f , let $\mathcal{L}_a(f) = \{\mathcal{L}_a(f; t) : a \leq t \leq b\}$ be the indefinite Left Young integral on $[a, b]$ defined by

$$\mathcal{L}_a(f; t) := (LY) \int_a^t df f^{\text{inv}}, \quad a \leq t \leq b,$$

provided the Left Young integral $(LY) \int_a^b df f^{\text{inv}}$ exists (see Definition 2.21 below). The following is Theorem 6.14 in [23, Part II].

Theorem 1 *Let $0 < p < 2$, let U be an evolution in $\mathbb{B}$ such that $U_a \equiv U(\cdot, a) \in \mathcal{W}_p([a, b]; \mathbb{B})$, and let the reciprocal function U_a^{inv} exists and is bounded. Then (a), (b) and (c) hold:*

- (a) *the indefinite Left Young integral $\mathcal{L}_a(U_a)$ on $[a, b]$ is defined and is in $\mathcal{W}_p([a, b]; \mathbb{B})$;*
- (b) *the product integral on $[a, b]$ with respect to $\mathcal{L}_a(U_a)$ is defined;*
- (c) *the function $\mathcal{L}_a(U_a)$ in $\mathcal{W}_p([a, b]; \mathbb{B})$ is unique up to an additive function such that*

$$U(t, s) = \mathfrak{Y}_s^t(\mathbb{I} + d\mathcal{L}_a(U_a)), \quad a \leq s \leq t \leq b.$$

The main result of this paper is a solution of a representation problem for a class of evolutions U in $\mathbb{R}$ such that $U_a \equiv U(\cdot, a)$ has the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$. Before formulating this result we summarize other results.

(λ, p) -decomposable function and $(2 - \epsilon)$ -semimartingale. Let $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$. As it was mentioned earlier, a function f having bounded p -variation also has the quadratic λ -variation provided a two sided jump of f occurs only at a point of a partition of λ . By the preceding remarks such functions in the class $\mathcal{Q}_\lambda[a, b]$ of all functions having the quadratic λ -variation may be considered as special: for such functions we have a calculus quite similar to the classical calculus for smooth functions (see [25]). Therefore when proving something for a

function having the quadratic λ -variation it is desirable to make the new result concerted with the calculus applicable to functions having bounded p -variation. To this aim we consider sums $g + h$ such that the p -variation is bounded for h and may be unbounded for g , but g still has the quadratic λ -variation. We also found useful to assume when $p > 1$ that g has bounded q -variation for some $q > 2$ such that $p^{-1} + q^{-1} > 1$. This condition is satisfied by our typical examples mentioned earlier. Under an additional mild condition on jumps of g and h , the sum $g + h$ also has the quadratic λ -variation (see Corollary 3.16 below). We also treat stochastic processes with a similar decomposability property except that the first component of such a decomposition has to be a local martingale.

Recall that $\mathcal{W}_p[a, b]$, $0 < p < \infty$, denotes the class of all real-valued functions f on a closed bounded interval $[a, b]$ with bounded p -variation defined by (1.5). Also, a real-valued function f on $[a, b]$ is regulated on $[a, b]$ if for $a \leq s < t \leq b$, there exist the limits $f(t-) := \lim_{u \uparrow t} f(u)$ and $f(s+) := \lim_{u \downarrow s} f(u)$. If for some $p < \infty$, f has bounded p -variation on $[a, b]$ then f is regulated on $[a, b]$. A point $t \in (a, b]$ is *left discontinuity point* of f if $\Delta^- f(t) := f(t) - f(t-) \neq 0$, and a point $s \in [a, b)$ is *right discontinuity point* of f if $\Delta^+ f(s) := f(s+) - f(s) \neq 0$. The class of all regulated functions on $[a, b]$ is denoted by $\mathcal{R}[a, b]$. For $1 \leq p < \infty$, let

$$\text{dual}(\mathcal{W}_p)[a, b] := \begin{cases} \cup \{\mathcal{W}_q[a, b] : \frac{1}{p} + \frac{1}{q} > 1\} & \text{if } 1 < p < \infty, \\ \mathcal{R}[a, b] & \text{if } p = 1. \end{cases} \quad (1.16)$$

Thus $f \in \text{dual}(\mathcal{W}_p)[a, b]$ with $1 < p < \infty$ if f has bounded q -variation for some $q < p/(p-1)$, and $f \in \text{dual}(\mathcal{W}_1)[a, b]$ if f is a regulated function. Now we are ready to define the class of (λ, p) -decomposable considered in Chapter 3.

Definition 2 Let f be a regulated function on $[a, b]$, let $\lambda \in \Lambda[a, b]$, and let $1 \leq p < 2$. We say that f is (λ, p) -decomposable on $[a, b]$, or $f \in \mathcal{D}_{\lambda, p}[a, b]$, if there exists a pair (g, h) of regulated functions on $[a, b]$ such that $g \in Q_\lambda[a, b] \cap \text{dual}(\mathcal{W}_p)[a, b]$, $h \in \mathcal{W}_p[a, b]$, $g(a) = h(a) = 0$ and $f \equiv C + g + h$ for some constant C . Any such pair (g, h) is called a (λ, p) -dual pair (λ, p) -decomposing f . The set of all (λ, p) -dual pairs which (λ, p) -decompose f is denoted by $D_{\lambda, p}(f)$.

Among all (λ, p) -dual pairs we select those pairs which have left or right discontinuities at points of partitions of λ . Given $\lambda = \{\lambda_m : m \geq 1\} \in \Lambda[a, b]$, we often assume to hold

$$\text{either } N_{(a, b)}(\Delta^+ f) := \{t \in (a, b) : (\Delta^+ f)(t) \neq 0\} \subset \cup \lambda \quad (1.17)$$

$$\text{or } N_{(a, b)}(\Delta^- f) := \{t \in (a, b) : (\Delta^- f)(t) \neq 0\} \subset \cup \lambda, \quad (1.18)$$

where $\cup \lambda := \cup_m \lambda_m$. We say that the *right discontinuity points of f are accessible by λ* if $N_{(a, b)}(\Delta^+ f) \subset \cup \lambda$. Similarly, we say that the *left discontinuity points of f are accessible by λ* if $N_{(a, b)}(\Delta^- f) \subset \cup \lambda$. Clearly, the first condition holds for each $\lambda \in \Lambda[a, b]$ if f is right-continuous, and the second one holds if f is left-continuous. In some cases we use a weaker assumption (see (1.8) above): we say that the *two-sided discontinuity points of f are accessible by λ* if

$$N_{(a, b)}(\Delta^- f) \cap N_{(a, b)}(\Delta^+ f) = \{t \in (a, b) : (\Delta^- f \Delta^+ f)(t) \neq 0\} \subset \cup \lambda. \quad (1.19)$$

Now let $f \in \mathcal{D}_{\lambda, p}[a, b]$ for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$. We denote by $D_{\lambda, p}^+(f)$ the set of all pairs $(g, h) \in D_{\lambda, p}(f)$ such that

$$N_{(a, b)}(\Delta^+ g) \cup N_{(a, b)}(\Delta^+ h) \subset \cup \lambda. \quad (1.20)$$

Similarly, $D_{\lambda,p}^-(f)$ denotes the set of all pairs $(g, h) \in D_{\lambda,p}(f)$ such that (1.20) holds with Δ^+ replaced by Δ^- .

Suppose that $f \equiv C + g + h$ for some $(g, h) \in D_{\lambda,p}^+(f) \cup D_{\lambda,p}^-(f)$. By Corollary 3.16, the function f has the quadratic λ -variation, and its bracket function is given by

$$[f]_\lambda(t) = [g]_\lambda^c(t) + \sum_{(a,t]} (\Delta^- f)^2 + \sum_{[a,t)} (\Delta^+ f)^2 \quad a \leq t \leq b. \quad (1.21)$$

Thus the representation $f \equiv C + g + h$ is not unique because $(f - f(a), 0) \in D_{\lambda,p}^+(f) \cup D_{\lambda,p}^-(f)$ is another (λ, p) -decomposition of f . Nevertheless several constructions defined in this paper for functions f of the class $\mathcal{D}_{\lambda,p}[a, b]$, do not depend on a representation of f by a (λ, p) -dual pair in $D_{\lambda,p}^+(f) \cup D_{\lambda,p}^-(f)$. For example, if $D_{\lambda,p}^+(f) \cup D_{\lambda,p}^-(f) \neq \emptyset$ then the continuous part of the bracket function $[f]_\lambda$ does not depend on a (λ, p) -decomposition. This follows from formula (1.21). Therefore we define $[f]_\lambda^c := [g]_\lambda^c$ with a g being such that $(g, h) \in D_{\lambda,p}^+(f) \cup D_{\lambda,p}^-(f)$. Below we define two integrals, the Left and Right λ -integrals, with respect to a function f of the class $\mathcal{D}_{\lambda,p}[a, b]$ which under mild additional assumptions also do not depend on nonuniqueness of (λ, p) -decomposability of f .

Another nice feature of the class of (λ, p) -decomposable functions is that it is closed under taking a composition with the C^2 class functions.

Theorem 3 *For $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, let f be a (λ, p) -decomposable function on $[a, b]$, and let ϕ be a C^2 class function. If $D_{\lambda,p}^+(f) \cup D_{\lambda,p}^-(f) \neq \emptyset$ then the composition $\phi \circ f$ is a (λ, p) -decomposable function.*

The proof of this statement given at the end of Section 3.2, shows that the composition $\phi \circ f$ has at least two different (λ, p) -decompositions if $D_{\lambda,p}^+(f) \cap D_{\lambda,p}^-(f) \neq \emptyset$.

Let $1 \leq p < 2$. We say that a stochastic process X is a p -semimartingale if $X - X(0) = M + A$, where $M(0) = A(0) = 0$, M is a local martingale and A is a stochastic process having almost all sample functions of bounded p -variation on each bounded time interval. In the case $p = 1$, X is usually called a semimartingale. We also say that X is a $(2 - \epsilon)$ -semimartingale if it is a p -semimartingale for some $1 \leq p < 2$. This class of processes may be considered as a stochastic analog of the class of (λ, p) -decomposable functions. In Chapter 4 we prove several elementary properties for such processes. Given a $(2 - \epsilon)$ -semimartingale X we define the stochastic $\ominus$ -integral with respect to X to be a sum of two integrals with respect to each component of a decomposition and show that values of such sums do not depend on a decomposition in most interesting cases. Analogous uniqueness result is proved with respect to the continuous part of a quadratic variation of a $(2 - \epsilon)$ -semimartingale. Finally in Chapter 4 we prove that in the class of all $(2 - \epsilon)$ -semimartingales, an extended stochastic Doléans exponential is a unique solution of the linear stochastic $\ominus$ -integral equation.

Integrals of functions having the quadratic λ -variation. As we mentioned earlier, the L. C. Young Theorem on Stieltjes integrability does not hold in general when the integrand and integrator both have the quadratic λ -variation. For such functions we use restricted Stieltjes type integrals defined to be limits, if exist, along the sequence of partitions λ . An existence of such an integral with a special form of the integrand can be proved by replicating a proof of Itô's formula (see Fölmer [34]). In this paper we follow the same route except that we do not assume functions to be right or left continuous, and use a Riemann-Stieltjes type integral to integrate with respect to a bracket function.

Let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ with $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ for $m \geq 1$. For regulated functions g and f on $[a, b]$, the *Left Cauchy λ -integral* $(LC) \int g d_\lambda f$ is defined on $[a, b]$ if there exists a regulated function Φ on $[a, b]$ such that $\Phi(a) = 0$ and for each $a \leq s < t \leq b$, letting $x_i^m = (t_i^m \wedge t) \vee s$ for $i = 0, \dots, n(m)$,

$$\Phi(t) - \Phi(s) = \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} g(x_{i-1}^m) [f(x_i^m) - f(x_{i-1}^m)],$$

$$\Delta^- \Phi(t) = (g_- \Delta^- f)(t) \quad \text{and} \quad \Delta^+ \Phi(s) = (g \Delta^+ f)(s).$$

If such Φ exists then we also say that g is Left Cauchy λ -integrable on $[a, b]$ with respect to f , and for $a \leq s < t \leq b$, let $(LC) \int_s^t g d_\lambda f := \Phi(t) - \Phi(s)$. In fact we define slightly more general integral when g may not be regulated (Definition 3.17). Such extension is used to prove the Black-Scholes formula in Section 5.3. Similarly the Right Cauchy λ -integral is defined (Definition 3.18).

The following statement is a corollary of Theorem 3.24. We omit its proof because it is symmetric to the proof of Proposition 3.25 concerning the Right Cauchy λ -integral.

Proposition 4 *Let f be a regulated function on $[a, b]$, and let $\lambda \in \Lambda[a, b]$ be such that the right discontinuity points of f are accessible by λ . The following statements about f are equivalent:*

- (a) f has the quadratic λ -variation on $[a, b]$;
- (b) for a C^1 class function ψ , the composition $\psi \circ f$ is Left Cauchy λ -integrable on $[a, b]$ with respect to f , and for any $a \leq s < t \leq b$,

$$\begin{aligned} (\Psi \circ f)(t) &= (\Psi \circ f)(s) + (LC) \int_s^t (\psi \circ f) d_\lambda f + \frac{1}{2} (RS) \int_s^t (\psi' \circ f) d[f]_\lambda^c \\ &\quad + \sum_{(s,t]} \left\{ \Delta^- (\Psi \circ f) - (\psi \circ f)_- \Delta^- f \right\} + \sum_{[s,t)} \left\{ \Delta^+ (\Psi \circ f) - (\psi \circ f) \Delta^+ f \right\}, \end{aligned}$$

where the two sums converge unconditionally and $\Psi(u) = \Psi(0) + \int_0^u \psi(x) dx$, $u \in \mathbb{R}$;

- (c) f is Left Cauchy λ -integrable on $[a, b]$ with respect to itself.

If any of the three statements holds, then for $a \leq t \leq b$,

$$(LC) \int_a^t f d_\lambda f = \frac{1}{2} \left\{ f^2(t) - f^2(a) - [f]_\lambda(t) \right\}.$$

Now we define an integral with respect to a (λ, p) -decomposable function. Let $\lambda \in \Lambda[a, b]$, let $1 \leq p < 2$ and let f be a (λ, p) -decomposable function. For a regulated function F and a set $A \subset D_{\lambda,p}(f)$, we say that the *Left λ -integral* $(L) \int F d_\lambda f$ is defined on $[a, b]$ with respect to A if for any $(g, h) \in A$, on the right side

$$(L) \int F d_\lambda f := (LC) \int F d_\lambda g + (LY) \int F dh$$

the Left Cauchy λ -integral and the Left Young integral are both defined on $[a, b]$, and sums of values at $a \leq s < t \leq b$ do not depend on $(g, h) \in A$. In Proposition 3.23 we show that the Left

λ -integral $(L) \int F d_\lambda f$ is defined on $[a, b]$ with respect to $D_{\lambda,p}^+(f) \neq \emptyset$ provided $F \in \text{dual}(\mathcal{W}_p)[a, b]$ and $(LC) \int F d_\lambda g$ is defined on $[a, b]$ for some $(g, h) \in D_{\lambda,p}^+(f)$. Analogous statement holds for the Right λ -integral defined symmetrically.

Next we summarize results concerning an extension of the product integral (1.12). Let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ with $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ for $m \geq 1$. Let f be a regulated function on $[a, b]$. By Proposition 3.39, the product λ -integral $\mathfrak{Y}(1 + d_\lambda f)$ is defined on $[a, b]$ if and only if there exists a regulated function H on $[a, b]$ such that $H(a) = 1$, $|H| \gg 0$ and for $a \leq s < t \leq b$, letting $x_i^m = (t_i^m \wedge t) \vee s$ for $i = 0, \dots, n(m)$,

$$H(t)/H(s) = \lim_{m \rightarrow \infty} \prod_{i=1}^{n(m)} [1 + f(x_i^m) - f(x_{i-1}^m)],$$

$$H(t)/H(t-) = 1 + (\Delta^- f)(t) \quad \text{and} \quad H(s+)/H(s) = 1 + (\Delta^+ f)(s).$$

If such H exists then for $a \leq s < t \leq b$, let $\mathfrak{Y}_s^t(1 + d_\lambda f) := H(t)/H(s)$. By Proposition 3.40, if the product integral $\mathfrak{Y}(1 + df)$ exists and is non-zero on $[a, b]$, and if the two-sided discontinuity points of f are accessible by λ , then the product λ -integral $\mathfrak{Y}(1 + d_\lambda f)$ is defined on $[a, b]$, and the two integrals have the same values.

Let f be a regulated function on $[a, b]$ having the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$. The Doléans exponential $\mathcal{E}_{\lambda,a}(f)$ on $[a, b]$ is defined by

$$\mathcal{E}_{\lambda,a}(f; t) := \begin{cases} \exp \{f(t) - f(a) - \frac{1}{2}[f]_\lambda^c(t)\} \prod_{[a,t]} (1 + \Delta f) e^{-\Delta f} & \text{if } t \in (a, b] \\ 1 & \text{if } t = a, \end{cases}$$

where the product on the right side is defined by (3.83). By Theorem 3.42, if the two-sided discontinuity points of f are accessible by λ , and if (1.15) holds, then the product λ -integral $\mathfrak{Y}(1 + d_\lambda f)$ is defined on $[a, b]$ and for $a \leq t \leq b$,

$$\mathfrak{Y}_a^t(1 + d_\lambda f) = \mathcal{E}_{\lambda,a}(f; t).$$

For $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, let f be a function on $[a, b]$ which is (λ, p) -decomposable and $D_{\lambda,p}^+(f)$ is nonempty. Thus f has the quadratic λ -variation. Consider the forward linear Left λ -integral equation

$$F(t) = 1 + (L) \int_a^t F d_\lambda f, \quad a \leq t \leq b, \quad (1.22)$$

provided the Left λ -integral is defined on $[a, b]$ with respect to $D_{\lambda,p}^+(f)$. Using a chain rule it is easy to check that the Doléans exponential $\mathcal{E}_{\lambda,a}(f)$ is a solution of (1.22). To show its uniqueness in the class $\text{dual}(\mathcal{W}_p)[a, b]$ we augment equation (1.22) as formulated in Definition 3.46. The augmented equation is solved by Theorem 48 in Section 3.6. Analogous result for the backward linear Right λ -integral equation is proved by Theorem 51 in the same Section 3.6.

Evolution and the quadratic λ -variation. Now we are prepared to formulate the main result. For a function f on $[a, b]$, $f \gg 0$ means that there is a constant $C > 0$ such that $f(t) \geq C$ for each $t \in [a, b]$. By Proposition 3.53, an *evolution* U in $\mathbb{R}$ is regulated if and only if $U_a \equiv U(\cdot, a)$ is a regulated function on $[a, b]$ and $|U_a| \gg 0$. Let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ with $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ for $m \geq 1$. By Proposition 3.55, a regulated evolution U in $\mathbb{R}$ has

the λ -generator if and only if there exists a regulated function G on $[a, b]$ such that $G(a) = 0$ and for each $a \leq s < t \leq b$, letting $x_i^m = (t_i^m \wedge t) \vee s$ for $i = 0, \dots, n(m)$,

$$G(t) - G(s) = \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} [U(x_{i-1}^m, x_i^m) - 1],$$

$$\Delta^- G(t) = U_a(t)/U_a(t-) - 1 \quad \text{and} \quad \Delta^+ G(s) = U_a(s+)/U_a(s) - 1.$$

The function G is called the λ -generator of U . This function generalizes the notion of the generator of a one-parameter semigroup as follows. Let U be a continuous translation-invariant evolution on $[a, b]$ as defined by (1.14). Then letting $T(y-z) := U(z, y)$ it follows that the family $T = \{T(u): u \geq 0\}$ is a continuous one-parameter semigroup. By the results of Mac Nerney [71], we have that for each $\lambda \in \Lambda[a, b]$, U has the λ -generator G such that $G(t) - G(s) = (t-s)T'(0)$ for $a \leq s < t \leq b$.

To define a mapping analogous to Mac Nerney's duality mapping $\mathcal{P}$ between $\mathcal{W}_1$ and $\mathcal{U}_1$, we use two sets of functions having the quadratic λ -variation. For $\lambda \in \Lambda[a, b]$, let

$$L_\lambda[a, b] := \{f \in Q_\lambda[a, b]: f(a) = 1, \quad f \gg 0 \quad \text{and} \quad N_{(a,b)}(\Delta^+ f) \subset \cup \lambda\} \quad \text{and} \quad (1.23)$$

$$E_\lambda[a, b] := \{g \in Q_\lambda[a, b]: g(a) = 0, \quad 1 + \Delta^- g \wedge \Delta^+ g \gg 0 \quad \text{and} \quad N_{(a,b)}(\Delta^+ g) \subset \cup \lambda\}. \quad (1.24)$$

We consider a regulated evolution U such that $U_a \in L_\lambda[a, b]$ for some $\lambda \in \Lambda[a, b]$. To show that U has the λ -generator and to give its representation, we use the Left Cauchy λ -integral and the product λ -integral. We say that the function $\mathcal{L}_\lambda f$ is defined on $[a, b]$ if the Left Cauchy λ -integral $(LC) \int f^{-1} d_\lambda f$ is defined on $[a, b]$, and then let $\mathcal{L}_\lambda f(t) := (LC) \int_a^t f^{-1} d_\lambda f$ for $t \in [a, b]$.

Theorem 5 *Let $\lambda \in \Lambda[a, b]$, and let U be a regulated evolution in $\mathbb{R}$ such that $U_a \equiv U(\cdot, a) \in L_\lambda[a, b]$. Then statements (a), (b) and (c) hold, where:*

- (a) *the function $\mathcal{L}_\lambda U_a$ is defined on $[a, b]$, it is in $E_\lambda[a, b]$ and it is the λ -generator of U ;*
- (b) *the product λ -integral $\mathfrak{Y}(1 + d_\lambda(\mathcal{L}_\lambda U_a))$ is defined on $[a, b]$;*
- (c) *the function $\mathcal{L}_\lambda U_a$ in $E_\lambda[a, b]$ is unique up to an additive constant such that*

$$U(t, s) = \mathfrak{Y}_s^t(1 + d_\lambda(\mathcal{L}_\lambda U_a)) \quad a \leq s \leq t \leq b. \quad (1.25)$$

Proof. Statement (a) holds by Proposition 3.56 and Theorem 3.57. Statement (b) holds by Theorem 3.58. By (3.169) of Theorem 3.58, we have that for all $a \leq s \leq t \leq b$,

$$\mathfrak{Y}_s^t(1 + d_\lambda(\mathcal{L}_\lambda U_a)) = U_a(t)/U_a(s) = U(t, s),$$

and so (1.25) holds. Suppose that $g \in E_\lambda[a, b]$ is such that $\mathfrak{Y}_a^t(1 + d_\lambda g) = U(t, a)$ for $t \in [a, b]$. Then $\mathcal{L}_\lambda U_a = g + C$ for some constant C by Theorem 3.59, proving uniqueness of the representation, and hence statement (c). The proof of Theorem 5 is complete. $\square$

In Chapter 5, the solution of the evolution representation problem is used to build up an asset pricing model. More specifically, stock price changes during a time period $[0, T]$ are considered as an evolution U such that $U_0 \equiv U(\cdot, 0) \in L_\lambda[0, T]$. By Theorem 5, the evolution U has the λ -generator which is used to model a return of a stock in our asset pricing model. In this chapter there are two new results: an almost sure approximation of a continuous time Black-Scholes model by a discrete time binomial model, and the option pricing formula when a stock price is modelled by a function having the quadratic λ -variation. The second result clarifies and solves the problem posed in Bick and Willinger [7].

Chapter 2

The p -variation and integrals of Stieltjes type

One thing at least we may say: no theory of integration is complete which does not enable us to define, by some kind of integration process, the integral of a continuous function with respect to itself, or, more generally, the integration of an ordinary function, such as the square, of a continuous function with respect to that continuous function, when the latter has not bounded variation.

W. H. Young. “The progress of mathematical analysis in the twentieth century” (Presidential Address). Proc. London Math. Soc. Ser. 2, Vol. 24 (1925), 421-434.

The problem of integration of functions having unbounded variation was approached in several different ways. Although only very few of these approaches have reached a development to a full fledged calculus. Integration theories dealing with suitable families of functions, such as stochastic processes or supports of probability distributions, are the most advanced ones. That is not surprising because of Stochastic Analysis ability to model uncertainty in natural and social sciences. The classical approach to integration of a single function having unbounded variation received less attention partly because there were no immediate applications, at least in other areas of analysis. We believe that further developments of the classical approach to integration have potentials to shed a new light on modelling problems of a real life.

2.1 Additive and multiplicative functions

It is well-known that the Lebesgue-Stieltjes integral and the Riemann-Stieltjes integral are both additive, but in a different sense. Namely, let ν be a function defined on the class of all subintervals, closed or open at either end, of an interval $[a, b]$. The function ν is additive over disjoint subintervals if $\nu(A \cup B) = \nu(A) + \nu(B)$ for any disjoint subintervals A, B of $[a, b]$ such that $A \cup B$ is a subinterval. In particular, for any $a \leq s < t < r \leq b$, we have

$$\nu([s, r]) = \nu([s, t]) + \nu([t, r]) = \nu([s, t]) + \nu((t, r]).$$

This is the additivity property enjoyed by the Lebesgue-Stieltjes integral. On the other hand, the Riemann-Stieltjes integral may be considered as a function μ defined on the simplex

$$S[a, b] := \{(s, t) : s, t \in [a, b], \quad s \leq t\}$$

of an interval $[a, b]$, and with this notation the integral possess the additivity property: for any $a \leq s \leq t \leq r \leq b$,

$$\mu(s, r) = \mu(s, t) + \mu(t, r).$$

As compared to the additivity over disjoint subintervals, the latter property amounts to the additivity over adjacent closed subintervals. The latter property also holds for all extended Stieltjes type integrals discussed in this paper, and so the resulting integration theories differ from the theory of integration with respect to a measure. Notice that an evolution U defined by (1.9), is just a multiplicative function when it is viewed as a function defined on the simplex $S[a, b]$.

In this paper we consider functions defined on the simplex of a lineary ordered set, which are additive or multiplicative. This approach provides a concise formulation of various properties related to a sequence λ of partitions of an interval $[a, b]$. For example, conditions (1.1) and (1.2) defining the quadratic λ -variation will become equivalent to a single property of an additive upper continuous function defined on the simplex of an extended interval to be defined next.

Regulated functions and extended intervals. Let J be an interval of a real line, which may be bounded or unbounded, and open or closed at either end. A real-valued function f defined on J is called *regulated on J* if the right limit $f(x+) := \lim_{y \downarrow x} f(y)$ exists and is finite for $x \in \bar{J}$ not equal to the right endpoint of J , and if the left limit $f(x-) := \lim_{y \uparrow x} f(y)$ exists and is finite for $x \in \bar{J}$ not equal to the left endpoint of J . See Section 2 in [23, Part III] for more information about regulated functions. The set of all regulated function on J will be denoted by $\mathcal{R}(J)$.

For a regulated function f on J , define a function $\Delta^+ f$ on J by $(\Delta^+ f)(x) := f(x+) - f(x)$ for all $x \in J$ except at the right endpoint, where it is defined to be zero. Similarly, define a function $\Delta^- f$ on J by $(\Delta^- f)(x) := f(x) - f(x-)$ for all $x \in J$ except at the left endpoint, where it is defined to be zero. Also, let $\Delta^\pm f := \Delta^+ f + \Delta^- f$ on J .

With each point x of a real line one can adjoin two symbols $x-$ and $x+$. Then a set formed by an interval with adjoined symbols extends naturally the domain of a regulated function. More formally, given a closed subinterval $[a, b]$ of an interval $J \subset \mathbb{R}$, define the *extended interval* $\llbracket a, b \rrbracket$ by

$$\llbracket a, b \rrbracket := (\{a\} \times \{0, +\}) \cup ((a, b) \times \{-, 0, +\}) \cup (\{b\} \times \{-, 0\}) \quad (2.1)$$

if $a < b$, and $\llbracket a, b \rrbracket := \emptyset$ if $a = b$. For $x \in [a, b]$, we identify $(x, -)$ with $x-$, $(x, 0)$ with x and $(x, +)$ with $x+$. With this identification, we have that an ordinary interval $[a, b]$ is a subset of the extended interval $\llbracket a, b \rrbracket$. Letting $z+ < x- < x < x+ < y-$ for all $z < x < y$ with $z, y \in [a, b]$, gives a linear ordering of points of $\llbracket a, b \rrbracket$. If f is a regulated function on $[a, b]$ then its natural extension on $\llbracket a, b \rrbracket$ again will be denoted by f . On $\llbracket a, b \rrbracket$ we have the topology induced by the linear ordering with a base given by the class of all sets $\{u: v < u < w\}$, $\{u: a \leq u < w\}$ and $\{u: v < u \leq b\}$ for $v, w \in \llbracket a, b \rrbracket$. The extended interval $\llbracket a, b \rrbracket$ is compact, not metrizable and is totally disconnected. The set $\mathcal{R}[a, b]$ can be identified with the set of all continuous real-valued functions on $\llbracket a, b \rrbracket$. This identification also has a Banach algebra aspect (see Theorem 5 of Berberian [4]).

Let $[a, b]$ be a closed interval with endpoints $a < b$. The *simplex of the extended interval* $\llbracket a, b \rrbracket$ is defined by

$$S\llbracket a, b \rrbracket := \{(u, v): u, v \in \llbracket a, b \rrbracket, \quad u \leq v\}.$$

In view of the above mentioned identification of $x \in [a, b]$ with $(x, 0)$ in (2.1), we can and do assume that the simplex $S[a, b]$ of the ordinary interval $[a, b]$ is a subset of the simplex $S[[a, b]]$ of the extended interval $[[a, b]]$. For $(u, v) \in S[[a, b]]$, let

$$[[u, v]] := \begin{cases} \{w \in [[a, b]] : u \leq w \leq v\} & \text{if } u \neq v, \\ \emptyset & \text{if } u = v. \end{cases}$$

This definition agree with (2.1) if $u, v \in [a, b]$ are ordinary points. In that case, that is, for ordinary points $(s, t) \in S[a, b]$, $[[s, t]]$ may be called the extended closed interval. Also, for ordinary points $(s, t) \in S[a, b]$, $s \neq t$, the extended open interval is defined by $[[s+, t-]]$, the extended left-open interval is defined by $[[s+, t]]$, and the extended right-open interval is defined by $[[s, t-]]$.

Functions on $S[[a, b]]$. Let $[a, b]$ be a closed interval with endpoints $a < b$. Any function μ defined on the simplex $S[[a, b]]$ of the extended interval $[[a, b]]$ is called a function on $S[[a, b]]$. Next we define additive, multiplicative and upper continuous functions on $S[[a, b]]$.

Definition 1 A function μ on $S[[a, b]]$ is called *additive* if

$$\begin{cases} \mu(u, v) = \mu(u, w) + \mu(w, v) & \text{for each } u, w, v \in [[a, b]] \text{ such that } u \leq w \leq v, \\ \mu(u, u) = 0 & \text{for each } u \in [[a, b]]. \end{cases}$$

Here the second condition follows from the first one. To give an example of an additive function on $S[[a, b]]$, let f be a regulated function on $[a, b]$. Define the function $\mu(f)$ on $S[[a, b]]$ by

$$\mu(f; u, v) := f(v) - f(u), \quad \text{for } (u, v) \in S[[a, b]]. \quad (2.2)$$

Clearly, $\mu(f)$ is additive function on $S[[a, b]]$ such that $\mu(f; t-, t) = \Delta^- f(t)$ for $t \in (a, b]$ and $\mu(f; s, s+) = \Delta^+ f(s)$ for $s \in [a, b)$. Other examples of additive functions on $S[[a, b]]$ will be given later on. As we already mentioned, the quadratic λ -variation is defined by means of a suitable additive function on $S[[a, b]]$ (see Definition 3 below).

Definition 2 A function μ on $S[[a, b]]$ is called *multiplicative* if

$$\begin{cases} \mu(u, v) = \mu(u, w)\mu(w, v) & \text{for each } u, w, v \in [[a, b]] \text{ such that } u \leq w \leq v, \\ \mu(u, u) = 1 & \text{for each } u \in [[a, b]]. \end{cases}$$

Here the second condition follows from the first one if $\mu(a, u) \neq 0$ for each $u \in [[a, b]]$. To give an example of a multiplicative function on $S[[a, b]]$, let f be a regulated function on $[a, b]$ such that $1/f$ is bounded on $[a, b]$. Define the function $\pi(f)$ on $S[[a, b]]$ by

$$\pi(f; u, v) := f(v)/f(u), \quad \text{for } (u, v) \in S[[a, b]]. \quad (2.3)$$

Clearly, $\pi(f)$ is multiplicative function on $S[[a, b]]$.

For extended intervals as for other sets, $[[u_k, v_k]] \uparrow [[u, v]]$ means that $u_1 \geq u_2 \geq \dots$, $v_1 \leq v_2 \leq \dots$ and $\cup_{k=1}^{\infty} [[u_k, v_k]] = [[u, v]]$, while $[[u_k, v_k]] \downarrow [[u, v]]$ means that $u_1 \leq u_2 \leq \dots$, $v_1 \geq v_2 \geq \dots$ and $\cap_{k=1}^{\infty} [[u_k, v_k]] = [[u, v]]$.

Definition 3 A function μ on $S[[a, b]]$ is called *upper continuous* if $\mu(u_k, v_k) \rightarrow \mu(u, v)$ for any sequence $(u, v), (u_1, v_1), (u_2, v_2), \dots \in S[[a, b]]$ such that $[[u_k, v_k]] \downarrow [[u, v]]$. If the same is true whenever $u = v$ then a function μ on $S[[a, b]]$ is called *upper continuous at $\emptyset$* (empty set).

For a function μ on $S[[a, b]]$, define the right distribution function R_μ and the left distribution function L_μ on $[[a, b]]$ respectively by

$$R_\mu(u) := \mu(a, u), \quad u \in [[a, b]] \quad \text{and} \quad L_\mu(u) := \mu(u, b), \quad u \in [[a, b]]. \quad (2.4)$$

By definition of the interval topology, a real-valued function f on $[[a, b]]$ is *continuous at $u \in [[a, b]]$* , which is not an endpoint, if for each $\epsilon > 0$ there are $u_1, u_2 \in [[a, b]]$ such that $u_1 < u < u_2$ and $|f(u) - f(v)| < \epsilon$ for each $v \in [[a, b]]$, $u_1 < v < u_2$. Continuity at an endpoint means the one-sided variant of the continuity just formulated. A function f is *continuous on $[[a, b]]$* if it is continuous at each $u \in [[a, b]]$. Any function f defined on $[[a, b]]$ is continuous at an ordinary point $t \in [a, b] \subset [[a, b]]$ because always one can take $u_1 := t-$ if $t > a$ and $u_2 := t+$ if $t < b$. Thus the continuity on $[[a, b]]$ is a restriction of a function on its behaviour on the left side of each point $t-$ and on the right side of each point $t+$.

Let μ be a function on an extended interval $[[a, b]]$. We say that μ is *nondegenerate* if $\mu(a, b) \neq 0$, and μ is *bounded* if $\|\mu\|_\infty := \sup\{|\mu(u, v)| : (u, v) \in S[[a, b]]\} < \infty$.

Theorem 4 Let a function μ on $S[[a, b]]$ be either additive or multiplicative, nondegenerate and bounded. Then the following statements are equivalent:

- (i) μ is upper continuous;
- (ii) μ is upper continuous at $\emptyset$;
- (iii) R_μ is continuous on $[[a, b]]$;
- (iv) when restricted to $[a, b]$, R_μ is regulated on $[a, b]$, $\lim_{s \uparrow t} R_\mu(s) = \mu(a, t-)$ for $t \in (a, b]$ and $\lim_{s \downarrow t} R_\mu(s) = \mu(a, t+)$ for $t \in [a, b)$;
- (v) L_μ is continuous on $[[a, b]]$;
- (vi) when restricted to $[a, b]$, L_μ is regulated on $[a, b]$, $\lim_{s \uparrow t} L_\mu(s) = \mu(t-, b)$ for $t \in (a, b]$ and $\lim_{s \downarrow t} L_\mu(s) = \mu(t+, b)$ for $t \in [a, b)$;
- (vii) $\mu(s_k+, t_k-) \rightarrow \mu(u, u)$ whenever extended open intervals $[[s_k+, t_k-]] \downarrow [[u, u]]$ for some $u \in [[a, b]]$, and

$$\forall \epsilon > 0, \quad \begin{cases} \text{card } \{t \in (a, b]: |\mu(t-, t) - \mu(t, t)| > \epsilon\} < \infty, \\ \text{card } \{t \in [a, b): |\mu(t, t+) - \mu(t, t)| > \epsilon\} < \infty. \end{cases} \quad (2.5)$$

Proof. (i) $\Leftrightarrow$ (ii). Clearly, (i) implies (ii). For the converse implication, let extended intervals $A_k := [[u_k, v_k]] \downarrow [[u, v]] =: A$. Then $A_k = B_k \cup A \cup C_k$ for extended intervals B_k, C_k such that $B_k \downarrow [[u, u]]$ and $C_k \downarrow [[v, v]]$. Thus $\mu(u_k, v_k) \rightarrow \mu(u, v)$ by additivity or multiplicativity.

(ii) $\Rightarrow$ (iii). Let $u \in [[a, b]]$. Due to above remarks following the definition of continuity on extended interval and by symmetry, one can assume that $u = t-$ for some $t \in (a, b]$. Let $\epsilon > 0$. Since $[[v, t-]] \downarrow [[t-, t-]]$ as $v \uparrow t-$, by the assumption there exists $a \leq u_1 < t-$ such that $|\mu(v, t-) - \mu(t-, t-)| < \epsilon$ for each $u_1 < v < t-$. Otherwise we could extract a

sequence $\llbracket v_k, t- \rrbracket \downarrow \llbracket t-, t- \rrbracket$ such that $\mu(v_k, t-) \not\rightarrow \mu(t-, t-)$. Let $u_2 := t$. If μ is additive then $|R_\mu(t-) - R_\mu(v)| = |\mu(v, t-)| < \epsilon$ for each $u_1 < v < u_2$. If μ is multiplicative then

$$|R_\mu(t-) - R_\mu(v)| = |\mu(a, v)| |\mu(v, t-) - 1| < \epsilon \|\mu\|_\infty$$

for each $u_1 < v < u_2$. Since $\epsilon > 0$ is arbitrary, R_μ is continuous at $t-$, and so R_μ is continuous on $\llbracket a, b \rrbracket$.

(iii) $\Rightarrow$ (iv). For $t \in (a, b]$, since R_μ is continuous at $t-$ and $R_\mu(t-) = \mu(a, t-)$, it follows that $\lim_{s \uparrow t} R_\mu(s) = \mu(a, t-)$. Similarly it follows that $\lim_{s \downarrow t} R_\mu(s) = \mu(a, t+)$ for $t \in [a, b)$. Thus R_μ is regulated when restricted to $[a, b]$.

(iv) $\Rightarrow$ (vii). Let $\bar{R}_\mu$ be the restriction of R_μ to $[a, b]$. Since $\bar{R}_\mu$ is regulated, $\bar{R}_\mu(t-)$ is defined and equals to $\mu(a, t-)$ for each $t \in (a, b]$ by the assumption. Similarly, $\bar{R}_\mu(t+) = \mu(a, t+)$ for each $t \in [a, b)$. First suppose that μ is additive. Since $\mu(t-, t) = \Delta^- \bar{R}_\mu(t)$, $\mu(t, t+) = \Delta^+ \bar{R}_\mu(t)$ and $\mu(t, t) = 0$, (2.5) holds by the analogous property for regulated functions. Let open extended intervals $\llbracket s_k+, t_k- \rrbracket \downarrow \emptyset$ as $k \rightarrow \infty$, where each $(s_k, t_k) \in S[a, b]$. Then either for some $t \in (a, b]$ and all sufficiently large k , $t_k = t$ and $s_k \uparrow t$, or for some $s \in [a, b)$ and all sufficiently large k , $s_k = s$ and $t_k \downarrow s$. Since μ is additive and $\bar{R}_\mu$ is regulated, $\mu(s_k+, t_k-) = \bar{R}_\mu(t-) - \bar{R}_\mu(s_k+) \rightarrow 0$ as $k \rightarrow \infty$ in the first case, and $\mu(s_k, t_k-) = \bar{R}_\mu(t_k-) - \bar{R}_\mu(s_k+) \rightarrow 0$ as $k \rightarrow \infty$ in the second case. Thus the statement (vii) holds when μ is additive. Now suppose that μ is multiplicative, and hence nondegenerate and bounded. For $t \in (a, b]$, we have

$$\bar{R}_\mu(t) - \bar{R}_\mu(t-) = \bar{R}_\mu(t-) \{\mu(t-, t) - 1\}. \quad (2.6)$$

We claim that $C := \inf_t |\bar{R}_\mu(t-)| > 0$. Suppose not. Then there exists an infinite sequence $s_j \rightarrow s \in [a, b]$ such that either $\bar{R}_\mu(s_j-) \rightarrow \bar{R}_\mu(s-) = 0$, or $\bar{R}_\mu(s_j-) \rightarrow \bar{R}_\mu(s+) = 0$. In either case due to multiplicativity of μ , it follows that $\mu(a, b) = 0$. The contradiction proves the claim. Then by (2.6), $|\Delta^- \bar{R}_\mu(t)| \geq C |\mu(t-, t) - 1|$ for $t \in (a, b]$. Similarly, one can conclude that $|\Delta^+ \bar{R}_\mu(t)| \geq C |\mu(t, t+) - 1|$ for $t \in [a, b)$ and some $C > 0$. Thus (2.5) holds by the analogous property for regulated functions. For $(s_k, t_k) \in S[a, b]$, since

$$\bar{R}_\mu(t_k-) - \bar{R}_\mu(s_k+) = \bar{R}_\mu(s_k+) \{\mu(s_k+, t_k-) - 1\},$$

one can conclude as in the additive case that $\mu(s_k+, t_k-) \rightarrow 1$ if $(s_k, t_k) \downarrow \emptyset$. Thus the statement (vii) holds when μ is multiplicative.

(vii) $\Rightarrow$ (ii). Let extended intervals $\llbracket u_k, v_k \rrbracket \downarrow \llbracket w, w \rrbracket$ for some $w \in \llbracket a, b \rrbracket$. Then either for some $s \in (a, b]$, $v_k = s- = w$ for all sufficiently large k , or for some $t \in [a, b)$, $u_k = t+ = w$ for all sufficiently large k . Using additivity if μ is additive, or continuity of the multiplication if μ is multiplicative, in each of the two cases $\mu(u_k, v_k) \rightarrow \mu(w, w)$ follows by the statement (vii).

The proof of implications (ii) $\Rightarrow$ (v) $\Rightarrow$ (vi) $\Rightarrow$ (vii) $\Rightarrow$ (ii) is based on symmetric arguments which we omit. The proof of Theorem 4 is complete. $\square$

Let f be a regulated function on $[a, b]$. Since the right distribution function $R_{\mu(f)} = f - f(a)$ of the additive function $\mu(f)$ on $\llbracket a, b \rrbracket$ defined by (2.2), satisfies statement (iv) of the preceding theorem, the function $\mu(f)$ is upper continuous. Similarly, the multiplicative function $\pi(f)$ defined by (2.3) is upper continuous, if in addition, f is bounded away from zero. In the sense of the following two statements, these two examples are canonical. Let $\mathcal{AC}\llbracket a, b \rrbracket$ be the set of all additive upper continuous functions on $S\llbracket a, b \rrbracket$, and let $R_{\mu, [a, b]}$ denotes the right distribution function R_μ restricted to $[a, b]$.

Corollary 5 *Let $a < b$. Then one-to-one linear operators between the vector spaces $\mathcal{AC}[[a, b]]$ and $\{f \in \mathcal{R}[a, b] : f(a) = 0\}$ are given by $f = R_{\mu, [a, b]}$ and $\mu = \mu(f)$.*

Proof. For $\mu \in \mathcal{AC}[[a, b]]$, let $f := R_{\mu, [a, b]}$. By statement (iv) of Theorem 4, $f \in \mathcal{R}[a, b]$ and $f(a) = \mu(a, a) = 0$. Conversely, for $f \in \mathcal{R}[a, b]$ such that $f(a) = 0$, let $\mu := \mu(f)$ be defined by (2.2). Since $R_{\mu(f), [a, b]} = f$, μ is upper continuous by the implication (iv) $\Rightarrow$ (i) of Theorem 4, and so $\mu \in \mathcal{AC}[[a, b]]$. Clearly, the two mappings are linear and one-to-one, proving the conclusion. $\square$

For a function f on $[a, b]$, we write $|f| \gg 0$ if there is a constant $C > 0$ such that $|f(t)| \geq C$ for all $t \in [a, b]$. Let $\mathcal{MC}[[a, b]]$ be the set of all nondegenerate, bounded, multiplicative and upper continuous functions defined on $S[[a, b]]$.

Corollary 6 *Let $a < b$. Then one-to-one mappings between the sets $\mathcal{MC}[[a, b]]$ and $\{f \in \mathcal{R}[a, b] : f(a) = 1 \text{ and } |f| \gg 0\}$ are given by $f = R_{\mu, [a, b]}$ and $\mu = \pi(f)$.*

Proof. For $\mu \in \mathcal{MC}[[a, b]]$, let $f := R_{\mu, [a, b]}$. By statement (iv) of Theorem 4, $f \in \mathcal{R}[a, b]$ and $f(a) = \mu(a, a) = 1$. Letting $C := \inf\{|\mu(a, t)| : t \in [a, b]\}$, we have that $C > 0$. If not, due to compactness of $[a, b]$, there is $u \in [a, b]$ such that $\mu(a, u) = 0$. Then $\mu(a, b) = \mu(a, u)\mu(u, b) = 0$, a contradiction proving that $|f(t)| = |\mu(a, t)| \geq C > 0$ for each $t \in [a, b]$. Conversely, for $f \in \mathcal{R}[a, b]$ such that $f(a) = 1$ and $|f| \gg 0$, let $\mu := \pi(f)$ be defined by (2.3). Then μ is nondegenerate, bounded and multiplicative function defined on $S[[a, b]]$. Since $R_{\mu(f), [a, b]} = f$, μ is upper continuous by the implication (iv) $\Rightarrow$ (i) of Theorem 4, and so $\mu \in \mathcal{MC}[[a, b]]$, proving the conclusion. $\square$

2.2 The Wiener class

Recall that $\Xi[a, b]$ denotes the set of all partitions of a closed interval of real numbers $[a, b]$. For $\kappa, \lambda \in \Xi[a, b]$, the relation $\kappa \subset \lambda$ means that the partition λ is a refinement of the partition κ . Let $\mathfrak{P}$ be the family of all sets $P(\kappa) := \{\lambda \in \Xi[a, b] : \lambda \supset \kappa\}$, $\kappa \in \Xi[a, b]$. Then $\mathfrak{P}$ is the direction in the sense of Definition A.1, which we call the *direction of partitions* of $[a, b]$. Recall also that $\mathcal{W}_p = \mathcal{W}_p[a, b]$, $0 < p < \infty$, denotes the set of all real-valued functions f on $[a, b]$ having bounded p -variation, that is, (1.5) holds. For a function f on $[a, b]$ and for $0 < p < \infty$, let

$$v_p^*(f) = v_p^*(f; [a, b]) := \limsup_{\kappa, \mathfrak{P}} s_p(f; \kappa) := \inf_{\kappa} \sup\{s_p(f; \lambda) : \lambda \in \Xi[a, b], \lambda \supset \kappa\},$$

$$\sigma_p^*(f) = \sigma_p^*(f; [a, b]) := \liminf_{\kappa, \mathfrak{P}} s_p(f; \kappa) := \sup_{\kappa} \inf\{s_p(f; \lambda) : \lambda \in \Xi[a, b], \lambda \supset \kappa\}.$$

Notation $s_p(f; \kappa)$ is defined by (1.3). For a regulated function f on $[a, b]$ and for $0 < p < \infty$, let

$$\sigma_p(f) = \sigma_p(f; [a, b]) := \sum_{[a, b]} |\Delta f|^p := \lim_{\sigma, \mathfrak{S}} \sum_{\sigma} |\Delta f|^p, \quad (2.7)$$

where the limit is defined in Definition A.5, and it always exists finite or infinite. Here and below, for a finite set $\sigma \subset [a, b]$,

$$\sum_{\sigma} |\Delta f|^p := \sum_{x \in \sigma} \left\{ |\Delta_a^- f(x)|^p + |\Delta_b^+ f(x)|^p \right\},$$

where $\Delta_a^- f(x) = f(x) - f(x-)$ if $x \in (a, b]$, $\Delta_a^- f(a) := 0$, and $\Delta_b^+ f(x) = f(x+) - f(x)$ if $x \in [a, b)$, $\Delta_b^+ f(b) := 0$. Then the following relations hold for the four quantities just defined.

Lemma 7 *Let f be a regulated function on $[a, b]$, and let $0 < p < \infty$. Then*

$$\sigma_p(f) \leq \sigma_p^*(f) \leq v_p^*(f) \leq v_p(f). \quad (2.8)$$

Proof. Suppose that $\sigma_p(f) < +\infty$. Then for each finite $\sigma \subset [a, b]$ and each $\epsilon > 0$, there exists a partition κ of $[a, b]$ such that $\sigma \subset \kappa$ and for each refinement λ of κ , $s_p(f; \lambda) \geq \sum_{\sigma} |\Delta f|^p - \epsilon$, and so $\sigma_p(f) \leq \sigma_p^*(f)$. If $\sigma_p(f) = +\infty$, then similarly it follows that $\sigma_p^*(f) = +\infty$. Therefore the first inequality in (2.8) holds. To prove the second inequality, one can assume that $v_p^*(f)$ is finite. For each $\kappa \in \Xi[a, b]$, let

$$I(\kappa) := \inf\{s_p(f; \lambda) : \lambda \in \Xi[a, b], \lambda \supset \kappa\} \quad \text{and} \quad S(\kappa) := \sup\{s_p(f; \lambda) : \lambda \in \Xi[a, b], \lambda \supset \kappa\}.$$

Suppose that $v_p^*(f) = \inf_{\kappa} S(\kappa) < \sigma_p^*(f) = \sup_{\kappa} I(\kappa)$. Then there exist $\kappa_1, \kappa_2 \in \Xi[a, b]$ such that $S(\kappa_1) < I(\kappa_2)$. Let $\kappa_3 := \kappa_1 \cup \kappa_2$. Since $\kappa_3 \supset \kappa_1$ and $\kappa_3 \supset \kappa_2$, it follows that

$$S(\kappa_3) \leq S(\kappa_1) < I(\kappa_2) \leq I(\kappa_3).$$

This contradiction proves the second inequality in (2.8). Since the third inequality is obvious, the proof of Lemma 7 is complete. $\square$

By Proposition 2.12 in [23, Part II], for a regulated function f on $[a, b]$ and for $0 < p < 1$, we have

$$\sigma_p(f) = \sigma_p^*(f) = v_p^*(f) = v_p(f).$$

In fact, each function f which has the p -variation on $[a, b]$ bounded for some $0 < p < 1$, is a pure jump function on $[a, b]$ (see Theorem 2.11 in [23, Part II]). Recall that a regulated function f on $[a, b]$ is called a *pure jump* function, if for each $a < x \leq b$, the directed function $\sum_{\mu \cap (a, x)} \{\Delta^- f + \Delta^+ f\}$ has a limit $\sum_{(a, x)} \{\Delta^- f + \Delta^+ f\}$, which satisfies the relation

$$f(x) = f(a) + \Delta^+ f(a) + \sum_{(a, x)} \{\Delta^- f + \Delta^+ f\} + \Delta^- f(x).$$

In the case $p = 1$, by Proposition 2.13 in [23, Part II], $v_1^*(f) = v_1(f)$ for any function f on $[a, b]$, while $\sigma_1(f) = v_1^*(f)$ if and only if f is a pure jump function. The situation again becomes different when $1 < p < \infty$. By Theorem 8 to be proved next, for a regulated function f , $\sigma_p^*(f) = \sigma_p(f)$ provided $1 < p < \infty$.

It appears that functions $f \in \mathcal{W}_p$, $1 < p < \infty$, such that $\sigma_p^*(f) = v_p^*(f)$, have especially nice properties. Define the *Wiener class* $\mathcal{W}_p^*$ by

$$\mathcal{W}_p^* := \mathcal{W}_p^*[a, b] := \{f \in \mathcal{W}_p[a, b] : \sigma_p^*(f) = v_p^*(f)\}.$$

By Theorem 11.4 of McShane and Botts [81, p. 55], $f \in \mathcal{W}_p^*$ if and only if the directed function $(s_p(f; \cdot), \mathfrak{P})$ has a limit $\lim_{\kappa, \mathfrak{P}} s_p(f; \kappa)$. Thus recalling (1.4), functions having the local p -variation constitute the Wiener class $\mathcal{W}_p^*$. This in conjunction with Theorem 8, yields that for a regulated function f and $1 < p < \infty$, $f \in \mathcal{W}_p^*$ if and only if the directed function $(s_p(f; \cdot), \mathfrak{P})$ converges and

$$\lim_{\kappa, \mathfrak{P}} s_p(f; \kappa) = \sigma_p(f).$$

Love and Young [68] defined the Wiener class $\mathcal{W}_p^*$ to be the set of all functions $f \in \mathcal{W}_p$ for which $v_p^*(f) = \sigma_p(f)$. Due to the following Theorem 8, the original and the present definitions of $\mathcal{W}_p^*$ give the same class of functions. Love and Young [68, §12, p. 32] showed that, for $1 < p < \infty$,

$$\cup\{\mathcal{W}_q : 1 \leq q < p\} \subset \mathcal{W}_p^* \subset \mathcal{W}_p.$$

Theorem 8 *Let f be a regulated function on $[a, b]$, and let $1 < p < \infty$. Then $\sigma_p^*(f) = \sigma_p(f)$.*

Proof. If $\sigma_p(f) = +\infty$ then the conclusion holds by Lemma 7. Suppose that $\sigma_p(f) < +\infty$. Let $\epsilon \in (0, 1)$. It is enough to show that for a given partition κ , there exists a refinement λ of κ such that

$$s_p(f; \lambda) \leq \sigma_p(f) + \epsilon. \quad (2.9)$$

Let κ be a partition of $[a, b]$, and let $K := (4^p/2)(1 + \text{Osc}(f))$. Choose a partition $\mu = \{y_j: j = 0, \dots, m\}$ of $[a, b]$ such that $\kappa \subset \mu$ and

$$\sum_{\mu} |\Delta f|^p \geq \sigma_p(f) - \epsilon/(3K). \quad (2.10)$$

Then choose a set $\{u_{j-1}, v_j: j = 1, \dots, m\} \subset [a, b]$ such that $y_{j-1} < u_{j-1} < v_j < y_j$ for $j = 1, \dots, m$ and

$$\left| \sum_{\mu} |\Delta f|^p - \sum_{j=1}^m \left\{ |f(u_{j-1}) - f(y_{j-1})|^p + |f(y_j) - f(v_j)|^p \right\} \right| < \epsilon/3. \quad (2.11)$$

Also, without loss of generality we can assume that f is continuous at each of the points $\{u_{j-1}, v_j: j = 1, \dots, m\}$. We claim that for each $j = 1, \dots, m$ there exists a partition λ_j of $[u_{j-1}, v_j]$ such that

$$s_p(f; \lambda_j) \leq K\sigma_p(f; [u_{j-1}, v_j]) + \epsilon/(3m). \quad (2.12)$$

Assuming that this is true, and letting $\lambda := \mu \cup (\cup_{j=1}^m \lambda_j)$, it then follows that (2.9) holds by (2.10), (2.11) and (2.12), as desired.

To prove the claim, let $j \in \{1, \dots, m\}$, and suppose that $\sigma_j := \sigma_p(f; [u_{j-1}, v_j]) > 0$. If $|f(v_j) - f(u_{j-1})|^p \leq K\sigma_j$ then taking $\lambda_j = \{u_{j-1}, v_j\}$, (2.12) holds. Otherwise, either $f(v_j) - f(u_{j-1}) > (K\sigma_j)^{1/p}$ or $f(v_j) - f(u_{j-1}) < -(K\sigma_j)^{1/p}$. We construct λ_j only in the first case because the second case is symmetric. It is enough to construct a partition λ_j^+ of $[u_{j-1}, v_j]$ having partition points t_+ with $t \in (u_{j-1}, v_j)$ and such that

$$s_p(f; \lambda_j^+) \leq K\sigma_j = K\sigma_p(f; [u_{j-1}, v_j]). \quad (2.13)$$

Approximating $f(t_+)$ by values of f at ordinary points, we get a partition λ_j such that $s_p(f; \lambda_j)$ is arbitrary close to $s_p(f; \lambda_j^+)$, and hence (2.12) follows. To begin the construction let $d_j := f(v_j) - f(u_{j-1})$. Since $d_j \leq \text{Osc}(f) < K < d_j^p/\sigma_j$, we have $0 < \sigma_j^{1/(p-1)} < d_j$. Choose the least integer $M \geq 2$ such that

$$d_j/M \leq \sigma_j^{1/(p-1)} < d_j. \quad (2.14)$$

If $f(t) - f(u_{j-1}) \leq d_j/M$ for each $t \in (u_{j-1}, v_j)$, then by continuity of f at v_j , $d_j \leq d_j/M$, a contradiction. Thus, there is a $t \in (u_{j-1}, v_j)$ such that $f(t) - f(u_{j-1}) > d_j/M$. Let

$$t_1 := \inf \{t \in (u_{j-1}, v_j): f(t) - f(u_{j-1}) > d_j/M\}.$$

Since f is continuous at u_{j-1} , $t_1 > u_{j-1}$. Then we have $f(t_1+) - f(u_{j-1}) \geq d_j/M$ and $f(t_1-) - f(u_{j-1}) \leq d_j/M$. Thus

$$d_j/M \leq f(t_1+) - f(u_{j-1}) \leq \Delta^\pm f(t_1) + d_j/M. \quad (2.15)$$

Since for each $t \in (u_{j-1}, v_j)$,

$$|\Delta^\pm f(t)| \leq 2^{p-1} [|\Delta^+ f(t)|^p + |\Delta^- f(t)|^p]^{1/p} \leq (K\sigma_j)^{1/p} < d_j.$$

we have $0 \leq \Delta^\pm f(t_1) \leq d_j$. Suppose that $d_j - d_j/M \leq f(t_1+) - f(u_{j-1}) \leq d_j + d_j/M$, that is,

$$-d_j/M \leq f(v_j) - f(t_1+) = d_j - [f(t_1+) - f(u_{j-1})] \leq d_j/M.$$

Letting $\lambda_j^+ := \{u_{j-1}, t_1+, v_j\}$, by (2.15) and (2.14), we then have

$$\begin{aligned} s_p(f; \lambda_j^+) &= |f(t_1+) - f(u_{j-1})|^p + |f(v_j) - f(t_1+)|^p \leq 2^{p-1} \{|\Delta^\pm f(t_1)|^p + 2(d_j/M)^p\} \\ &\leq 4^{p-1} \{\sigma_j + \text{Osc}(f)(d_j/M)^{p-1}\} \leq K\sigma_p(\mu; [u_{j-1}, v_j]), \end{aligned}$$

and so (2.13) holds. If $M = 2$ then the construction stops, proving the claim. Otherwise, we can assume that $M > 2$ and $f(t_1+) - f(u_{j-1}) < d_j - d_j/M$. Suppose that for some integer $2 \leq k < M$, we have constructed $u_{j-1} =: t_0 < t_1 < \dots < t_{k-1} < v_j$ such that $f(t_{k-1}+) - f(u_{j-1}) < d_j - d_j/M$ and for $i = 1, \dots, k-1$,

$$d_j/M \leq f(t_i+) - f(t_{i-1}+) \leq \Delta^\pm f(t_i) + d_j/M. \quad (2.16)$$

Thus $f(v_j) - f(t_{k-1}+) = d_j - [f(t_{k-1}+) - f(u_{j-1})] > d_j/M$. Let

$$t_k := \inf \{t \in (t_{k-1}, v_j) : f(t) - f(t_{k-1}+) > d_j/M\},$$

which exists since f is continuous at v_j , and $t_k > t_{k-1}$. As in (2.15), we have that (2.16) holds with $i = k$, and so,

$$f(t_k+) - f(u_{j-1}) = [f(t_{k-1}+) - f(u_{j-1})] + [f(t_k+) - f(t_{k-1}+)] < d_j + \Delta^\pm f(t_k).$$

Suppose that $f(t_k+) - f(u_{j-1}) \geq d_j - d_j/M$, and so,

$$-\Delta^\pm f(t_k) \leq f(v_j) - f(t_k+) = d_j - [f(t_k+) - f(u_{j-1})] \leq d_j/M.$$

Letting $\lambda_j^+ = \{t_i : i = 0, \dots, k+1\}$ with $t_{k+1} := v_j$ (recall that f is continuous at u_{j-1}, v_j), by (2.16) and (2.12), we then have

$$\begin{aligned} s_p(f; \lambda_j^+) &= \sum_{i=1}^{k+1} |f(t_i+) - f(t_{i-1}+)|^p \\ &\leq 2^{p-1} \left\{ \sum_{i=1}^k |\Delta^\pm f(t_i)|^p + |\Delta^\pm f(t_k)|^p + (k+1)(d_j/M)^p \right\} \\ &\leq (4^p/2) \{\sigma_j + \text{Osc}(f)(d_j/M)^{p-1}\} \leq K\sigma_p(f; [u_{j-1}, v_j]), \end{aligned}$$

so (2.13) holds. Otherwise, $f(t_k+) - f(u_{j-1}) < d_j - d_j/M$. On the other hand, by (2.16) with $i = 1, \dots, k$,

$$f(t_k+) - f(u_{j-1}) = \sum_{i=1}^k [f(t_i+) - f(t_{i-1}+)] \geq k(d_j/M),$$

which yields that $k \leq M - 1$. We continue construction which stops after at most M steps, proving the claim for all j such that $\sigma_j > 0$.

Let $j \in \{1, \dots, m\}$ be such that $\sigma_j = 0$, if such j exists. If $f(v_j) = f(u_{j-1})$ then (2.12) holds with $\lambda_j = \{u_{j-1}, v_j\}$. Otherwise, either $f(v_j) > f(u_{j-1})$ or $f(v_j) < f(u_{j-1})$. Again, we construct λ_j only in the first case because the second case is symmetric. Let $M \geq 2$ be the least integer such that $(1/M)^{p-1} \leq \epsilon/(3m\text{Osc}(f)^p)$. Let $t_0 := u_{j-1}$, and for $i = 1, \dots, M$, let

$$t_i := \inf\{t \in [t_{i-1}, v_j]: f(t) = f(u_{j-1}) + (i/M)[f(v_j) - f(u_{j-1})]\}.$$

Then taking $\lambda_j = \{t_i: i = 0, \dots, M\}$, we have

$$s_p(f; \lambda_j) = \sum_{i=1}^M |f(t_i) - f(t_{i-1})|^p = M^{1-p} |f(v_j) - f(u_{j-1})|^p < \epsilon/(3m),$$

so (2.12) holds in the case $\sigma_j = 0$. The proof of Theorem 8 is now complete. $\square$

The following characterization of the Wiener class is often used in what follows. We extend definition (1.5) of the p -variation over a closed interval $J = [a, b]$ to the case when J is open or closed at either end, by

$$v_p(f; J) := \sup \{s_p(f; \kappa): \kappa = \{x_i: i = 0, \dots, n\}, x_0, x_n \in J\}.$$

Lemma 9 *Let f be a regulated function on $[a, b]$, and let $1 < p < \infty$. Then the following statements are equivalent: (a) $f \in \mathcal{W}_p^*$;*

(b) for every $\epsilon > 0$, there is a partition $\{z_j: j = 0, \dots, m\}$ of $[a, b]$ such that

$$\sum_{j=1}^m v_p(f; (z_{j-1}, z_j)) < \epsilon; \quad (2.17)$$

(c) $\sigma_p(f) < \infty$, and for every $\epsilon > 0$, there is a partition λ of $[a, b]$ such that

$$\sum_{i=1}^n |f(x_{i-}) - f(x_{i-1}+)|^p < \epsilon \quad (2.18)$$

for each refinement $\{x_i: i = 0, \dots, n\}$ of λ .

Proof. (a) $\Rightarrow$ (b). Let $f \in \mathcal{W}_p^*$ and $\epsilon > 0$. By the definitions of $v_p^*(f)$ and $\sigma_p(f)$, there exists a partition $\kappa = \{z_j: j = 0, \dots, m\}$ of $[a, b]$ such that

$$\sum_{j=1}^m v_p(f; [z_{j-1}, z_j]) = \sup \{s_p(f; \lambda): \lambda \in \Xi[a, b], \lambda \supset \kappa\} < v_p^*(f) + \epsilon/2$$

$$\text{and} \quad \sum_{j=1}^m \{|\Delta^+ f(z_{j-1})|^p + |\Delta^- f(z_j)|^p\} > \sigma_p(f) - \epsilon/2.$$

Let $\{u_{j-1}, v_j: j = 1, \dots, m\}$ be a set of points in (a, b) such that $z_{j-1} < u_{j-1} < v_j < z_j$ for $j = 1, \dots, m$. Then we have, for $j = 1, \dots, m$,

$$v_p(f; [u_{j-1}, v_j]) \leq v_p(f; [z_{j-1}, z_j]) - \{v_p(f; [z_{j-1}, u_{j-1}]) + v_p(f; [v_j, z_j])\}.$$

For each $j = 1, \dots, m$, letting $u_{j-1} \downarrow z_{j-1}$ and $v_j \uparrow z_j$, by Lemma 2.19 of Dudley and Norvaiša [23, Part II], it follows that

$$\begin{aligned} \sum_{j=1}^m v_p(f; (z_{j-1}, z_j)) &\leq \sum_{j=1}^m \left\{ v_p(f; [z_{j-1}, z_j]) - |\Delta^+ f(z_{j-1})|^p - |\Delta^- f(z_j)|^p \right\} \\ &< v_p^*(f) + \epsilon/2 - \sigma_p(f) + \epsilon/2 = \epsilon. \end{aligned}$$

The last equality holds by the definition of the Wiener class $\mathcal{W}_p^*$ and by Theorem 8, proving statement (b).

(b) $\Rightarrow$ (a). Let statement (b) hold. Then it is easy to see that $f \in \mathcal{W}_p$. Assume however that $f \notin \mathcal{W}_p^*$. Therefore, since (2.8) always holds, $v_p^*(f) - \sigma_p^*(f) \geq C$ for some positive constant C . Let $\kappa = \{z_j: j = 0, \dots, m\} \in \Xi[a, b]$. Then we have

$$\sum_{j=1}^m \left\{ |\Delta^+ f(z_{j-1})|^p + v_p(f; (z_{j-1}, z_j)) + |\Delta^- f(z_j)|^p \right\} \geq v_p^*(f).$$

By Theorem 8, it then follows that

$$\begin{aligned} \sum_{j=1}^m v_p(f; (z_{j-1}, z_j)) &\geq v_p^*(f) - \sum_{j=1}^m \left\{ |\Delta^+ f(z_{j-1})|^p + |\Delta^- f(z_j)|^p \right\} \\ &\geq v_p^*(f) - \sigma_p(f) = v_p^*(f) - \sigma_p^*(f) \geq C > 0. \end{aligned}$$

Since κ is arbitrary, (2.17) can't hold for each $\epsilon > 0$. This contradiction implies that $f \in \mathcal{W}_p^*$.

(b) $\Rightarrow$ (c). Let statement (b) hold. Then it is easy to see that $f \in \mathcal{W}_p$, and hence $\sigma_p(f) < \infty$ by (2.8). Let $\epsilon > 0$ and let $\lambda := \{z_j: j = 0, \dots, m\} \in \Xi[a, b]$ be such that (2.17) holds. It is clear that (2.17) also holds when the partition λ is replaced by any its refinement. Let $\{x_i: i = 0, \dots, n\} \in \Xi[a, b]$ be a refinement of λ . Then we have

$$\begin{aligned} \sum_{i=1}^n |f(x_{i-1}-) - f(x_{i-1}+)|^p &= \sum_{i=1}^n \lim_{y_{i-1} \downarrow x_{i-1}, y_i \uparrow x_i} |f(y_i) - f(y_{i-1})|^p \\ &\leq \sum_{i=1}^n v_p(f; (x_{i-1}, x_i)) < \epsilon. \end{aligned}$$

Therefore statement (c) holds.

(c) $\Rightarrow$ (b). Let $\epsilon > 0$ and let $\lambda \in \Xi[a, b]$ be such that (2.18) holds for all its refinements. Since $\sigma_p(f) < \infty$, there exists a finite set $\nu \subset [a, b]$ such that

$$\sum_{x \in \mu} \left\{ |\Delta^- f(x)|^p + |\Delta^+ f(x)|^p \right\} < \epsilon$$

whenever a finite set $\mu \subset (a, b)$ and $\mu \cap \nu = \emptyset$. Let $\{z_j: j = 0, \dots, m\} := \lambda \cup \nu \in \Xi[a, b]$ and, for each $j = 1, \dots, m$, let $\kappa(j) := \{x_i^j: i = 0, \dots, n(j)\}$ be a set of points in (a, b) such that $z_{j-1} < x_0^j < \dots < x_{n(j)}^j < z_j$. Then we have

$$\begin{aligned} \sum_{j=1}^m s_p(f; \kappa(j)) &\leq 4^{p-1} \sum_{j=1}^m \sum_{i=1}^{n(j)} \left\{ |\Delta^+ f(x_{i-1}^j)|^p + |\Delta^- f(x_i^j)|^p \right\} \\ &\quad + 4^{p-1} \sum_{j=1}^m \sum_{i=1}^{n(j)} |f(x_i^j-) - f(x_{i-1}^j+)|^p < 4^p \epsilon/2. \end{aligned}$$

Since all partitions $\kappa(j)$ of (z_{j-1}, z_j) are arbitrary, it follows that

$$\sum_{j=1}^m v_p(f; (z_{j-1}, z_j)) \leq 4^p \epsilon / 2.$$

This proves statement (b). The proof of Lemma 9 is complete. $\square$

2.3 Extended Riemann-Stieltjes integrals

The refinement Riemann-Stieltjes and the Henstock-Kurzweil integrals are the best known extensions of the Riemann-Stieltjes integral. We refer to Graves [46], Hildebrandt [49] and McLeod [77] for detail expositions of the properties of the three integrals. Less well known are extensions of the Riemann-Stieltjes integral suggested by several members of the family of Young. This section is just a quick glance at relations between several different extensions of the Riemann-Stieltjes integral in the light of the p -variation conditions originated in the Theorem on Stieltjes integrability of L. C. Young [113, p. 264].

The Riemann-Stieltjes integral, or the RS integral, of a function f with respect to a function g is defined to exist and equal the limit of Riemann-Stieltjes sums as the mesh of partitions tends to zero. The RS integral exists first, if $f \in \mathcal{W}_p$, $g \in \mathcal{W}_q$ with $p^{-1} + q^{-1} > 1$, and second if f and g have no a discontinuity at the same point. The first condition cannot be replaced in general by the condition $p^{-1} + q^{-1} = 1$, and the second condition is necessary for the existence of the RS integral. In this section we recall several extensions of the RS integral which exist when either one of the two conditions, or both of them, fail to hold for a pair of functions (f, g) . We will say that an integral I_2 is an $(\mathcal{A}, \mathcal{B})$ extension of an integral I_1 , or shortly

$$I_1 \xrightarrow{(\mathcal{A}, \mathcal{B})} I_2, \tag{2.19}$$

if the following two statements hold:

- (a) for a class $\mathcal{A}$ of pairs of functions, if the I_1 integral exists then the I_2 integral so does;
- (b) for a class $\mathcal{B} \subset \mathcal{A}$, the converse implication holds, and the two integrals have the same value.

Typically, $\mathcal{A}$ is a class of pairs of regulated functions, and $\mathcal{B}$ is a subclass of the class $\mathcal{D}_p$ of all pairs (f, g) such that $f \in \mathcal{W}_p$, $g \in \mathcal{W}_q^*$ with $p^{-1} + q^{-1} = 1$, or vice versa. A summary of relations (2.19) between extended Riemann-Stieltjes integrals is given at the end of this section.

Notation 10 Let $[a, b]$ be a closed interval of real numbers. A set $\kappa = \{x_i: i = 0, \dots, n\}$ of points in $[a, b]$ is called a *partition of $[a, b]$* if $a = x_0 < x_1 < \dots < x_n = b$. For $i = 1, \dots, n$, a point $y_i \in [x_{i-1}, x_i]$ attached to a subinterval $[x_{i-1}, x_i]$ is called a *tag*, and the set $\tau = \tau(\kappa) = \{([x_{i-1}, x_i], y_i): i = 1, \dots, n\}$ is called a *tagged partition* of $[a, b]$ associated to κ . A tagged partition $\tau = \tau(\kappa)$ is a refinement of a partition λ if $\kappa \supset \lambda$. Let $\Xi[a, b]$ and $\Theta[a, b]$ be the sets of all partitions and all tagged partitions of $[a, b]$, respectively. Given a partition λ of $[a, b]$, let $R(\lambda)$ be the set of all tagged partitions which are refinements of λ . Let $\mathfrak{T}$ be the family of all sets $R = R(\lambda)$, $\lambda \in \Xi[a, b]$. Then $\mathfrak{T}$ is a direction in $\Theta[a, b]$ in the sense of Definition A.1.

The refinement Riemann-Stieltjes integral. Let f and g be real-valued functions defined on $[a, b]$. The *Riemann-Stieltjes sum* is a function $S_{RS} = S_{RS}(f, g)$ defined on $\Theta[a, b]$ by

$$S_{RS}(\tau) = S_{RS}(f, g; \tau) := \sum_{i=1}^n f(y_i)[g(x_i) - g(x_{i-1})]$$

for a tagged partition $\tau = \{([x_{i-1}, x_i], y_i) : i = 1, \dots, n\} \in \Theta[a, b]$. Then $(S_{RS}, \mathfrak{T})$ is the directed function. The *refinement Riemann-Stieltjes integral* $(RRS) \int_a^b f dg$ is defined to exist and equal the limit of the directed function $(S_{RS}, \mathfrak{T})$, that is,

$$(RRS) \int_a^b f dg := \lim_{\mathfrak{T}, \tau} S_{RS}(f, g; \tau) \quad (2.20)$$

provided the limit exists. Therefore $(RRS) \int_a^b f dg$ is defined to equal A if given $\epsilon > 0$ there exists a partition λ of $[a, b]$ such that $|S_{RS}(f, g; \tau) - A| < \epsilon$ for each tagged partition τ which is a refinement of λ . It is easy to ascertain that if $(RRS) \int_a^b f dg$ exists then for each $a \leq x < y \leq b$,

$$\lim_{z \downarrow x} [f(z) - f(x)][g(z) - g(x)] = \lim_{z \uparrow y} [f(z) - f(y)][g(z) - g(y)] = 0.$$

Therefore f and g cannot have common discontinuities on the same side at the same point provided their refinement Riemann-Stieltjes integral exists. This restriction does not allow to integrate an indicator function with respect to itself.

Theorem 11 *Let g and f be bounded functions on $[a, b]$. The integral $\int_a^b g df$ exists as the Riemann-Stieltjes integral if and only if it exists as the refinement Riemann-Stieltjes integral and the two functions g, f have no common discontinuities on $[a, b]$.*

The proof of this theorem is given in [46, p. 264] and in [49, p. 51]. We use extensively the preceding theorem when dealing with chain rules in the next chapter for functions having the quadratic λ -variation.

The refinement Young-Stieltjes integral. One way to resolve the problem of common discontinuities for the refinement Riemann-Stieltjes integral is to modify the Riemann-Stieltjes sum as was suggested by W.H. Young (1914)¹. Let g be a regulated function on $[a, b]$. The *refinement Young-Stieltjes integral* $(RYS) \int_a^b f dg$ is defined to be the limit, if it exists, as in (2.20) except that the Riemann-Stieltjes sum $S_{RS}(f, g; \tau)$ is replaced by the *Young-Stieltjes sum*

$$S_{YS}(f, g; \tau) := \sum_{i=1}^n \left\{ [f \Delta^+ g](x_{i-1}) + f(y_i)[g(x_i-) - g(x_{i-1}+)] + [f \Delta^- g](x_i) \right\}, \quad (2.21)$$

and the direction is restricted to all tagged partitions $\tau = \{([x_{i-1}, x_i], y_i) : i = 1, \dots, n\}$ of $[a, b]$ such that $x_{i-1} < y_i < x_i$ for $i = 1, \dots, n$. Clearly, the refinement Young-Stieltjes integral exists if either f , or g (or both) is a step function. Also, the refinement Young-Stieltjes integral exists if the refinement Riemann-Stieltjes integral so does, and the two integrals have the same value. This is so because any Young-Stieltjes sum (2.21) can be approximated arbitrarily closely by Riemann-Stieltjes sums based on refinements of $\{x_i : i = 1, \dots, n\} \in \Xi[a, b]$. More information about the refinement Young-Stieltjes integral one can find in [49, pp. 88-95].

¹In this work W.H. Young developed the integral equivalent to the Lebesgue-Stieltjes integral in a way which was latter extended by Daniell (1918)

The Central Young integral. L. C. Young [113] suggested another way to bypass the problem of common discontinuities for the refinement Riemann-Stieltjes integral. His integral is defined when both the integrand and integrator are regulated functions, and for such functions extends the refinement Young-Stieltjes integral (see Proposition 3.17 in [23, Part II]). For a regulated function f on $[a, b]$, define its left continuous modification $f_-^{(a)}$ on $[a, b]$ by

$$f_-^{(a)}(x) := f_-(x) := f(x-) \quad \text{for } a < x \leq b, \text{ and } f_-^{(a)}(a) := f(a).$$

Similarly, define the right continuous modification $f_+^{(b)}$ on $[a, b]$ by

$$f_+^{(b)}(x) := f_+(y) := f(y+) \quad \text{for } a \leq y < b, \text{ and } f_+^{(b)}(b) := f(b).$$

Let g and f be regulated functions on $[a, b]$. Define the Y_1 integral on $[a, b]$ by

$$(Y_1) \int_a^b f dg := (RRS) \int_a^b g_+^{(b)} df_-^{(a)} - \sum_{[a,b]} \Delta^+ f [g_+ - g_-^{(c)}] + [f \Delta^- g](b) \quad (2.22)$$

provided the RRS integral exists, and the sum converges unconditionally. Also, define the Y_2 integral on $[a, b]$ by

$$(Y_2) \int_a^b f dg := (RRS) \int_a^b g_-^{(a)} df_+^{(b)} + [f \Delta^+ g](a) + \sum_{(a,b]} \Delta^- f [g_+^{(b)} - g_-]$$

provided the RRS integral exists, and the sum converges unconditionally. The Y_1 and Y_2 integrals satisfy standard properties of integrals, such as linearity and additivity over adjacent intervals. Also, the Y_1 integral exists if and only if the Y_2 integral so does, and the integrals have the same value. Therefore for regulated functions g and f , the *Central Young* integral on $[a, b]$ is defined by

$$(CY) \int_a^b f dg := (Y_1) \int_a^b f dg = (Y_2) \int_a^b f dg$$

provided either the Y_1 or the Y_2 integral exists on $[a, b]$. All these facts are proved in [23, Section 3 of Part II] (see also [24] for simpler proofs). The integrals Y_1 and Y_2 defined Dudley [22] to clarify the original definition of L. C. Young [113]. The word “Central” in the name of the Central Young integral is used because there are its two other modifications: the Left Young and Right Young integrals discussed in Section 2.5 below.

The Henstock-Kurzweil integral. The above two extensions of the refinement Riemann-Stieltjes integral are defined for regulated functions. This restriction is not a problem if one wishes to apply these integrals to stochastic processes, which usually have regulated, and hence bounded sample functions. However there is another extension of the refinement Riemann-Stieltjes integral which may exist when neither the integrand nor the integrator are regulated, or bounded functions. Ward [108] working with the Perron-Stieltjes integral, modified its definition so that the new integral extends the refinement Riemann-Stieltjes integral and is defined for functions which need not be regulated. An equivalent definition of Ward’s extension was given indepepdently by Kurzweil [60] and Henstock [47]. The resulting integral is often called the Henstock-Kurzweil integral. Namely, for real-valued functions g and f on $[a, b]$, the Henstock-Kurzweil integral $(HK) \int_a^b f dg$ is defined to exist and equal A if given $\epsilon > 0$ there exists a

positive function $\delta(\cdot)$ on $[a, b]$ such that Riemann-Stieltjes sums $S_{RS}(g, f; \tau)$ differ from A at most by ϵ for all tagged partitions $\tau = \{([x_{i-1}, x_i], y_i): i = 1, \dots, n\}$ such that $y_i - \delta(y_i) \leq x_{i-1} \leq y_i \leq x_i \leq y_i + \delta(y_i)$ for $i = 1, \dots, n$. Several comparison results between the Henstock-Kurzweil and the refinement Young-Stieltjes integral are given in [23, Appendix F in Part I]. In particular, the Henstock-Kurzweil integral extends strictly the Lebesgue-Stieltjes integral. Most likely this fact made the Henstock-Kurzweil integral so popular among those currently working in Real Analysis. In this paper the Henstock-Kurzweil integral is used to improve earlier results related to the Black-Scholes formula in Financial Mathematics (see Theorem 5.13).

The L. C. Young Stieltjes integrability theorem. As it is mentioned in the introduction, in terms of the p -variation, the first and best possible existence conditions for the extended Riemann-Stieltjes integrals are due to L. C. Young [113] in 1936. Two years latter he published in [115] a different proof of a more general result using the ϕ -variation property of a function, an extension of the p -variation. In the first paper L. C. Young used, what we now call the Y_1 integral, while in the second one he used the refinement Young-Stieltjes integral. By Theorem F.2 in [23, Part I], L. C. Young's Stieltjes integrability theorem also applies to the Henstock-Kurzweil integral.

We formulate the L. C. Young Stieltjes integrability theorem in a form suitable for further references in the present notes. Recall that $\mathcal{W}_\infty[a, b]$ is the class $\mathcal{R}[a, b]$ of all regulated functions on $[a, b]$.

Theorem 12 *Let $1 \leq p < \infty$, and q be such that $p^{-1} + q^{-1} > 1$ if $p > 1$ and $q = \infty$ if $p = 1$. For $f \in \mathcal{W}_p[a, b]$ and $g \in \mathcal{W}_q[a, b]$, the integral $\int_a^b f dg$ exists*

- (a) *as the Riemann-Stieltjes integral if f and g have no discontinuities at the same point;*
- (b) *as the refinement Riemann-Stieltjes integral if f and g have no common discontinuities on the same side at the same point;*
- (c) *always as the refinement Young-Stieltjes integral, as the Central Young integral and as the Henstock-Kurzweil integral.*

In whichever of the five senses the integral exists, there exists a constant K if $p > 1$ which depends only on p and q such that the inequality

$$\left| \int_a^b \{g - g(y)\} df \right| \leq \begin{cases} \text{Osc}(g; [a, b]) v_1(f; [a, b]) & \text{if } p = 1, \\ K v_q(g; [a, b]) v_p(f; [a, b]) & \text{if } 1 < p < \infty, \end{cases} \quad (2.23)$$

holds for any $y \in [a, b]$.

In the case $p = 1$ the conclusion is well-known and easy to prove. In the case $p > 1$ the stated theorem with the Central Young integral in statement (c), follows from the theorem on Stieltjes integrability of L. C. Young [113]. Statement (c) for the other two integrals then follows from Corollary 3.20 of [23, Part II] and from Theorem F.2 of [23, Part I]. There are other proofs of the same result with a one of the three integrals in statement (c) and with a slightly different constant K in (2.23), in addition to already mentioned work of L. C. Young [115]. Necessary and sufficient Stieltjes integrability conditions in terms of the ϕ -variation have been established by D'yačkov [27]. These results also can be found [24].

Suppose that a function g is integrable with respect to a function f in the sense of any of the five integrals defined so far, and suppose that inequality (2.23) holds for some $1 \leq p < \infty$. Let $\Phi(x) := C + \int_a^x f dg$, $x \in [a, b]$, be the corresponding indefinite integral. Since each of the five integrals is additive over adjacent intervals, then in the case $p > 1$, we have the bound

$$s_p(\Phi; \kappa) \leq 2^{p-1} \left\{ K^p v_q(g)^{p/q} + \|g\|_{\sup}^p \right\} v_p(f)$$

for any partition κ of $[a, b]$, and so the p -variation of Φ is bounded in this case. A similar bound holds in the case $p = 1$, and so we get:

Corollary 13 *Under the assumptions of the preceding theorem, each of the five indefinite integrals Φ has bounded p -variation.*

The symmetric Young-Stieltjes integral. L. C. Young [113] showed that the p -variation conditions (1.6) in his Theorem on Stieltjes integrability cannot be replaced in general by the same condition with $p^{-1} + q^{-1} = 1$. However, if one restricts to integrals of the form

$$\int_a^b (\phi \circ f) df \tag{2.24}$$

where the integrand is a composition of the integrator with a smooth function ϕ , then the conditions (1.6) can be weakened. For example, by Proposition 4.4 of [85], the integral (RYS) $\int_a^b f df$ exists and satisfies the relation

$$(RYS) \int_a^b f df = \frac{1}{2} \left\{ f(b)^2 - f(a)^2 + \sum_{(a,b]} \{\Delta^- f\}^2 - \sum_{[a,b)} \{\Delta^+ f\}^2 \right\}$$

if and only if $f \in \mathcal{W}_2^*[a, b]$. It is interesting that one can modify Young-Stieltjes sums further so that the resulting integral, called the symmetric Young-Stieltjes integral, of the form (2.24) exists for some functions with infinite 2-variation. In particular, the symmetric Young-Stieltjes integral exists almost surely for a sample function of Brownian motion with respect to itself. At the same time the new integral agrees with the refinement Young-Stieltjes integral if the conditions (1.6) of L. C. Young's theorem are satisfied for functions with suitable values at their discontinuity points.

Let g and f be regulated functions on $[a, b]$. If all tags of a tagged partition $\tau = \{([x_{i-1}, x_i], y_i) : i = 1, \dots, n\}$ of $[a, b]$ satisfy the condition $x_{i-1} < y_i < x_i$ for $i = 1, \dots, n$, then we call τ a *Young tagged partition*. For a Young tagged partition τ , let

$$C_{YS}(g, f; \tau) := \sum_{i=1}^n \left\{ [\Delta^+ g \Delta^+ f](x_{i-1}) - [\Delta g \Delta f](x_{i-1}, y_i) + [\Delta g \Delta f](y_i, x_i) - [\Delta^- g \Delta^- f](x_i) \right\},$$

and let

$$\begin{aligned} T_{YS}(g, f; \tau) := S_{YS}(g, f; \tau) + \frac{1}{2} C_{YS}(g, f; \tau) = \frac{1}{2} \sum_{i=1}^n \left\{ [(g_+ + g) \Delta^+ f](x_{i-1}) \right. \\ \left. + [g_+(x_{i-1}) + g(y_i)] \Delta f(x_{i-1}, y_i) + [g(y_i) + g_-(x_i)] \Delta f(y_i, x_i) + [(g + g_-) \Delta^- f](x_i) \right\}, \end{aligned} \tag{2.25}$$

where $\Delta f(u, v) := f(v) - f(u)$, $\Delta f[u, v] := f(v) - f(u)$, and $[\Delta g \Delta f](J) := \Delta g(J) \Delta f(J)$. Let $R(\lambda)$ be the set of all Young tagged partitions which are refinements of a partition λ , and let $\mathfrak{R}$ be the family of all sets $R(\lambda)$, $\lambda \in \Xi[a, b]$. Then $\mathfrak{R}$ is the direction in the sense of Definition A.1.

Definition 14 Let g, f be regulated functions on $[a, b]$. Define the *symmetric Young-Stieltjes* integral $(SYS) \int_a^b g df$ to exist and equal the limit of the directed function $(T_{YS}(g, f; \cdot), \mathfrak{A})$. Also, define the *Young-Stieltjes quadratic covariation* $C_{YS}(g, f)$ on $[a, b]$ to exist and equal the limit of the directed function $(C_{YS}(g, f; \cdot), \mathfrak{A})$.

Sufficient existence conditions for the symmetric Young-Stieltjes integral are formulated in the next section. Here we present a formal relation between symmetric and refinement Young-Stieltjes integrals.

Theorem 15 Let g, f be regulated functions on $[a, b]$ satisfying the relations

$$\begin{cases} f = (f_+ + f_-)/2 \text{ on } (a, b) \\ g = (g_+ + g_-)/2 \text{ on } (a, b), g(a+) = g(a) \text{ and } g(b-) = g(b). \end{cases} \quad (2.26)$$

Then the following holds:

- (a) The integral $(RYS) \int_a^b f dg$ exists if and only if both the integral $(SYS) \int_a^b f dg$ exists and the quadratic covariation $C_{YS}(f, g) = 0$.
- (b) Assume in addition that for $p^{-1} + q^{-1} = 1$, one of the two functions f, g is in $\mathcal{W}_p^*[a, b]$ and the other one is in $\mathcal{W}_q[a, b]$. Then the integrals $(RYS) \int_a^b f dg$ and $(SYS) \int_a^b f dg$ both exist or not simultaneously, and if they exist then both have the same value.

Statements (a) and (b) are Theorem 5.5 and Proposition 5.6 proved in [85].

Summary. The relations (2.19) between extended Riemann-Stieltjes integrals can be summarized as follows:

$$RS \xrightarrow{(\mathcal{A}_1, \mathcal{B}_1)} RRS \xrightarrow{(\mathcal{A}_2, \mathcal{B}_2)} RYS \begin{cases} \xrightarrow{(\mathcal{A}_3, \mathcal{B}_3)} CY \\ \xrightarrow{(\mathcal{A}_4, \mathcal{B}_4)} SYS \\ \xrightarrow{\mathcal{A}_5} HK \end{cases}$$

Legend: Let $\mathcal{D}_p = \{(f, g): f \in \mathcal{W}_p, g \in \mathcal{W}_{p'}\}$ with $1/p + 1/p' = 1$. Then we have:

1. $\begin{cases} \mathcal{A}_1 = \{(f, g): f, g \in \mathbb{R}^{[a, b]}\} \\ \mathcal{B}_1 = \{(f, g): \forall x \in [a, b], \text{ either } f \text{ or } g \text{ is continuous at } x\}; \end{cases}$
2. $\begin{cases} \mathcal{A}_2 = \{(f, g): f \in \mathbb{R}^{[a, b]}, g \in \mathcal{R}[a, b]\} \\ \mathcal{B}_2 = \{(f, g): \text{ either } \Delta^+ f = \Delta^- g = 0 \text{ or } \Delta^- f = \Delta^+ g = 0\}; \end{cases}$
3. $\mathcal{A}_3 = \{(f, g): f, g \in \mathcal{R}[a, b]\}$ and $\mathcal{B}_3 = \mathcal{D}_p$;
4. $\mathcal{A}_4 = \mathcal{A}_3$ and $\mathcal{B}_4 = \mathcal{D}_p \cap \{(f, g): f = (f^+ + f^-)/2 \text{ and } g = (g^+ + g^-)/2\}$;
5. $\mathcal{A}_5 = \mathcal{D}_p$.

The refinement Riemann-Stieltjes integral with monotone integrator. The refinement Riemann-Stieltjes and the Riemann-Stieltjes integrals with respect to a monotone function appear in chain rules for a function having the quadratic λ -variation in the next chapter. Therefore we include here few remarks concerning these integrals. In text-books a Stieltjes type integral is often defined by restricting to monotone integrators, or to integrators having bounded variation. In that case the refinement Riemann-Stieltjes integral can be defined without explicit use of the limit under refinements of partitions, so that there may not be seen obvious connection between such integrals. Usually in text-books, an extension of the Riemann integral developed by Darboux, Peano and others is used. A Stieltjes type extension of such an integral was due to Pollard [94]. We state it next as the necessary and sufficient condition for the existence of the refinement Riemann-Stieltjes integral in the case when h is nondecreasing and f is bounded.

Let f be a bounded function on $[a, b]$ and let h be a nondecreasing function on $[a, b]$. For a partition $\kappa = \{x_i: i = 0, \dots, n\}$ of $[a, b]$, define the *lower and upper Riemann-Stieltjes sums*, respectively, by

$$\begin{cases} L_{RS}(\kappa) := L_{RS}(f, h; \kappa) := \sum_{i=1}^n m(f; [x_{i-1}, x_i])[h(x_i) - h(x_{i-1})] \\ U_{RS}(\kappa) := U_{RS}(f, h; \kappa) := \sum_{i=1}^n M(f; [x_{i-1}, x_i])[h(x_i) - h(x_{i-1})], \end{cases}$$

where, for any set J in the domain of f ,

$$m(f; J) := \inf\{f(x): x \in J\} \quad \text{and} \quad M(f; J) := \sup\{f(x): x \in J\}. \quad (2.27)$$

Since h is nondecreasing, it follows that $L_{RS}(\lambda) \leq U_{RS}(\kappa)$ and $U_{RS}(\lambda) \geq U_{RS}(\kappa)$ whenever $\kappa \supset \lambda$ for $\kappa, \lambda \in \Xi[a, b]$. Also, letting

$$\begin{cases} L_{RS} := L_{RS}(f, h; [a, b]) := \sup\{L_{RS}(f, h; \kappa): \kappa \in \Xi[a, b]\} \\ U_{RS} := U_{RS}(f, h; [a, b]) := \inf\{U_{RS}(f, h; \kappa): \kappa \in \Xi[a, b]\} \end{cases}$$

we have the relation $L_{RS} \leq U_{RS}$. Then one says that the *Darboux integral* $(D) \int_a^b f dh$ is defined and equals $b \in \mathbb{R}$ if $L_{RS} = U_{RS} = b$.

Let $\mathfrak{T}$ be the direction in the set of all tagged partitions of $[a, b]$ (see Notation 10), and let $S_{RS}(f, h; \tau)$ be the Riemann-Stieltjes sum based on a tagged partition $\tau \in \Theta[a, b]$. Since h is non decreasing it is easy to see that

$$L_{RS}(f, h; [a, b]) = \lim_{\mathfrak{T}} \inf S_{RS}(f, h) \quad \text{and} \quad U_{RS}(f, h; [a, b]) = \lim_{\mathfrak{T}} \sup S_{RS}(f, h),$$

where the right sides are defined by (A.1). By Theorem A.4, the integral $\int_a^b f dh$ exists as the Darboux integral if and only if it exists as the refinement Riemann-Stieltjes integrals, and then both integrals have the same value. It seems that the definition of an integral in terms of the limit in the sense of refinements of partitions is preferable because it is not restricted to integrands having bounded variation.

The refinement Young-Stieltjes integral with monotone integrator. As in the case of the refinement Riemann-Stieltjes integral, the refinement Young-Stieltjes integration with respect to monotone functions afford certain simplifications. Let f be a bounded function on $[a, b]$ and let h be a nondecreasing function on $[a, b]$. For $\kappa = \{x_i: i = 0, \dots, n\} \in \Xi[a, b]$, define the lower and upper Young sums, respectively, by

$$L_Y(f, h; \kappa) := \mathfrak{S}(f, h; \kappa) + \sum_{i=1}^n m(f; (x_{i-1}, x_i))[h(x_i-) - h(x_{i-1}+)] \quad (2.28)$$

$$U_Y(f, h; \kappa) := \mathfrak{S}(f, h; \kappa) + \sum_{i=1}^n M(f; (x_{i-1}, x_i)) [h(x_i-) - h(x_{i-1}+)], \quad (2.29)$$

where m, M are defined by (2.27), and

$$\mathfrak{S}(f, h; \kappa) := \sum_{i=0}^n f(x_i) [h_+^{(b)}(x_i) - h_-^{(a)}(x_{i-1})] = [f \Delta^+ f](a) + \sum_{i=1}^{n-1} [f \Delta^\pm h](x_i) + [f \Delta^- h](b).$$

Again as in the preceding paragraph, let $\mathfrak{T}$ be the direction in the set of all tagged partitions of $[a, b]$, and let $S_{YS}(f, h; \tau)$, $\tau \in \Theta[a, b]$, be the Young-Stieltjes sum defined by (2.21). Since h is nondecreasing function, we have that

$$\begin{cases} L_Y := L_Y(f, h; [a, b]) := \sup\{L_Y(f, h; \kappa) : \kappa \in Q([a, b])\} = \liminf_{\mathfrak{T}} S_{YS}(f, h) \\ U_Y := U_Y(f, h; [a, b]) := \inf\{U_Y(f, h; \kappa) : \kappa \in Q([a, b])\} = \limsup_{\mathfrak{T}} S_{YS}(f, h) \end{cases}$$

Then by Theorem A.4, the following holds:

Theorem 16 *Let f be a bounded function on $[a, b]$ and let h be a nondecreasing function on $[a, b]$. Then the integral (RYS) $\int_a^b f dh$ exists if and only if*

$$A := L_Y(f, h; [a, b]) = U_Y(f, h; [a, b]), \quad (2.30)$$

and then A is the value of the RYS integral.

Ross [98, §35], [97] defined the integral $\int_a^b f dh := A$ if (2.30) holds and proved several its properties based on this definition. However, he did not relate the integral so defined to the refinement Young-Stieltjes integral. Love [65, Section 5] does relate the integral of Ross with the refinement Young-Stieltjes integral but makes no a reference to its origin. In fact, as earlier as 1936, essentially the same integral have been developed by Glivenko [44, Section V].

2.4 Chain rules for extended Riemann-Stieltjes integrals

Here by a chain rule we mean an integral representation of the composition $\phi \circ g$ of two functions g and ϕ such that ϕ is smooth and g has possibly unbounded variation. The best known example of a chain rule is Itô's formula giving a stochastic integral representation of a smooth function with a semimartingale.

The refinement Young-Stieltjes integral representation. Let $B = \{B(t) : t \geq 0\}$ be a standard Brownian motion on a complete probability space $(\Omega, \mathcal{F}, \text{Pr})$. For any $p > 2$, we have that $v_p(B(\cdot, \omega); [0, 1]) < \infty$ for almost all $\omega \in \Omega$, that is the p -variation index for almost all sample functions of a Brownian motion is 2. Therefore it follows from the L. C. Young Stieltjes integrability theorem (Theorem 12) that the Riemann-Stieltjes integral

$$(RS) \int_0^1 f dB(\cdot, \omega) \quad (2.31)$$

is defined for almost all $\omega \in \Omega$ and for any function f having bounded p -variation for some $p < 2$. Since $v_2(B(\cdot, \omega); [0, 1]) = +\infty$ for almost all $\omega \in \Omega$, we cannot take f in (2.31) from the class $\mathcal{W}_2$, still less can we take f to be a sample function of a Brownian motion (see Section 3.8 below).

The condition $p^{-1} + q^{-1} > 1$ in Theorem 12 is best possible in general. It can be improved to the condition $p^{-1} + q^{-1} = 1$ provided one restricts the class of integrands to functions of a special form. That is, the existence part of the preceding theorem can be extended when the integrand f is the composition $\phi \circ g$ for some smooth function ϕ . Instead of the inequality (2.23), a chain rule formula then holds.

Next is a chain rule for a composition representation by the refinement Young-Stieltjes integral; its proof is in [85, Theorem 1.1].

Theorem 17 *For $0 < \alpha \leq 1$, let $g \in \mathcal{W}_{1+\alpha}^*[a, b]$, and let ϕ be differentiable with a derivative ϕ' satisfying a Lipschitz condition of order α . The refinement Young-Stieltjes integral (RYS) $\int_a^b (\phi' \circ g) dg$ is defined, and its value is determined by the relation*

$$\begin{aligned} (\phi \circ g)(b) - (\phi \circ g)(a) &= (RYS) \int_a^b (\phi' \circ g) dg \\ &+ \sum_{(a,b]} \left\{ \Delta^-(\phi \circ g) - (\phi' \circ g) \Delta^- g \right\} + \sum_{[a,b)} \left\{ \Delta^+(\phi \circ g) - (\phi' \circ g) \Delta^+ g \right\}, \end{aligned} \quad (2.32)$$

where the two sums are unconditional.

The statement of the preceding theorem also holds for the Central Young and Henstock-Kurzweil integrals in place of the refinement Young-Stieltjes integral. In Theorem 17, g cannot be taken to be a sample function of a Brownian motion. To extend the preceding chain rule formula to functions $g \in \mathcal{W}_p$ with $p < 3$ one can use the symmetric Young-Stieltjes integral. The proof of the next theorem is in [85, Theorem 1.4].

Theorem 18 *For $0 < \alpha \leq 1$, let $g \in \mathcal{W}_{2+\alpha}^*([a, b])$, and let ϕ be twice differentiable with a second derivative satisfying a Lipschitz condition of order α . The symmetric Riemann-Stieltjes integral (SRS) $\int_a^b (\phi' \circ g) dg$ is defined, and its value is determined by the relation*

$$\begin{aligned} (\phi \circ g)(b) - (\phi \circ g)(a) &= (SRS) \int_a^b (\phi' \circ g) dg \\ &+ \sum_{(a,b]} \left\{ \Delta^-(\phi \circ g) - (\phi' \circ g) \Delta^- g + \frac{\Delta^-(\phi' \circ g)}{2} \Delta^- g \right\} + \sum_{[a,b)} \left\{ \Delta^+(\phi \circ g) - (\phi' \circ g) \Delta^+ g - \frac{\Delta^+(\phi' \circ g)}{2} \Delta^+ g \right\}. \end{aligned}$$

Integration by parts formula. Using the multivariate variant of the chain rule for the refinement Young-Stieltjes integral we get an integration by parts formula: for $g, f \in \mathcal{W}_2^*[a, b]$, if $(RYS) \int_a^b g df$ exists then so does $(RYS) \int_a^b f dg$ and

$$(RYS) \int_a^b g df + (RYS) \int_a^b f dg = f(b)g(b) - f(a)g(a) + \sum_{(a,b]} \Delta^- f \Delta^- g - \sum_{[a,b)} \Delta^+ f \Delta^+ g, \quad (2.33)$$

where the two sums converge unconditionally (see [85, Corollary 4.3]). This formula under similar conditions was known already to L. C. Young, who used it in the proof of other results (see p. 611 in [115]). According to Hildebrandt [48, p. 276], formula (2.33) was proved by de Finetti and Jacob [32] under the assumption that g and f have bounded variation on $[a, b]$. We do not know at this writing whether the assumption $g, f \in \mathcal{W}_2^*$ could be replaced just by the assumption that the two sums in (2.33) converge unconditionally. However, a simple integration by parts formula holds for the symmetric Young-Stieltjes integral.

Theorem 19 *Let g, f be regulated functions on $[a, b]$. For $\sharp = SYS$, if $(\sharp) \int_a^b g df$ exists then so does $(\sharp) \int_a^b f dg$ and*

$$(\sharp) \int_a^b g df + (\sharp) \int_a^b f dg = f(b)g(b) - f(a)g(a). \quad (2.34)$$

Proof. Let $\tau = \{([x_{i-1}, x_i], y_i) : i = 0, \dots, n\}$ be a Young tagged partition of $[a, b]$. By definition (2.25) of *SYS* sums, we have

$$\begin{aligned} & T_{YS}(g, f; \tau) + T_{YS}(f, g; \tau) \\ &= \sum_{i=1}^n \left\{ \Delta^+(gf)(x_{i-1}) + \Delta(gf)(x_{i-1}, y_i) + \Delta(gf)[y_i, x_i] + \Delta^-(gf)(x_i) \right\} = \Delta(gf)[a, b]. \end{aligned}$$

Thus the directed function $(T_{YS}(g, f; \cdot), \mathfrak{R})$ has a limit if and only if the directed function $(T_{YS}(f, g; \cdot), \mathfrak{R})$ does, and (2.34) holds provided both limits exist. The proof is complete. $\square$

Under the conditions of Theorem 15, relating the symmetric and refinement Young-Stieltjes integral, we get the same integration by parts formula for the refinement Young-Stieltjes integral.

Corollary 20 *Let g, f be regulated functions on $[a, b]$ satisfying (2.26). For $\sharp = RYS$, if $(\sharp) \int_a^b g df$ exists then so does $(\sharp) \int_a^b f dg$ and (2.34) holds.*

Proof. Suppose that $(RYS) \int_a^b g df$ exists. Then by statement (a) of Theorem 15, $(SYS) \int_a^b g df$ exists, and

$$0 = C_{YS}(g, f) = C_{YS}(f, g). \quad (2.35)$$

By the form of the *SYS* sums, it also follows that the two integrals have the same value. Hence by the preceding theorem, $(SYS) \int_a^b f dg$ exists and the relations

$$(RYS) \int_a^b g df = (SYS) \int_a^b g df = \Delta(gf)[a, b] - (SYS) \int_a^b f dg.$$

hold. This in conjunction with (2.35) allow us to apply statement (a) of Theorem 15 in the opposite direction to conclude that $(RYS) \int_a^b f dg$ exists and (2.34) holds with $\sharp = RYS$. The proof of Corollary 20 is complete. $\square$

It is easy to see that if regulated functions g, f have discontinuities satisfying condition (2.26), then the two sums in formula (2.33) disappear and we get the integration by parts formula (2.34) with $\sharp = RYS$ without the assumption $g, f \in \mathcal{W}_2^*$.

If instead of condition (2.26), the regulated functions g, f on $[a, b]$ satisfy the same conditions except that jumps of g at the endpoints are allowed, then statement (a) of Theorem 15 holds with the value

$$C_{YS}(g, f) = \Delta^+ g(a) \Delta^+ f(a) - \Delta^- g(b) \Delta^- f(b) := A.$$

Following then the arguments of the proof of the preceding corollary, we get that the integration by parts formula

$$(RYS) \int_a^b g df + (RYS) \int_a^b f dg = f(b)g(b) - f(a)g(a) - A$$

holds for the refinement Young-Stieltjes integral. This is a result recently proved by Love [66, Theorem 25]. We notice that the R^3S integral in Love's work is the refinement Young-Stieltjes integral.

2.5 The Left and Right Young integrals

Next we define two different versions of the Central Young integral, that is the Left and Right Young integrals. These two integrals are better suited to solving linear integral equations (see Section 5.4 in [23, Part II]). For a regulated function f on $[a, b]$, the left-continuous function $f_-^{(a)}$ and the right-continuous function $f_+^{(b)}$ are defined by

$$f_-^{(a)}(x) := f_-(x) := f(x-) \quad \text{for } a < x \leq b, \quad f_-^{(a)}(a) := f(a) \quad (2.36)$$

and

$$f_+^{(b)}(x) := f_+(x) := f(x+) \quad \text{for } a \leq x < b, \quad f_+^{(b)}(b) := f(b), \quad (2.37)$$

respectively.

Definition 21 Let f and g be regulated functions on $[a, b]$. Define the *Left Young integral*, or the *LY* integral, by

$$(LY) \int_a^b g df := (RRS) \int_a^b g_-^{(a)} df_+^{(b)} + [g\Delta^+ f](a) + \sum_{(a,b)} \Delta^- g \Delta^+ f$$

provided $a < b$, the *RRS* integral exists and the sum converges unconditionally. We say that g is *LY* integrable with respect to f on $[a, b]$. Define the *Right Young integral*, or the *RY* integral, by

$$(RY) \int_a^b g df := (RRS) \int_a^b g_+^{(b)} df_-^{(a)} - \sum_{(a,b)} \Delta^+ g \Delta^- f + [g\Delta^- f](b)$$

provided $a < b$, the *RRS* integral exists and the sum converges unconditionally. We say that g is *RY* integrable with respect to f on $[a, b]$. If $a = b$ then both integrals are defined as 0.

The Left Young and Right Young integrals are defined in [23] (see relation (3.44) there) for Banach algebra valued functions. The present variants of the two integrals are special cases of the integrals with two integrands used to extend the Duhamel formula for product integrals. It is easy to see that if f is right-continuous then

$$(LY) \int_a^b g df = (RRS) \int_a^b g_-^{(a)} df \quad (2.38)$$

for each regulated function g . Similarly, if f is left-continuous then

$$(RY) \int_a^b g df = (RRS) \int_a^b g_+^{(b)} df.$$

The Left Young and Right Young integrals can be approximated by Left and Right Cauchy sums, respectively. For $\kappa = \{x_i: i = 0, \dots, n\} \in \Xi[a, b]$, define the Left Cauchy sum

$$S_{LC}(g, f; \kappa) := \sum_{i=1}^n g(x_{i-1})[f(x_i) - f(x_{i-1})],$$

and the Right Cauchy sum

$$S_{RC}(g, f; \kappa) := \sum_{i=1}^n g(x_i)[f(x_i) - f(x_{i-1})].$$

These sums define respectively the Left Cauchy-Stieltjes integral (*LCS*) $\int_a^b g df$ and the Right Cauchy-Stieltjes integral (*RCS*) $\int_a^b g df$ in the sense of refinements of partitions. Let $\Lambda[a, b]$ be the set of all nested sequences $\lambda = \{\lambda_m: m \geq 1\}$ of partitions of $[a, b]$ such that $\cup_m \lambda_m$ is dense in $[a, b]$.

Theorem 22 *For $1 \leq p < \infty$, let $f \in \mathcal{W}_p[a, b]$ and $g \in \text{dual}(\mathcal{W}_p)[a, b]$. Then g is Left Young integrable with respect to f on $[a, b]$ and the relation*

$$\lim_{m \rightarrow \infty} S_{LC}(g, f; \lambda_m) = (LY) \int_a^b g df \quad (2.39)$$

holds for each $\{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ such that

$$\{x \in (a, b): [\Delta^- g \Delta^+ f](x) \neq 0\} \subset \cup_m \lambda_m. \quad (2.40)$$

Also, g is Right Young integrable with respect to f on $[a, b]$ and the relation

$$\lim_{m \rightarrow \infty} S_{RC}(g, f; \lambda_m) = (RY) \int_a^b g df$$

holds for each $\{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ such that

$$\{x \in (a, b): [\Delta^+ g \Delta^- f](x) \neq 0\} \subset \cup_m \lambda_m.$$

Proof. We prove the theorem only for the Left Young integral and in the case $p = 1$ because a proof for the Right Young integral is similar and a proof for the case $p > 1$ is given in [83, Theorem 3]. Since $f_+^{(b)}$ has bounded variation and $g_-^{(a)}$ is regulated, the integral $(RRS) \int_a^b g_-^{(a)} df_+^{(b)}$ exists by Theorem 12.(b). Clearly the sum $\sum_{(a,b)} |\Delta^- g \Delta^+ f|$ converges absolutely, and hence unconditionally. Thus the integral $(LY) \int_a^b g df$ is defined. To prove relation (2.39) consider a sequence of partitions $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$ satisfying the stated assumptions. For each $m \geq 1$, let $g_m(x) := g(x_{i-1}^m)$ if $x \in [x_{i-1}^m, x_i^m]$ for some $i \in \{1, \dots, n(m)\}$ and let $g_m(b) := g(b)$. Using notation (2.36) and (2.37) for $[a, b] = [x_{i-1}^m, x_i^m]$, for each $i = 1, \dots, n(m)$, we have

$$(RRS) \int_{x_{i-1}^m}^{x_i^m} (g_m)_{-}^{(x_{i-1}^m)} df_{+}^{(x_i^m)} = g(x_{i-1}^m)[f(x_i^m) - f(x_{i-1}^m+)].$$

Using additivity of the *LY* integral proved in [83, Theorem 5], and since $\Delta^-(g_m) = 0$ on (x_{i-1}^m, x_i^m) , it then follows that

$$\begin{aligned} (LY) \int_a^b g_m df &= \sum_{i=1}^{n(m)} \left\{ (RRS) \int_{x_{i-1}^m}^{x_i^m} (g_m)_{-}^{(x_{i-1}^m)} df_{+}^{(x_i^m)} + [g_m \Delta^+ f](x_{i-1}^m) \right\} \\ &= S_{LCS}(g, f; \lambda_m). \end{aligned} \quad (2.41)$$

On the other hand, since g_m has only finitely many jumps, we have the representation

$$\begin{aligned} (LY) \int_a^b g_m df &= (RRS) \int_a^b (g_m)_{-}^{(a)} df_{+}^{(b)} + \sum_{i=1}^{n(m)-1} [\Delta^- g \Delta^+ f](x_i^m) \\ &\quad + \sum_{i=1}^{n(m)-1} [g(x_i^m-) - g(x_{i-1}^m)] \Delta^+ f(x_i^m) + [g \Delta^+ f](a). \end{aligned} \quad (2.42)$$

To prove

$$\lim_{m \rightarrow \infty} (RRS) \int_a^b (g_m)_-^{(a)} df_+^{(b)} = (RRS) \int_a^b g_-^{(a)} df_+^{(b)}. \quad (2.43)$$

we use the Osgood theorem on term by term integration (see e.g. [49, Theorem II.15.6]). It is clear that g_m are uniformly bounded. To show that $(g_m)_-^{(a)}$ converge to $g_-^{(a)}$, let $x \in (a, b]$. For each $m \geq 1$, there is $i \leq n(m)$ such that $x_{i-1}^m < x \leq x_i^m$. Then $g_m(x-) = g(x_{i-1}^m) \rightarrow g(x-)$ as $m \rightarrow \infty$. Thus all the hypotheses of the Osgood theorem are satisfied, and hence (2.43) holds. To prove

$$\lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)-1} [\Delta^- g \Delta^+ f](x_i^m) = \sum_{(a,b)} [\Delta^- g \Delta^+ f] \quad (2.44)$$

notice that each finite sum of terms $[\Delta^- g \Delta^+ f](x)$ is included into approximating sum for sufficiently large m by (2.40). To prove

$$\lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)-1} [g(x_i^m -) - g(x_{i-1}^m)] \Delta^+ f(x_i^m) = 0 \quad (2.45)$$

notice that each term tends to zero as $m \rightarrow \infty$ and any finite sum of them is bounded by $\sum |\Delta^+ f|$. Now (2.39) follows from (2.41) – (2.45). The proof is complete. $\square$

Corollary 23 *Under the hypotheses of Theorem 22, the LCS and RCS integrals exist and equal respectively to the LY and RY integrals.*

Proof. Suppose that the conclusion concerning the integral $(LCS) \int_a^b g df$ does not hold. Then there exists $\epsilon > 0$ such that for any $\lambda \in \Xi[a, b]$ the least upper bound of $|S_{LCS}(g, f; \kappa) - (LY) \int_a^b g df|$ over all refinements κ of λ is bigger than ϵ . Then one can construct recursively a sequence $\{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ satisfying (2.40) but not (2.39). A contradiction proves the first part of the corollary. The proof of the second one is similar.

Next is the Love-Young inequality for the Left Young and Right Young integrals. Recall that $\mathcal{W}_\infty[a, b]$ is the class $\mathcal{R}[a, b]$ of all regulated functions on $[a, b]$.

Theorem 24 *Let $1 \leq p < \infty$, and q be such that $p^{-1} + q^{-1} > 1$ if $p > 1$ and $q = \infty$ if $p = 1$. For $f \in \mathcal{W}_p[a, b]$ and $g \in \mathcal{W}_q[a, b]$, the integrals $(LY) \int_a^b g df$ and $(RY) \int_a^b g df$ are defined. Moreover, there exists a constant K if $p > 1$ which depends only on p and q such that the inequality*

$$\left| (\sharp) \int_a^b [g - g(y)] df \right| \leq \begin{cases} \text{Osc}(g; [a, b]) v_1(f; [a, b]) & \text{if } p = 1, \\ K v_q(g; [a, b]) v_p(f; [a, b]) & \text{if } 1 < p < \infty, \end{cases}$$

holds for $\sharp = LY, \sharp = RY$ and for any $y \in [a, b]$.

The proof follows easily from Theorem 12, Hölder's inequality and Theorem 22.

We finish with a formulation of several useful properties of the Left Young and Right Young integrals used in this paper.

Theorem 25 *Let g, f be regulated functions on $[a, b]$, and let $a \leq c \leq b$. For $\sharp = LY$ or RY , $(\sharp) \int_a^b g df$ exists if and only if both (sharp) $\int_a^c g df$ and (sharp) $\int_c^b g df$ exist, and then*

$$(\sharp) \int_a^b g df = (\sharp) \int_a^c g df + (\sharp) \int_c^b g df.$$

The direct proof of the preceding additivity property is given in [83, Theorem 5].

Next is an integration by parts formula for the Left Young and Right Young integrals.

Theorem 26 *Let g, f be regulated functions on $[a, b]$. If either of the two integrals $(LY) \int_a^b g df$ and $(RY) \int_a^b f dg$ exists then both exist, and*

$$(LY) \int_a^b g df + (RY) \int_a^b f dg = f(b)g(b) - f(a)g(a).$$

The proof of the theorem is given in [83, Theorem 7].

Let Φ and Ψ be *indefinite* Left Young and Right Young integrals defined respectively by

$$\Phi(x) := \text{const} + (LY) \int_a^x g dh \quad \text{and} \quad \Psi(y) := \text{const} + (RY) \int_y^b g dh$$

for $x, y \in [a, b]$. By the following statement Φ and Ψ , are regulated functions.

Proposition 27 *For regulated functions g and h on $[a, b]$ the following hold:*

- (a) *If g is Left Young integrable with respect to h on $[a, b]$ then Φ is a regulated function on $[a, b]$, and for $a \leq y < x \leq b$,*

$$(\Delta^- \Phi)(x) = [g_- \Delta^- h](x) \quad \text{and} \quad (\Delta^+ \Phi)(y) = [g \Delta^+ h](y).$$

- (b) *If g is Right Young integrable with respect to h on $[a, b]$ then Ψ is a regulated function on $[a, b]$, and for $a \leq y < x \leq b$,*

$$(\Delta^- \Psi)(x) = -[g \Delta^- h](x) \quad \text{and} \quad (\Delta^+ \Psi)(y) = -[g_+ \Delta^+ h](y).$$

The proof of the proposition is given in [83, Proposition 8].

Next is the *substitution rule* for the refinement Riemann-Stieltjes integrals. It is used in Chapter 4 and to derive the corresponding results for the Left Young and Right Young integrals.

Proposition 28 *Let $h \in \mathcal{W}_p[a, b]$ and $f, g \in \mathcal{W}_q[a, b]$ for some $p, q > 0$ with $p^{-1} + q^{-1} > 1$. Suppose that the pairs h, g and h, f have no common discontinuities on the left and on the right at the same point. Then g and fg are RRS integrable with respect to h , f is RRS integrable with respect to the indefinite RRS integral $\Phi(y) := \text{const} + (RRS) \int_a^y g dh$, $y \in [a, b]$, and*

$$(RRS) \int_a^b f d\Phi = (RRS) \int_a^b fg dh.$$

The proof of this proposition is given in [83, Proposition 9].

Next are substitution rules for the Left Young and Right Young integrals.

Theorem 29 *Let $h \in \mathcal{W}_p[a, b]$ and $f, g \in \mathcal{W}_q[a, b]$ for some $p, q > 0$ with $p^{-1} + q^{-1} > 1$. Then g and fg are LY integrable with respect to h , f is LY integrable with respect to the indefinite LY integral Ψ and*

$$(LY) \int_a^b f d\Psi = (LY) \int_a^b fg dh.$$

Similarly, g and fg are RY integrable with respect to h , f is RY integrable with respect to the indefinite RY integral Φ and

$$(RY) \int_a^b f d\Phi = (RY) \int_a^b fg dh.$$

The proof of this theorem is given in [83, Theorem 10].

The final statement provides chain rules for the Left Young and Right Young integral representations of a composition. Its proof is given in [83, Theorems 11 and 14].

Theorem 30 *For $\alpha \in [0, 1]$, let $f \in \mathcal{W}_{1+\alpha}^*[a, b]$ let ϕ be differentiable with the first derivative satisfying a Hölder condition of order α , and let ψ be a regulated function on $[a, b]$. Then the two equalities*

$$\begin{aligned} (LY) \int_a^b \psi d(\phi \circ f) &= (LY) \int_a^b \psi(\phi' \circ f) df \\ &+ \sum_{(a,b]} h_- [\Delta^-(\phi \circ f) - (\phi' \circ f)_- \Delta^- f] + \sum_{[a,b)} h [\Delta^+(\phi \circ f) - (\phi'_l \circ f) \Delta^+ f] \end{aligned}$$

and

$$\begin{aligned} (RY) \int_a^b \psi d(\phi \circ f) &= (RY) \int_a^b \psi(\phi' \circ f) df \\ &+ \sum_{(a,b]} h [\Delta^-(\phi \circ f) - (\phi' \circ f) \Delta^- f] + \sum_{[a,b)} h_+ [\Delta^+(\phi \circ f) - (\phi'_l \circ f)_+ \Delta^+ f] \end{aligned}$$

hold both meaning that the two integrals exist provided any one of the two exists, and the two sums converge unconditionally.

Chapter 3

Quadratic variation and restricted integrals of Stieltjes type

In mathematics, fruitful points of view are more powerful tools than so called key theorems. To say more precisely, they are eyes of a researcher, ardently trying to understand the essence of mathematical objects.

If the reality we seek to understand is rich and complicated then to grasp its depth and exquisiteness, we need as many such "eyes" as possible.

A. Grothendieck. "Recoltes et Semailles".

Recall that a regulated function f on $[a, b]$ has the local 2-variation, that is belongs to the Wiener class $\mathcal{W}_2^*[a, b]$, if and only if the limit

$$\lim_{\kappa, \mathfrak{P}} s_2(f; \kappa) \quad \left(= \sum_{(a,b]} (\Delta^- f)^2 + \sum_{[a,b)} (\Delta^+ f)^2 \right) \quad (3.1)$$

exists in the sense of refinements of partitions of the interval $[a, b]$ (see (1.4) and Section 2.2). In this chapter we consider a class of functions such that instead of (3.1), the limit exists along a nested sequence λ of partitions densely filling the interval $[a, b]$. We call such a property of f , if it holds, the quadratic λ -variation of f . In a similar manner we modify a Stieltjes type integral. Namely, we consider two variants of a λ -integral defined to be a limit, if it exists, of special type Riemann-Stieltjes sums along a sequence λ . The two integrals, when they exist, are called the Left Cauchy and Right Cauchy λ -integrals, respectively. Existence of the quadratic λ -variation of a function f implies the existence of the two λ -integrals of a composition $\phi \circ f$ with respect to f provided ϕ is a smooth function. In Section 3.6, the extended Doléans exponential is shown to be the unique solution to the augmented linear λ -integral equation. The main result is in Section 3.7, where the evolution representation problem is solved for a class of functions having the quadratic λ -variation.

3.1 The quadratic λ -variation and λ -covariation

The quadratic λ -variation: definition and examples. We start with a notation related to partitions of intervals.

Notation 1 Let J be a bounded nondegenerate interval of real numbers, open or closed at either end. A *partition* of J is a finite increasing sequence $\{t_0, t_1, \dots, t_n\}$ of points of J , where t_0 is the left endpoint of J if it is left closed and t_n is the right endpoint of J if it is right closed. The class of all sequences $\lambda = \{\lambda_m: m \geq 1\}$ of partitions $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ of J such that $\lambda_m \subset \lambda_{m+1}$ for $m = 1, 2, \dots$ and $\cup_m \lambda_m$ is dense in the closure of J is denoted by $\Lambda(J)$.

Next we define a partition of an extended subinterval $\llbracket u, v \rrbracket$, $(u, v) \in S\llbracket a, b \rrbracket$, induced by a partition of an ordinary interval $[a, b]$.

Notation 2 Let $\kappa = \{t_i: i = 0, \dots, n\}$, $n \geq 2$, be a partition of $[a, b]$, and let $(u, v) \in S\llbracket a, b \rrbracket$. First suppose that $u \in \{x, x+\}$ and $v \in \{y-, y\}$ for some $a \leq x < y \leq b$. If between u and v there are no points of κ , then let $\kappa \cap \llbracket u, v \rrbracket := \{u, v\}$. Otherwise denote by $i(u)$ and $i(v)$ the two indices in $\{1, \dots, n\}$ such that $t_{i(u)-1} \leq u < t_{i(u)}$ and $t_{i(v)-1} < v \leq t_{i(v)}$, respectively. Denoting $x_{i(u)-1} := u$, $x_{i(v)} := v$ and $x_i := t_i$ for $i(u) \leq i < i(v)$, let

$$\kappa \cap \llbracket u, v \rrbracket := \{u, t_{i(u)}, \dots, t_{i(v)-1}, v\} = \{x_{i(u)-1}, \dots, x_{i(v)}\}.$$

Now suppose that $u = x-$ and $v \in \{y-, y\}$ for some $a < x \leq y \leq b$. If $x \in \kappa$ then $\kappa \cap \llbracket u, v \rrbracket$ is defined as before. If $x \notin \kappa$ then let $\kappa \cap \llbracket u, v \rrbracket := (\kappa \cup \{x\}) \cap \llbracket u, v \rrbracket$. Finally suppose that $v = y+$ for some $a \leq y < b$. If $y \in \kappa$ then $\kappa \cap \llbracket u, v \rrbracket$ is defined as before. If $y \notin \kappa$ then let $\kappa \cap \llbracket u, v \rrbracket := (\kappa \cup \{y\}) \cap \llbracket u, v \rrbracket$. Then $\kappa \cap \llbracket u, v \rrbracket$ is called the *trace partition* of $\llbracket u, v \rrbracket$ induced by κ .

Notice that the trace partition of $\llbracket u, v \rrbracket$ consists of ordinary points except possibly for the endpoints u and v . If $(u, v) \in S[a, b]$, that is, if $[u, v]$ is an ordinary interval then we write $\kappa \cap [u, v]$ instead of $\kappa \cap \llbracket u, v \rrbracket$. Recalling notation (1.3) with $p = 2$, for a partition $\kappa = \{x_i: i = 0, \dots, n\}$ and any $a \leq t \leq b$, we have

$$s_2(f; \kappa \cap [a, t]) = \sum_{i=1}^n [f(x_i \wedge t) - f(x_{i-1} \wedge t)]^2.$$

Also by the definition of the trace partition, we have

$$\begin{cases} s_2(f; \kappa \cap \llbracket s-, s \rrbracket) = s_2(f; \{s-, s\}) = (\Delta^- f(s))^2, \\ s_2(f; \kappa \cap \llbracket t, t+ \rrbracket) = s_2(f; \{t, t+\}) = (\Delta^+ f(t))^2 \end{cases}$$

for any partition κ of $[a, b]$ and for any $a \leq t < s \leq b$.

Definition 3 Let f be a regulated function on $[a, b]$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. We say that f has the *quadratic λ -variation* on $[a, b]$ if there exists an additive upper continuous function $\alpha_\lambda(f)$ on $S\llbracket a, b \rrbracket$ such that for each $(u, v) \in S\llbracket a, b \rrbracket$,

$$\alpha_\lambda(f; u, v) = \lim_{m \rightarrow \infty} s_2(f; \lambda_m \cap \llbracket u, v \rrbracket). \quad (3.2)$$

If f has the quadratic λ -variation on $[a, b]$ then the right distribution function of $\alpha_\lambda(f)$ restricted to $[a, b]$ is called the *bracket function* of f and is denoted by $[f]_\lambda$, that is, $[f]_\lambda(t) = \alpha_\lambda(f; a, t)$ for $t \in [a, b]$ (cf. (2.4)).

Alternatively the quadratic λ -variation on an interval $[a, b]$ can be defined solely in terms of a regulated function on $[a, b]$ as follows:

Proposition 4 *Let f be a regulated function on $[a, b]$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. The function f has the quadratic λ -variation on $[a, b]$ if and only if there exists a regulated function $H = H_\lambda$ on $[a, b]$ such that $H(a) = 0$ and for each $(s, t) \in S[a, b]$,*

$$H(t) - H(s) = \lim_{m \rightarrow \infty} s_2(f; \lambda_m \cap [s, t]), \quad (3.3)$$

$$\Delta^- H(t) = (\Delta^- f(t))^2 \quad \text{and} \quad \Delta^+ H(s) = (\Delta^+ f(s))^2. \quad (3.4)$$

If the two statements hold then $[f]_\lambda = H$. Moreover, condition (3.3) for each $(s, t) \in S[a, b]$, can be replaced by the condition

$$H(t) = \lim_{m \rightarrow \infty} s_2(f; \lambda_m \cap [a, t]), \quad (3.5)$$

for each $t \in [a, b]$, provided two-sided discontinuity points of f are accessible by λ , that is (1.19) holds.

Proof. First suppose that there exists a regulated function H on $[a, b]$, equal to zero at a , and such that (3.3) and (3.4) hold. Define the function $\alpha_\lambda(f)$ on $S[[a, b]]$ by $\alpha_\lambda(f; u, v) := H(v) - H(u)$ for $(u, v) \in S[[a, b]]$. Then $\alpha_\lambda(f)$ is additive and upper continuous function on $S[[a, b]]$ by Theorem 2.4. We prove (3.2) only for the cases $(u, v) = (a, t-)$ and $(u, v) = (a, t+)$ because the proofs for the other cases are similar. To this aim let $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$, $m = 1, 2, \dots$, and let $t \in (a, b]$. For each $m \geq 1$, there is an index $i(t) \in \{1, \dots, n(m) - 1\}$ such that $x_{i(t)}^m < t \leq x_{i(t)+1}^m$. Then by (3.3) and (3.4), we have

$$\begin{aligned} \lim_{m \rightarrow \infty} s_2(f; \lambda_m \cap [[a, t-]]) &= \lim_{m \rightarrow \infty} \left\{ s_2(f; \lambda_m \cap [a, t]) - [f(t) - f(x_{i(t)}^m)]^2 + [f(t-) - f(x_{i(t)}^m)]^2 \right\} \\ &= H(t) - (\Delta^- f(t))^2 = H(t-) = \alpha_\lambda(f; a, t-), \end{aligned}$$

proving (3.2) for $(u, v) = (a, t-) \in S[[a, b]]$. Now let $t \in [a, b]$. By the definition of the trace partition, we have for each $m \geq 1$,

$$s_2(f; \lambda_m \cap [[a, t+]]) = s_2(f; \lambda_m \cap [a, t]) + (\Delta^+ f(t))^2,$$

proving (3.2) for $(u, v) = (a, t+) \in S[[a, b]]$.

To prove the converse implication, let H be the restriction to $[a, b]$ of the right distribution function of $\alpha_\lambda(f)$, that is, $H(t) := \alpha_\lambda(f; a, t)$ for $t \in [a, b]$. By statement (iv) of Theorem 2.4 and (3.2), H is regulated on $[a, b]$, is zero at a , and satisfies properties (3.3), (3.4).

To prove the last part of the conclusion, let $(s, t) \in S[a, b]$ be such that $s < t$, and let (3.5) holds for each $t \in [a, b]$. For each $m \geq 1$, let $i(s) = i_m(s) \in \{1, \dots, n(m)\}$ be the index such that $x_{i(s)-1}^m < s \leq x_{i(s)}^m$. Then we have

$$\begin{aligned} s_2(f; \lambda_m \cap [s, t]) &= s_2(f; \lambda_m \cap [a, t]) - s_2(f; \lambda_m \cap [a, s]) \\ &\quad + [f(x_{i(s)}^m) - f(s)]^2 + [f(s) - f(x_{i(s)-1}^m)]^2 - [f(x_{i(s)}^m) - f(x_{i(s)-1}^m)]^2. \end{aligned}$$

Since a two-sided discontinuity point eventually is a partition point, by (3.5), the right side converges to $H(t) - H(s)$ as $m \rightarrow \infty$, proving (3.3). The proof of Proposition 4 is now complete. $\square$

Suppose that for some $\lambda \in \Lambda[a, b]$, a regulated function f on $[a, b]$ has the quadratic λ -variation. By (3.3), the bracket function $H = [f]_\lambda$ is nondecreasing on $[a, b]$, and so it has a decomposition into a sum of a pure jump function and a continuous function. By (3.4), the pure jump function of $[f]_\lambda$ is given by $\sum_{(a, t]} \{\Delta^- f\}^2 + \sum_{[a, t)} \{\Delta^+ f\}^2$, $t \in [a, b]$. The difference

between $[f]_\lambda$ and its pure jump function is nondecreasing continuous function on $[a, b]$, and is denoted by $[f]_\lambda^c$, so that for each $a \leq t \leq b$,

$$[f]_\lambda(t) = [f]_\lambda^c(t) + \sum_{(a,t]} \{\Delta^- f\}^2 + \sum_{[a,t)} \{\Delta^+ f\}^2. \quad (3.6)$$

The notion of the quadratic λ -variation is similar to the one defined by Föllmer [34] for a regulated and right-continuous function on $[0, \infty)$. We recall his definition in the case when a regulated and right-continuous function f is defined on a closed interval $[a, b]$. For a sequence $\lambda = \{\lambda_m: m \geq 1\}$ of partitions $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$ of $[a, b]$ such that the mesh tends to zero as $m \rightarrow \infty$, let

$$\xi_m := \sum_{i=1}^{n(m)} [f(x_i^m) - f(x_{i-1}^m)]^2 \epsilon_{x_{i-1}^m},$$

where ϵ_x is a point mass at x . The function f has the quadratic variation on $[a, b]$ in the sense of Föllmer provided a Radon measure ξ exists on Borel sets of $[a, b]$ such that point measures ξ_m converge weakly to ξ as $m \rightarrow \infty$, and $\xi(\{t\}) = (\Delta^- f(t))^2$ for each $a < t \leq b$. By Theorem 11 below, if for such a function, the quadratic λ -variation exists then Föllmer's quadratic variation also exists, and the measure ξ is the Lebesgue-Stieltjes measure on $[a, b]$ generated by the bracket function $[f]_\lambda$.

Example 5 If f has the local 2-variation on $[a, b]$ and the two-sided discontinuity points of f are accessible by $\lambda \in \Lambda[a, b]$, then f has the quadratic λ -variation, and its bracket function $[f]_\lambda$ is the pure jump function given by

$$[f]_\lambda(t) = \sum_{(a,t]} \{\Delta^- f\}^2 + \sum_{[a,t)} \{\Delta^+ f\}^2 \quad (3.7)$$

for $a \leq t \leq b$ (see Corollary 15 below). If f is either right-continuous or left-continuous then assumption (1.19) always holds because in such cases the set on the left side is empty.

Relation (3.7) may be compared with the property (3.1) defining the Wiener class $\mathcal{W}_2^*$. Given $\lambda \in \Lambda[0, 1]$, almost every sample function $B(\cdot, \omega)$ of a standard Brownian motion B has the quadratic λ -variation on $[0, 1]$ with the bracket function $[B(\cdot, \omega)]_\lambda(t) = t$, $0 \leq t \leq 1$, and since it is continuous $B(\cdot, \omega)$ cannot have the local 2-variation (see Theorem 2.8). Next we give an example of a non-random function with the same properties.

Let f be a real-valued function defined on $[0, 1]$, let $0 < H < 1$, and let $r \geq 4$ be an integer. Following Kôno [57] and [58], the function f is called *self-affine with the scale parameter H to base r* if for any $i = 0, \dots, r^m - 1$, $m = 1, 2, \dots$, there are $\epsilon_{i,m} \in \{1, -1\}$ and the relation

$$f(ir^{-m} + h) - f(ir^{-m}) = \epsilon_{i,m} r^{-mH} f(r^m h) \quad (3.8)$$

holds for each $0 \leq h < r^{-m}$. A bounded self-affine function is continuous if and only if (3.8) holds for all $0 \leq h \leq r^{-m}$. For any rational number $0 < H < 1$, one can construct a continuous self-affine function of the scale parameter H satisfying $0 \leq f \leq 1$, $f(0) = 0$ and $f(1) = 1$. Typical examples of self-affine functions, such as the coordinate functions of the Peano curve, can be found in [57]. By the definition, a continuous self-affine function f of the scale parameter H is Hölder continuous of order H , and for each $t, s \in [0, 1]$,

$$|f(t) - f(s)| \leq 5M|t - s|^H, \quad (3.9)$$

where $M := \sup_{0 \leq t \leq 1} |f(t)|$. Also, a non-trivial self-affine function with the scale parameter H does not satisfy the Hölder continuity of order $H' > H$ at any point (see Theorem 1 in [57]). A more general class of self-affine functions were considered by Kamae [54] and Bertoin [6].

Example 6 Let f be a continuous self-affine function on $[0, 1]$ with the scale parameter $1/2$ to base r for some integer $r \geq 4$, such that $f(1) = 1$, and let $\lambda_m := \{ir^{-m}: i = 0, \dots, r^m\}$. Then $\lambda := \{\lambda_m: m \geq 1\} \in \Lambda[0, 1]$, and f has the quadratic λ -variation with the bracket function $[f]_\lambda(t) = t$ for $0 \leq t \leq 1$. By (3.9) and Theorem 2.8, it then follows that $f \in \mathcal{W}_2 \setminus \mathcal{W}_2^*$.

Let $X = \{X(t): t \geq 0\}$ be a square integrable continuous martingale on a filtered complete probability space $(\Omega, \mathcal{F}, \Pr, \{\mathcal{F}_t: t \geq 0\})$ satisfying the usual hypotheses. Meyer [78],[79] proved that there exists a unique increasing stochastic process $[X] = \{[X](t): t \geq 0\}$ such that almost surely,

$$E\{[X(t) - X(s)]^2 | \mathcal{F}_s\} = E\{[X](t) | \mathcal{F}_s\} - [X](s)$$

for $0 < s < t < \infty$. It can be shown that for $0 < T < \infty$, almost every sample function of X has the quadratic λ -variation on $[0, T]$ for some $\lambda \in \Lambda[0, T]$ as follows. We say that a sequence $\{\tau_i: i = 1, 2, \dots\}$ of stopping times is a *random partition of $[0, \infty)$ composed by stopping times* if almost surely, $\tau_0 = 0$, $\tau_i \leq \tau_{i+1}$ and $\tau_i \rightarrow +\infty$ with $i \rightarrow \infty$. Kunita and Watanabe [59, Theorem 1.3] proved that there exists a sequence $\lambda = \{\lambda_m: m \geq 1\}$ of random partitions of $[0, \infty)$ composed by stopping times $\lambda_m = \{\tau_i^m: i = 1, 2, \dots\}$ such that almost surely, $\lambda_m \subset \lambda_{m+1}$, $\max_i |\tau_{i+1}^m - \tau_i^m| \rightarrow 0$ as $m \rightarrow \infty$, and

$$\lim_{m \rightarrow \infty} \sum_{i=0}^{\infty} [X(\tau_{i+1}^m \wedge t) - X(\tau_i^m \wedge t)]^2 = [X](t) - [X](0) \quad \text{for } t \geq 0.$$

For $\omega \in \Omega$ and $0 < T < \infty$, we have $\lambda_m(\omega) \cap [0, T] = \{\tau_i^m(\omega) \wedge T: i = 0, 1, \dots\}$, $m \geq 1$, and $\lambda(\omega, T) := \{\lambda_m(\omega) \cap [0, T]: m \geq 1\} \in \Lambda[0, T]$. Therefore by the second part of Proposition 4, we have:

Example 7 Let X be a square integrable continuous martingale on a complete probability space $(\Omega, \mathcal{F}, \Pr)$. There exist a sequence λ of random partitions of $[0, \infty)$ composed by stopping times and a subset $\Omega_0 \subset \Omega$ such that $\Pr(\Omega_0) = 1$, and for each $0 < T < \infty$ and $\omega \in \Omega_0$, $X(\cdot, \omega)$ has the quadratic $\lambda(\omega, T)$ -variation on $[0, T]$ with the bracket function

$$[X(\cdot, \omega)]_{\lambda(\omega, T)}(t) = [X](t, \omega) - [X](0, \omega) \quad \text{for } 0 \leq t \leq T.$$

Let k be a positive integer, and let B be a standard Brownian motion. A stochastic process $X = \{X(t): 0 \leq t \leq 1\}$ is called a *weak Brownian motion of order k* if for every k -tuple $(t_1, t_2, \dots, t_k)$, the equality

$$(X(t_1), X(t_2), \dots, X(t_k)) = (B(t_1), B(t_2), \dots, B(t_k))$$

holds in distribution. If $k \geq 4$ the process X has the same quadratic variation as a standard Brownian motion B , even so X may not be a semimartingale. For this and many other properties of a weak Brownian motion see Föllmer, Wu and Yor [36].

Example 8 Let $X = \{X(t): 0 \leq t \leq 1\}$ be a weak Brownian motion of order $k \geq 4$, and let $\lambda \in \Lambda[0, 1]$. By Proposition 2.1 of [36] and its proof, X admits a continuous version $\bar{X}$ on a complete probability space $(\Omega, \mathcal{F}, \Pr)$ such that for almost all $\omega \in \Omega$, $\bar{X}(\cdot, \omega)$ has the quadratic λ -variation with the bracket function $[\bar{X}(\cdot, \omega)]_\lambda(t) = t$ for $0 \leq t \leq 1$.

Integrals with respect to the bracket function. Since the bracket function $[f]_\lambda$, if exists, is nondecreasing, any continuous and bounded function is Riemann-Stieltjes integrable with respect to $[f]_\lambda$. We show next that any regulated function is Left Young and Right Young integrable with respect to a function V of bounded variation, and prove an integral representation corresponding to the decomposition of V into the continuous and pure jump parts (see Definition 2.21 for the definitions of the Left and Right Young integrals).

Lemma 9 *Let ψ be a regulated function on $[a, b]$, and let V be a function on $[a, b]$ of bounded variation such that for $a \leq x \leq b$,*

$$V(x) = V^c(x) + \Delta^+ V(a) + \sum_{(a,x)} \{\Delta^- V + \Delta^+ V\} + \Delta^- V(x), \quad a \leq x \leq b,$$

where V^c is a continuous function. Then ψ is Left Young integrable with respect to V , it is Riemann-Stieltjes integrable with respect to V^c , and

$$(LY) \int_a^b \psi dV = (RS) \int_a^b \psi dV^c + \sum_{(a,b]} \psi_- \Delta^- V + \sum_{[a,b)} \psi \Delta^+ V. \quad (3.10)$$

Also, ψ is Right Young integrable with respect to V , it is Riemann-Stieltjes integrable with respect to V^c , and

$$(RY) \int_a^b \psi dV = (RS) \int_a^b \psi dV^c + \sum_{(a,b]} \psi \Delta^- V + \sum_{[a,b)} \psi_+ \Delta^+ V. \quad (3.11)$$

Remark. We notice that using the Central Young integral in the preceding statement instead of the Left or Right Young integrals, gives the usual representation:

$$(CY) \int_a^b \psi dV = (RS) \int_a^b \psi dV^c + \sum_{(a,b]} \psi \Delta^- V + \sum_{[a,b)} \psi \Delta^+ V.$$

Proof. Due to boundedness of the variation of V , the sum in the definition of $(LY) \int_a^b \psi dV$ converges absolutely, so that it is enough to prove the existence of the RRS integral of $\psi_-^{(a)}$ with respect to $V_+^{(b)}$. Recall that $\Delta_a^- V$ and $\Delta_b^+ V$ are defined on $[a, b]$ so that $\Delta_a^- V(a) = 0$ with $\Delta_a^- V = \Delta^- V$ on $(a, b]$, and $\Delta_b^+ V(b) = 0$ with $\Delta_b^+ V = \Delta^+ V$ on $[a, b)$, respectively. Then $V_+^{(b)} = V^c + U$ on $[a, b]$, where $U(x) := \sum_{[a,x]} \{\Delta_a^- V + \Delta_b^+ V\}$ for $a \leq x \leq b$. First, we prove that $\psi_-^{(a)}$ is RRS integrable with respect to U and

$$(RRS) \int_a^b \psi_-^{(a)} dU = \sum_{(a,b]} \psi_- \{\Delta^- V + \Delta_b^+ V\} =: A. \quad (3.12)$$

To this aim let $\epsilon > 0$. There exists a partition $\mu = \{z_j: j = 0, \dots, m\}$ of $[a, b]$ such that

$$\|\psi\|_\infty \sum_{j=1}^m \sum_{(z_{j-1}, z_j)} \{|\Delta^- V| + |\Delta^+ V|\} < \epsilon \quad \text{and} \quad \left| A - \sum_{\mu \setminus \{a\}} \psi_- \{\Delta^- V + \Delta_b^+ V\} \right| < \epsilon.$$

Then choose $\{u_j: j = 1, \dots, m\} \subset (a, b)$ such that $z_{j-1} < u_j < z_j$ for $j = 1, \dots, m$ and

$$\max_{1 \leq j \leq m} \sup_{u_j \leq x < z_j} |\psi_-(x) - \psi_-(z_j)| < \epsilon/v_1(V).$$

Let $\tau = \{([x_{i-1}, x_i], y_i) : i = 1, \dots, n\}$ be a tagged partition of $[a, b]$ such that $\{u_j, z_j : j = 1, \dots, m\} \subset \{x_i : i = 1, \dots, n\}$. For $j = 1, \dots, m$, let $i(j)$ be the index in $\{1, \dots, n\}$ such that $x_{i(j)} = z_j$, and let $I(j) := \{i(j-1) + 1, \dots, i(j)\}$. Then we have

$$\begin{aligned} |S_{RS}(\psi_-^{(a)}, U; \tau) - A| &\leq \epsilon + \left| \sum_{j=1}^m \sum_{i \in I(j)} \psi_-(y_i) \sum_{(x_{i-1}, x_i]} \{\Delta^- V + \Delta_b^+ V\} - \sum_{\mu \setminus \{a\}} \psi_- \{\Delta^- V + \Delta_b^+ V\} \right| \\ &\leq \epsilon + \|\psi\|_\infty \sum_{j=1}^m \sum_{(z_{j-1}, z_j)} \{|\Delta^- V| + |\Delta^+ V|\} + \sum_{j=1}^m |\psi_-(y_{i(j)}) - \psi_-(z_j)| |\Delta^- V(z_j) + \Delta^+ V(z_j)| < 3\epsilon. \end{aligned}$$

Therefore $\psi_-^{(a)}$ is *RRS* integrable with respect to U , and (3.12) holds. Since ψ is regulated, we have that $(RS) \int_a^b \psi dV^c$ exists and $(RS) \int_a^b \Delta_a^- \psi dV^c = 0$. This in conjunction with (3.12) yields that $\psi_-^{(a)}$ is *RRS* integrable with respect to $V_+^{(b)}$ and

$$(RRS) \int_a^b \psi_-^{(a)} dV_+^{(b)} = (RS) \int_a^b \psi dV^c + \sum_{(a,b]} \psi_- \{\Delta^- V + \Delta_b^+ V\}.$$

Adding the preceding sum with the sum from the definition of the Left Young integral we get

$$\sum_{(a,b]} \psi_- \{\Delta^- V + \Delta_b^+ V\} + \{\psi \Delta^+ V\}(a) + \sum_{(a,b)} \Delta^- \psi \Delta^+ V = \sum_{(a,b]} \psi_- \Delta^- V + \sum_{[a,b)} \psi \Delta^+ V.$$

Therefore ψ is Left Young integrable with respect to V , and (3.10) holds.

The proof of the second part of the lemma is similar. Indeed, using the same notation as before we have $V_-^{(a)} = V^c + U$, where $U(a) := 0$ and $U(x) := \sum_{[a,x)} \{\Delta_a^- V + \Delta_b^+ V\}$ for $a < x \leq b$. Then one can show similarly that $\psi_+^{(b)}$ is *RRS* integrable with respect to U , and

$$(RRS) \int_a^b \psi_+^{(b)} dU = \sum_{[a,b)} \psi_+ \{\Delta_a^- V + \Delta^+ V\}.$$

Again, adding the preceding sum with the sum from the definition of the Right Young integral we get

$$\sum_{[a,b)} \psi_+ \{\Delta_a^- V + \Delta^+ V\} + \{\psi \Delta^- V\}(b) - \sum_{(a,b)} \Delta^+ \psi \Delta^- V = \sum_{(a,b]} \psi \Delta^- V + \sum_{[a,b)} \psi_+ \Delta^+ V.$$

This yields the existence of the Right Young integral and relation (3.11). The proof of Lemma 9 is now complete. $\square$

To prove an approximation of an integral with respect to a bracket function, we use the following convergence theorem; its proof is given in [49, Theorem II.15.4].

Lemma 10 *Let g and $\{g_m : m \geq 1\}$ be nondecreasing functions on $[a, b]$ such that for each $a \leq x \leq b$, $g_m(x) \rightarrow g(x)$ as $m \rightarrow \infty$. If $(RRS) \int_a^b h dg$ and $(RRS) \int_a^b h dg_m$, $m \geq 1$, all exist, then*

$$\lim_{m \rightarrow \infty} (RRS) \int_a^b h dg_m = (RRS) \int_a^b h dg.$$

The next approximation theorem extends property (3.3) in the definition of the quadratic λ -variation.

Theorem 11 *Let $\lambda \in \Lambda[a, b]$, let f have the quadratic λ -variation on $[a, b]$, and let h be a regulated function on $[a, b]$. If the right discontinuity points of f are accessible by λ , then*

$$\lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} h(x_{i-1}^m) [f(x_i^m) - f(x_{i-1}^m)]^2 = (LY) \int_a^b h d[f]_\lambda, \quad (3.13)$$

where $\lambda_m = \{x_i^m : i = 0, \dots, n(m)\}$, $m \geq 1$, constitute λ . Also with the same notation, if the left discontinuity points of f are accessible by λ , then

$$\lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} h(x_i^m) [f(x_i^m) - f(x_{i-1}^m)]^2 = (RY) \int_a^b h d[f]_\lambda. \quad (3.14)$$

Proof. To prove (3.13) let $\lambda \in \Lambda[a, b]$ be such that f has the quadratic λ -variation and the right discontinuity points of f are accessible by λ , that is, (1.17) holds. Since the bracket function $[f] := [f]_\lambda$ has bounded variation, the Left Young integral in (3.13) exists by Lemma 9. We show that each sum on the left side of (3.13) is the Left Young integral with respect to a suitable nondecreasing step function, and use Lemma 10 to prove (3.13). For each $m \geq 1$, let F_m be the nondecreasing step function on $[a, b]$ defined by

$$F_m(y) := \begin{cases} 0, & \text{if } y = a \\ \sum_{i=1}^k [f(x_i^m) - f(x_{i-1}^m)]^2, & \text{if } y \in (x_{k-1}^m, x_k^m] \text{ for some } k = 1, \dots, n(m). \end{cases}$$

For each $m \geq 1$ and $1 \leq k \leq n(m)$, the function $(F_m)_+^{(x_k^m)}$ on $[x_{k-1}^m, x_k^m]$ (see (2.37) for the notation) is constant, and $\Delta^+ F_m = 0$ on (x_{k-1}^m, x_k^m) . Thus h is Left Young integrable with respect to F_m on $[x_{k-1}^m, x_k^m]$, and the equality

$$\begin{aligned} (LY) \int_{x_{k-1}^m}^{x_k^m} h dF_m &= (RRS) \int_{x_{k-1}^m}^{x_k^m} h_-^{(x_{k-1}^m)} d(F_m)_+^{(x_k^m)} \\ &\quad + \{h \Delta^+ F_m\}(x_{k-1}^m) + \sum_{(x_{k-1}^m, x_k^m)} \Delta^- h \Delta^+ F_m \\ &= h(x_{k-1}^m) [f(x_k^m) - f(x_{k-1}^m)]^2 \end{aligned}$$

holds for each $m \geq 1$ and $1 \leq k \leq n(m)$. Since the Left Young integral is additive over consecutive intervals (Theorem 2.25), h is Left Young integrable with respect to F_m over $[a, b]$, and the equality

$$\sum_{i=1}^{n(m)} h(x_{i-1}^m) [f(x_i^m) - f(x_{i-1}^m)]^2 = (LY) \int_a^b h dF_m$$

holds for each $m \geq 1$. Therefore to prove (3.13) it is enough to show that

$$\lim_{m \rightarrow \infty} (LY) \int_a^b h dF_m = (LY) \int_a^b h d[f]. \quad (3.15)$$

First, we show that for each $a \leq y \leq b$,

$$\lim_{m \rightarrow \infty} F_m(y) = [f](y). \quad (3.16)$$

Indeed this follows just by the definitions of the functions in the case $y \in \cup_m \lambda_m$. Otherwise, if $y \in [a, b] \setminus \cup_m \lambda_m$, then $x_{k-1}^m < y < x_k^m$ for some $k = k(m)$, and

$$\begin{aligned} \limsup_{m \rightarrow \infty} |F_m(y) - [f](y)| &\leq \limsup_{m \rightarrow \infty} |[f(x_k^m) - f(x_{k-1}^m)]^2 - [f(y) - f(x_{k-1}^m)]^2| \\ &= |[f(y+) - f(y-)]^2 - [f(y) - f(y-)]^2| = 0 \end{aligned}$$

because f is right-continuous at y in this case by (1.17). Therefore (3.16) holds for each $a \leq y \leq b$. Second, in addition to (3.16), for $a \leq y < b$,

$$\lim_{m \rightarrow \infty} F_m(y+) = [f](y+). \quad (3.17)$$

Indeed, if $y \in \cup_m \lambda_m$ then $y = x_k^m$ for all large enough m and some $1 \leq k < n(m)$, so that $F_m(x_k^m+) = F_m(x_k^m) + [f(x_{k+1}^m) - f(x_k^m)]^2 \rightarrow [f](y) + (\Delta^+ f(y))^2 = [f](y+)$ as $m \rightarrow \infty$. If $y \in [a, b] \setminus \cup_m \lambda_m$ then $x_{k-1}^m < y < x_k^m$ and $F_m(y+) = F_m(y) \rightarrow [f](y) = [f](y+)$ as $m \rightarrow \infty$ because f is right-continuous at y in this case by (1.17). Therefore (3.17) holds for each $y \in [a, b]$. This in conjunction with Lemma 10 yields that

$$\lim_{m \rightarrow \infty} (RRS) \int_a^b h_-^{(a)} d(F_m)_+^{(b)} = (RRS) \int_a^b h_-^{(a)} d[f]_+^{(b)}.$$

Since $\lim_{m \rightarrow \infty} \Delta^+ F_m(a) = \Delta^+ [f](a)$, we are left to show that

$$\lim_{m \rightarrow \infty} \sum_{(a,b)} \Delta^- h \Delta^+ F_m = \sum_{(a,b)} \Delta^- h \Delta^+ [f]. \quad (3.18)$$

Given $\epsilon > 0$ choose a finite set $\nu \subset (a, b)$ such that

$$\left| \sum_{(a,b)} \Delta^- h \Delta^+ [f] - \sum_{\nu} \Delta^- h \Delta^+ [f] \right| < \epsilon \quad \text{and} \quad |\Delta^- h(y)| < \epsilon$$

for each $y \notin \nu \cap (\cup_m \lambda_m)$. For each $m \geq 1$, let $\mu_m := \{x_k^m : k = 1, \dots, n(m) - 1\} \setminus \nu$ and $\nu_m := \{x_k^m : k = 1, \dots, n(m) - 1\} \cap \nu$. Then we have

$$\begin{aligned} &\limsup_{m \rightarrow \infty} \left| \sum_{(a,b)} \Delta^- h \Delta^+ F_m - \sum_{(a,b)} \Delta^- h \Delta^+ [f] \right| \\ &\leq \epsilon + \limsup_{m \rightarrow \infty} \left| \sum_{x_k^m \in \mu_m} \Delta^- h(x_k^m) [f(x_{k+1}^m) - f(x_k^m)]^2 \right| \\ &+ \limsup_{m \rightarrow \infty} \left| \sum_{x_k^m \in \nu_m} \Delta^- h(x_k^m) [f(x_{k+1}^m) - f(x_k^m)]^2 - \{\Delta^+ f(x_k^m)\}^2 \right| \leq \epsilon + \epsilon[f](b). \end{aligned}$$

Since ϵ is arbitrary (3.18), and hence (3.15) holds. The proof of (3.13) is complete.

Next we prove (3.14). For each $m \geq 1$, let G_m be a nondecreasing step function on $[a, b]$ defined by

$$G_m(y) := \begin{cases} 0, & \text{if } y \in [a, y_1) \\ \sum_{i=1}^{k-1} [f(x_i^m) - f(x_{i-1}^m)]^2, & \text{if } y \in [x_{k-1}^m, x_k^m) \text{ for some } k = 2, \dots, n(m) \\ \sum_{i=1}^{n(m)} [f(x_i^m) - f(x_{i-1}^m)]^2, & \text{if } y = b. \end{cases}$$

For each $m \geq 1$ and $1 \leq k \leq n(m)$, the function $(G_m)_{-}^{(x_{k-1}^m)}$ on $[x_{k-1}^m, x_k^m]$ (see (2.36) for the notation) is constant, and $\Delta^- G_m = 0$ on (x_{k-1}^m, x_k^m) . Thus h is Right Young integrable with respect to G_m on $[x_{k-1}^m, x_k^m]$, and the equality

$$\begin{aligned} (RY) \int_{x_{k-1}^m}^{x_k^m} h dG_m &= (RRS) \int_{x_{k-1}^m}^{x_k^m} h_+^{(x_k^m)} d(G_m)_{-}^{(x_{k-1}^m)} \\ &\quad - \sum_{(x_{k-1}^m, x_k^m)} \Delta^+ h \Delta^- G_m + (h \Delta^- G_m)(x_k^m) \\ &= h(x_k^m) [f(x_k^m) - f(x_{k-1}^m)]^2 \end{aligned}$$

holds for each $m \geq 1$ and $1 \leq k \leq n(m)$. Since the Right Young integral is additive over consecutive intervals (Theorem 2.25), h is Right Young integrable with respect to G_m over $[a, b]$, and the equality

$$\sum_{i=1}^{n(m)} h(x_i^m) [f(x_i^m) - f(x_{i-1}^m)]^2 = (RY) \int_a^b h dG_m$$

holds for each $m \geq 1$. Therefore to prove (3.14) it is enough to show that

$$\lim_{m \rightarrow \infty} (RY) \int_a^b h dG_m = (RY) \int_a^b h d[f]. \quad (3.19)$$

First, we show that for each $a \leq y \leq b$,

$$\lim_{m \rightarrow \infty} G_m(y) = [f](y). \quad (3.20)$$

Indeed this follows just by the definitions of the functions in the case $y \in \cup_m \lambda_m$. Otherwise, if $y \in [a, b] \setminus \cup_m \lambda_m$, then $x_{k-1}^m < y < x_k^m$ for some $k = k(m)$, and

$$\limsup_{m \rightarrow \infty} |G_m(y) - [f](y)| \leq \limsup_{m \rightarrow \infty} [f(y) - f(x_{k-1}^m)]^2 = (\Delta^- f(y))^2 = 0$$

because f is left-continuity at y in this case by (1.18). Therefore (3.20) holds for each $a \leq y \leq b$. Second, in addition to (3.20), for $a < y \leq b$,

$$\lim_{m \rightarrow \infty} G_m(y-) = [f](y-). \quad (3.21)$$

Indeed, if $y \in \cup_m \lambda_m$ then $y = x_k^m$ for all large enough m and some $1 \leq k < n(m)$, so that $G_m(x_k^m-) = G_m(x_k^m) - [f(x_k^m) - f(x_{k-1}^m)]^2 \rightarrow [f](y) - (\Delta^- f(y))^2 = [f](y-)$ as $m \rightarrow \infty$. If $y \in (a, b] \setminus \cup_m \lambda_m$ then $x_{k-1}^m < y < x_k^m$ and $G_m(y-) = G_m(y) \rightarrow [f](y) = [f](y-)$ as $m \rightarrow \infty$ because f is left-continuity at y in this case by (1.18). Therefore (3.21) holds for each $a \leq y < b$. This in conjunction with Lemma 10 yields that

$$\lim_{m \rightarrow \infty} (RRS) \int_a^b h_+^{(b)} d(G_m)_{-}^{(a)} = (RRS) \int_a^b h_+^{(b)} d[f]_{-}^{(a)}.$$

Since $\lim_{m \rightarrow \infty} \Delta^- G_m(b) = \Delta^- [f](b)$, we are left to show that

$$\lim_{m \rightarrow \infty} \sum_{(a,b)} \Delta^+ h \Delta^- G_m = \sum_{(a,b)} \Delta^+ h \Delta^- [f]. \quad (3.22)$$

Given $\epsilon > 0$ choose a finite set $\nu \subset (a, b)$ such that

$$\left| \sum_{(a,b)} \Delta^+ h \Delta^- [f] - \sum_{\nu} \Delta^+ h \Delta^- [f] \right| < \epsilon \quad \text{and} \quad |\Delta^+ h(y)| < \epsilon$$

for each $y \notin \nu \cap (\cup_m \lambda_m)$. For each $m \geq 1$, let $\mu_m := \{x_k^m : k = 1, \dots, n(m) - 1\} \setminus \nu$ and $\nu_m := \{x_k^m : k = 1, \dots, n(m) - 1\} \cap \nu$. Then we have

$$\begin{aligned} & \limsup_{m \rightarrow \infty} \left| \sum_{(a,b)} \Delta^+ h \Delta^- G_m - \sum_{(a,b)} \Delta^+ h \Delta^- [f] \right| \\ & \leq \epsilon + \limsup_{m \rightarrow \infty} \left| \sum_{x_k^m \in \mu_m} \Delta^+ h(x_k^m) [f(x_k^m) - f(x_{k-1}^m)]^2 \right| \\ & + \limsup_{m \rightarrow \infty} \left| \sum_{x_k^m \in \nu_m} \Delta^+ h(x_k^m) \left[[f(x_k^m) - f(x_{k-1}^m)]^2 - \Delta^- f^2(x_k^m) \right] \right| \leq \epsilon + \epsilon[f](b). \end{aligned}$$

Since ϵ is arbitrary (3.22), and hence (3.19) holds. The proof of Theorem 11 is complete. $\square$

The quadratic λ -covariation. Let f_1 and f_2 be two real-valued functions defined on an interval $[a, b]$. For a partition $\kappa = \{x_i : i = 0, \dots, n\}$ of $[a, b]$, let

$$C(f_1, f_2; \kappa) := \sum_{i=1}^n [f_1(x_i) - f_1(x_{i-1})][f_2(x_i) - f_2(x_{i-1})].$$

Next we extend the notion of the quadratic λ -variation of a single function to a pair of functions by replacing sums s_2 of squares of increments in its definition by sums C just defined.

Definition 12 Let f_1, f_2 be regulated functions on $[a, b]$, and let $\lambda = \{\lambda_m : m \geq 1\} \in \Lambda[a, b]$. We say that the 2-vector function (f_1, f_2) has the *quadratic λ -covariation on $[a, b]$* if there exists an additive upper continuous function $\alpha_\lambda(f_1, f_2)$ on $S[[a, b]]$ such that for each $(u, v) \in S[[a, b]]$

$$\alpha_\lambda(f_1, f_2; u, v) = \lim_{m \rightarrow \infty} C(f_1, f_2; \lambda_m \cap [u, v]).$$

If the 2-vector function (f_1, f_2) has the quadratic λ -covariation on $[a, b]$ then the right distribution function of $\alpha_\lambda(f_1, f_2)$ restricted to $[a, b]$ is called the *bracket function* of (f_1, f_2) , and is denoted by $[f_1, f_2]_\lambda$.

As in the case of the quadratic λ -variation, the quadratic λ -covariation on $[a, b]$ can be defined solely in terms of a regulated function on $[a, b]$. The proof of the following statement is similar to the proof of the Proposition 4, and therefore is omitted.

Proposition 13 Let f_1, f_2 be regulated functions on $[a, b]$, and let $\lambda = \{\lambda_m : m \geq 1\} \in \Lambda[a, b]$. The 2-vector function (f_1, f_2) has the quadratic λ -covariation on $[a, b]$ if and only if there exists a regulated function $B = B_\lambda$ on $[a, b]$ such that $B(a) = 0$ and for each $(s, t) \in S[a, b]$,

$$B(t) - B(s) = \lim_{m \rightarrow \infty} C(f_1, f_2; \lambda_m \cap [s, t]), \quad (3.23)$$

$$\Delta^- B(t) = \{\Delta^- f_1 \Delta^- f_2\}(t) \quad \text{and} \quad \Delta^+ B(s) = \{\Delta^+ f_1 \Delta^+ f_2\}(s). \quad (3.24)$$

If the two statements hold then $[f_1, f_2]_\lambda = B$. Moreover, condition (3.23) for each $(s, t) \in S[a, b]$, can be replaced by the condition

$$B(t) = \lim_{m \rightarrow \infty} C(f_1, f_2; \lambda_m \mathbin{\mathbb{M}} [a, t])$$

for each $t \in [a, b]$, if in addition, λ is such that

$$\{x \in (a, b): (\Delta^- f_1 \Delta^+ f_2 + \Delta^+ f_1 \Delta^- f_2)(x) \neq 0\} \subset \cup_m \lambda_m. \quad (3.25)$$

If in the preceding definition $f_1 = f_2 = f$, then the 2-vector function (f, f) has the quadratic λ -covariation if and only if f has the quadratic λ -variation, and then $[f, f]_\lambda = [f]_\lambda$. In Section 3.3 we extend the notion of the quadratic λ -variation of a real-valued function to a vector valued function. The notion of the quadratic λ -covariation of a 2-vector function is used to define the quadratic λ -variation of a d -vector valued function with $d \geq 2$.

For notation simplicity let $\mathcal{W}_\infty^*[a, b] := \mathcal{R}[a, b]$.

Proposition 14 For $p^{-1} + q^{-1} = 1$ and $p \geq 1$, let $f_1 \in \mathcal{W}_p[a, b]$ and $f_2 \in \mathcal{W}_q^*[a, b]$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ be such that (3.25) holds. Then the 2-vector function (f_1, f_2) has the quadratic λ -covariation on $[a, b]$, and its bracket function $[f_1, f_2]_\lambda$ is a pure jump function given by

$$[f_1, f_2]_\lambda(y) = \sum_{(a, y]} \Delta^- f_1 \Delta^- f_2 + \sum_{(a, y]} \Delta^+ f_1 \Delta^+ f_2, \quad \text{for } a \leq y \leq b. \quad (3.26)$$

Proof. By Hölder's inequality, the two sums $\sum_{(a, b]} \Delta^- f_1 \Delta^- f_2$ and $\sum_{(a, b]} \Delta^+ f_1 \Delta^+ f_2$ converge absolutely. Therefore the function on the right side of (3.26) is of bounded variation, and its jumps agree with those of the bracket function of (f_1, f_2) if it exists (cf. (3.24)). Let $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$ for each $m \geq 1$, and let $a \leq u < v \leq b$. We have to prove that

$$\lim_{m \rightarrow \infty} C(f_1, f_2; \lambda_m \mathbin{\mathbb{M}} [u, v]) = \sum_{(u, v]} \Delta^- f_1 \Delta^- f_2 + \sum_{(u, v]} \Delta^+ f_1 \Delta^+ f_2. \quad (3.27)$$

Let $\epsilon > 0$. There exists a partition $\kappa = \{z_k: k = 0, \dots, l\}$ of $[u, v]$ such that

$$\left| \sum_{\kappa \cap (u, v]} \Delta^- f_1 \Delta^- f_2 + \sum_{\kappa \cap (u, v]} \Delta^+ f_1 \Delta^+ f_2 - \sum_{(u, v]} \Delta^- f_1 \Delta^- f_2 - \sum_{(u, v]} \Delta^+ f_1 \Delta^+ f_2 \right| < \epsilon$$

and either $\max_{1 \leq k \leq l} \text{Osc}(f_2; (z_{k-1}, z_k)) < \epsilon/v_1(f_1)$ if $p = 1$, or

$$\sum_{k=1}^l v_q(f_2; (z_{k-1}, z_k)) < \epsilon^q/v_p(f_1)^{q/p} \quad \text{if } p > 1.$$

The last relation holds by Lemma 2.9. Let $I := \{k = 1, \dots, l-1: z_k \in \cup_m \lambda_m\}$. If $k \in I$ then $z_k = x_{i(k)}^m$ for some $i(k) \in \{0, \dots, n(m)\}$ and for all sufficiently large m . Thus for $k \in I$, we have

$$\lim_{m \rightarrow \infty} [\Delta_{i(k)}^m f_1 \Delta_{i(k)}^m f_2 + \Delta_{i(k)+1}^m f_1 \Delta_{i(k)+1}^m f_2] = (\Delta^- f_1 \Delta^- f_2)(z_k) + (\Delta^+ f_1 \Delta^+ f_2)(z_k),$$

where $\Delta_j^m \rho := \rho(x_j^m) - \rho(x_{j-1}^m)$ if x_j^m is defined, and $:= 0$ otherwise. If $k \in \{1, \dots, l-1\} \setminus I$ then $x_{i(k)-1}^m < z_k < x_{i(k)}^m$ for some $i(k) \in \{1, \dots, n(m)\}$ and for all sufficiently large m . By assumption (3.25), it follows that

$$[f_1(z_k+) - f_1(z_k-)][f_2(z_k+) - f_2(z_k-)] = (\Delta^- f_1 \Delta^- f_2)(z_k) + (\Delta^+ f_1 \Delta^+ f_2)(z_k).$$

Thus for $k \in \{1, \dots, l-1\} \setminus I$, we have

$$\lim_{m \rightarrow \infty} \Delta_{i(k)}^m f_1 \Delta_{i(k)}^m f_2 = (\Delta^- f_1 \Delta^- f_2)(z_k) + (\Delta^+ f_1 \Delta^+ f_2)(z_k).$$

For $k = 0$ and $k = l$, that is at the endpoints of $[u, v]$, recalling notation $\lambda_m \cap [u, v] = \{u, x_{i(u)}^m, \dots, x_{i(v)}^m, v\}$, we have

$$\lim_{m \rightarrow \infty} [f_1(x_{i(u)}^m) - f_1(u)][f_2(x_{i(u)}^m) - f_2(u)] = (\Delta^+ f_1 \Delta^+ f_2)(u)$$

and

$$\lim_{m \rightarrow \infty} [f_1(v) - f_1(x_{i(v)}^m)][f_2(v) - f_2(x_{i(v)}^m)] = (\Delta^- f_1 \Delta^- f_2)(v).$$

Let m_0 be the first integer such that $\{z_k: k \in I\} \subset \lambda_{m_0}$ and each pair (z_{k-1}, z_k) of points from κ is separated by at least two points from λ_{m_0} . Let $x_{i(u)-1} := u$ and $x_{i(v)+1} := v$. For $m \geq m_0$, let $I_1(m)$ be the set of all indices $i(k), i(k)+1$ with $k \in I$, all indices $i(k)$ with $k \in \{1, \dots, l-1\} \setminus I$, and the indices $i(u), i(v)+1$. Then we have

$$\lim_{m \rightarrow \infty} \sum_{i \in I_1(m)} \Delta_i^m f_1 \Delta_i^m f_2 = \sum_{\kappa \cap (u, v]} \Delta^- f_1 \Delta^- f_2 + \sum_{\kappa \cap [u, v)} \Delta^+ f_1 \Delta^+ f_2.$$

For $m \geq m_0$, let $I_2(m) := \{i(u), \dots, i(v)+1\} \setminus I_1(m)$. Then by Hölder's inequality, we have

$$\left| \sum_{i \in I_2(m)} \Delta_i^m f_1 \Delta_i^m f_2 \right| \leq \left(\sum_{k=1}^l v_q(f_2; (z_{k-1}, z_k)) \right)^{1/q} v_p(f_1)^{1/p} < \epsilon$$

for all $m \geq m_0$ if $p > 1$. If $p = 1$ then we simply have the bound

$$\left| \sum_{i \in I_2(m)} \Delta_i^m f_1 \Delta_i^m f_2 \right| \leq \max_{1 \leq k \leq l} \text{Osc}(f_2; (z_{k-1}, z_k)) v_1(f_1) < \epsilon$$

for all $m \geq m_0$. Therefore

$$\limsup_{m \rightarrow \infty} \left| C(f_1, f_2; \lambda_m \cap [u, v]) - \sum_{(a, y]} \Delta^- f_1 \Delta^- f_2 - \sum_{[a, y)} \Delta^+ f_1 \Delta^+ f_2 \right| \leq 2\epsilon.$$

Since $\epsilon > 0$ is arbitrary (3.27) holds, proving Proposition 14. $\square$

In Proposition 14 taking $f_1 = f_2$ we get:

Corollary 15 *Let $\lambda \in \Lambda[a, b]$. If f has the local 2-variation on $[a, b]$ and two-sided discontinuity points of f are accessible by λ , then f has the quadratic λ -variation, and its bracket function $[f]_\lambda$ is a pure jump function given by*

$$[f]_\lambda(y) = \sum_{(a, y]} (\Delta^- f)^2 + \sum_{[a, y)} (\Delta^+ f)^2, \quad \text{for } a \leq y \leq b.$$

For the next corollary recall Definition 1.2 of the class $\mathcal{D}_{\lambda, p}$ of (λ, p) -decomposable functions.

Corollary 16 *Let $f \in \mathcal{D}_{\lambda, p}[a, b]$ for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$. If $(g, h) \in D_{\lambda, p}^+(f) \cup D_{\lambda, p}^-(f) \neq \emptyset$ then f has the quadratic λ -variation, and its bracket function is given by*

$$[f]_\lambda(x) = [g]_\lambda^c(x) + \sum_{(a, x]} \{\Delta^- f\}^2 + \sum_{[a, x)} \{\Delta^+ f\}^2, \quad \text{for } a \leq x \leq b. \quad (3.28)$$

Proof. Let $\lambda = \{\lambda_m: m \geq 1\}$ and $1 \leq p < 2$ be such that $f \equiv C + g + h$ for some $(g, h) \in D_{\lambda, p}^+(f)$. Since

$$\{x \in (a, b): (\Delta^- h \Delta^+ h)(x) \neq 0\} \subset \{x \in (a, b): (\Delta^+ h)(x) \neq 0\} \subset \cup_m \lambda_m, \quad (3.29)$$

by the preceding corollary, h has the quadratic λ -variation with the bracket function

$$[h]_\lambda(x) = \sum_{(a, x]} \{\Delta^- h\}^2 + \sum_{[a, x)} \{\Delta^+ h\}^2, \quad \text{for } a \leq x \leq b.$$

Since $g \in \mathcal{W}_q^*[a, b]$ with $p^{-1} + q^{-1} = 1$, and

$$\begin{aligned} & \{x \in (a, b): (\Delta^- g \Delta^+ h + \Delta^+ g \Delta^- h)(x) \neq 0\} \\ & \subset \{x \in (a, b): (\Delta^+ g)(x) \neq 0\} \cup \{x \in (a, b): (\Delta^+ h)(x) \neq 0\} \subset \cup_m \lambda_m, \end{aligned} \quad (3.30)$$

by Proposition 14 applied to $f_1 = g$ and $f_2 = h$, the 2-vector function (g, h) has the quadratic λ -covariation with the bracket function

$$[g, h]_\lambda(x) = \sum_{(a, x]} \Delta^- g \Delta^- h + \sum_{[a, x)} \Delta^+ g \Delta^+ h, \quad \text{for } a \leq x \leq b.$$

Let H be the function on $[a, b]$ defined by the right side of (3.28). Since g has the quadratic λ -variation, it follows that for $(s, t) \in S[a, b]$,

$$\begin{aligned} & \lim_{m \rightarrow \infty} s_2(f; \lambda_m \cap [s, t]) \\ &= [g]_\lambda(t) - [g]_\lambda(s) + 2 \left\{ \sum_{(s, t]} \Delta^- g \Delta^- h + \sum_{[s, t)} \Delta^+ g \Delta^+ h \right\} + \sum_{(s, t]} \{\Delta^- h\}^2 + \sum_{[s, t)} \{\Delta^+ h\}^2 \\ &= [g]_\lambda^c(t) - [g]_\lambda^c(s) + \sum_{(s, t]} \{\Delta^-(g + h)\}^2 + \sum_{[s, t)} \{\Delta^+(g + h)\}^2 = H(t) - H(s). \end{aligned}$$

Also since $\Delta^- H(t) = \{\Delta^- f(t)\}^2$ for $a < t \leq b$, and $\Delta^+ H(s) = \{\Delta^+ f(s)\}^2$ for $a \leq s < b$, by Proposition 4, f has the quadratic λ -variation with the bracket function (3.28) in the case $(g, h) \in D_{\lambda}^+(f) \neq \emptyset$. If $(g, h) \in D_{\lambda}^-(f) \neq \emptyset$ then the proof is the same because (3.29) and (3.30) hold in this case with Δ^+ in the middle terms replaced by Δ^- . The proof of Corollary 16 is complete. $\square$

In (3.28) we have that $[g]_\lambda^c$ does not depend on a (λ, p) -decomposition $f \equiv C + g + h$ such that $(g, h) \in D_{\lambda, p}^+(f) \cup D_{\lambda, p}^-(f)$. Therefore for $f \in \mathcal{D}_{\lambda, p}[a, b]$ with $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, such that $D_{\lambda, p}^+(f) \cup D_{\lambda, p}^-(f) \neq \emptyset$, the continuous part of the bracket function $[f]_\lambda$ is uniquely defined.

3.2 The Left and Right λ -integrals

In this section we define two integrals with respect to a (λ, p) -decomposable function. Let $f \equiv C + g + h$, where (g, h) is a (λ, p) -dual pair for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$. The two integrals with respect to f are defined to be a sum of an integral with respect to g and an integral with respect to h . Since h has bounded p -variation with $1 \leq p < 2$, we use the Left Young and Right Young integrals to integrate with respect to h . To integrate with respect to a function having the quadratic λ -variation, we introduce the Left Cauchy and Right Cauchy λ -integrals which are extensions of the Left Young and Right Young integrals, respectively.

The Left and Right Cauchy λ -integrals. Let ϕ and g be real-valued functions defined on a closed interval $[a, b]$. For a partition $\kappa = \{x_i: i = 0, \dots, n\}$ of $[a, b]$, let

$$S_{LC}(\phi, g; \kappa) := \sum_{i=1}^n \phi(x_{i-1})[g(x_i) - g(x_{i-1})] \quad \text{and} \quad S_{RC}(\phi, g; \kappa) := \sum_{i=1}^n \phi(x_i)[g(x_i) - g(x_{i-1})].$$

Definition 17 Let g be a regulated function on $[a, b]$, let ϕ be a function on $[a, b]$ such that a left limit $\phi(t-)$ exists at $t \in (a, b]$ whenever $\Delta^-g(t) \neq 0$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. We say that the *Left Cauchy λ -integral (LC) $\int \phi d_\lambda g$ is defined on $[a, b]$* if there exists a regulated function Φ on $[a, b]$ such that $\Phi(a) = 0$ and for each $a \leq u < v \leq b$,

$$\Phi(v) - \Phi(u) = \lim_{m \rightarrow \infty} S_{LC}(\phi, g; \lambda_m \cap [u, v]), \quad (3.31)$$

$$\Delta^- \Phi(v) = \begin{cases} (\phi \Delta^- g)(v) & \text{if } (\Delta^- g)(v) \neq 0, \\ 0 & \text{if } (\Delta^- g)(v) = 0 \end{cases} \quad \text{and} \quad \Delta^+ \Phi(u) = (\phi \Delta^+ g)(u). \quad (3.32)$$

If a regulated function Φ on $[a, b]$ exists and satisfies the same conditions except that (3.31) and (3.32) do not hold for $v = b$, then we say that the *Left Cauchy λ -integral (LC) $\int \phi d_\lambda g$ is defined on $[a, b)$* .

In Theorem 5.13 below the Left Cauchy λ -integral (LC) $\int \phi d_\lambda g$ on $[a, b]$ is used when g is a suitable continuous function on $[a, b]$ and ϕ is a function on $[a, b]$ having no the left limit at the right-endpoint b . In other places we use this integral when both the integrator and integrand are regulated in which case a simple characterization of the integral holds. But first we define the Right Cauchy λ -integral.

Definition 18 Let g be a regulated function on $[a, b]$, let ϕ be a function on $[a, b]$ such that a right limit $\phi(t+)$ exists at $t \in [a, b)$ whenever $\Delta^+g(t) \neq 0$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. We say that the *Right Cauchy λ -integral (RC) $\int \phi d_\lambda g$ on $[a, b]$* exists if there exists a regulated function Ψ on $[a, b]$ such that $\Psi(a) = 0$ and for each $a \leq u < v \leq b$,

$$\Psi(v) - \Psi(u) = \lim_{m \rightarrow \infty} S_{RC}(\phi, g; \lambda_m \cap [u, v]),$$

$$\Delta^- \Psi(v) = (\phi \Delta^- g)(v) \quad \text{and} \quad \Delta^+ \Psi(u) = \begin{cases} (\phi \Delta^+ g)(u) & \text{if } (\Delta^+ g)(u) \neq 0, \\ 0 & \text{if } (\Delta^+ g)(u) = 0. \end{cases}$$

The Left Cauchy and Right Cauchy λ -integrals, if exist, are additive functions on $S[a, b]$. For notation simplicity, the following is proved under stronger assumptions than necessary.

Proposition 19 Let g, ϕ be regulated functions on $[a, b]$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. The Left Cauchy λ -integral (LC) $\int \phi d_\lambda g$ is defined on $[a, b]$ if and only if there exists an additive upper continuous function μ_L on $S[[a, b]]$ such that for each $(u, v) \in S[[a, b]]$,

$$\mu_L(u, v) = \lim_{m \rightarrow \infty} S_{LC}(\phi, g; \lambda_m \cap [[u, v]]). \quad (3.33)$$

Also, the Right Cauchy λ -integral (RC) $\int \phi d_\lambda g$ is defined on $[a, b]$ if and only if there exists an additive upper continuous function μ_R on $S[[a, b]]$ such that for each $(u, v) \in S[[a, b]]$,

$$\mu_R(u, v) = \lim_{m \rightarrow \infty} S_{RC}(\phi, g; \lambda_m \cap [[u, v]]).$$

Proof. We prove only the first part of the proposition because the proof of the second part is symmetric. First suppose that there exists a regulated function Φ on $[a, b]$, equal zero at a and satisfying (3.31) and (3.32). Define a function μ_L on $S[[a, b]]$ by $\mu_L(u, v) := \Phi(v) - \Phi(u)$ for $(u, v) \in S[[a, b]]$. Then μ_L is additive and upper continuous function on $S[[a, b]]$ by Theorem 2.4. We prove (3.33) only for the cases $(u, v) = (a, t-)$ and $(u, v) = (a, t+)$ because the proofs for the other cases are similar. To this aim let $\lambda_m = \{x_i^m : i = 0, \dots, n(m)\}$, $m = 1, 2, \dots$, and first let $t \in (a, b]$. For each $m \geq 1$, there is an index $i(t) = i_m(t) \in \{1, \dots, n(m) - 1\}$ such that $x_{i(t)}^m < t \leq x_{i(t)+1}^m$. Then by (3.31) and (3.32), we have

$$\begin{aligned} & \lim_{m \rightarrow \infty} S_{LC}(\phi, g; \lambda_m \cap [a, t-]) \\ &= \lim_{m \rightarrow \infty} \left\{ S_{LC}(\phi, g; \lambda_m \cap [a, t]) - \phi(x_{i(t)}^m)[g(t) - g(x_{i(t)}^m)] + \phi(x_{i(t)}^m)[g(t-) - g(x_{i(t)}^m)] \right\} \\ &= \Phi(t) - \phi(t-) \Delta^- g(t) = \Phi(t-) = \mu_L(a, t-), \end{aligned}$$

proving (3.33) for $(u, v) = (a, t-) \in S[[a, b]]$. Now let $t \in [a, b)$. By the definition of the trace partition (Notation 2), we have for each $m \geq 1$,

$$S_{LC}(\phi, g; \lambda_m \cap [a, t+]) = S_{LC}(\phi, g; \lambda_m \cap [a, t]) + (\phi \Delta^+ g)(t).$$

Letting $m \rightarrow \infty$, it follows that (3.33) holds for $(u, v) = (a, t+) \in S[[a, b]]$.

To prove the converse implication, let Φ be the restriction to $[a, b]$ of the right distribution function of μ_L , that is, $\Phi(t) := \mu_L(a, t)$ for $t \in [a, b]$. By statement (iv) of Theorem 2.4 and (3.33), Φ is regulated on $[a, b]$, is zero at a , and satisfies properties (3.31), (3.32). The proof of Proposition 19 is complete. $\square$

The next property identifies a difference between values of the Left Cauchy and Right Cauchy λ -integrals, provided the integrand and integrator possess the quadratic λ -covariation.

Proposition 20 *Let a 2-vector function (f_1, f_2) on $[a, b]$ have the quadratic λ -covariation for some $\lambda \in \Lambda[a, b]$. The λ -integral $(LC) \int f_1 d_\lambda f_2$ on $[a, b]$ exists if and only if so does the λ -integral $(RC) \int f_1 d_\lambda f_2$ on $[a, b]$, and if the two λ -integrals exist then for $a \leq x \leq b$,*

$$[f_1, f_2]_\lambda(x) = (RC) \int_a^x f_1 d_\lambda f_2 - (LC) \int_a^x f_1 d_\lambda f_2.$$

Proof. The conclusion follows from the relation $C(f_1, f_2; \kappa) = S_{RC}(f_1, f_2; \kappa) - S_{LC}(f_1, f_2; \kappa)$, valid for any partition κ . $\square$

The following extends an integration by parts formula for the Left Cauchy and Right Cauchy λ -integrals, provided the integrand and integrator possess the quadratic λ -covariation.

Proposition 21 *For $\lambda \in \Lambda[a, b]$, let a 2-vector function (f_1, f_2) on $[a, b]$ have the quadratic λ -covariation. For $\sharp$ equal to LC or RC , the λ -integral $(\sharp) \int f_1 d_\lambda f_2$ on $[a, b]$ exists if and only if so does the integral $(\sharp) \int f_2 d_\lambda f_1$ on $[a, b]$, and if the four λ -integrals exist then for $a \leq x \leq b$*

$$\begin{aligned} [f_1, f_2]_\lambda(x) &= (f_1 f_2)(x) - (f_1 f_2)(a) - (LC) \int_a^x f_1 d_\lambda f_2 - (LC) \int_a^x f_2 d_\lambda f_1 \\ &= - \left[(f_1 f_2)(x) - (f_1 f_2)(a) - (RC) \int_a^x f_1 d_\lambda f_2 - (RC) \int_a^x f_2 d_\lambda f_1 \right]. \end{aligned}$$

Proof. The conclusion follows from the relations

$$\begin{aligned} C(f_1, f_2; \kappa) &= (f_1 f_2)(x_n) - (f_1 f_2)(x_0) - S_{LC}(f_1, f_2; \kappa) - S_{LC}(f_1, f_2; \kappa) \\ &= -\left[(f_1 f_2)(x_n) - (f_1 f_2)(x_0) - S_{RC}(f_1, f_2; \kappa) - S_{RC}(f_1, f_2; \kappa)\right], \end{aligned}$$

valid for any partition $\kappa = \{x_i: i = 0, \dots, n\}$. $\square$

The Left and Right λ -integrals. For $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, let f be a (λ, p) -decomposable function on $[a, b]$, that is, $f \equiv C + g + h$ for a (λ, p) -dual pair (g, h) . Next two integrals with respect to f are defined to be a sum of two integrals with respect to g and h , respectively. Since the component h is less erratic than the component g , one may expect that an integral with respect to h may exist in a stronger sense for a large enough class of integrands. Therefore we use the Left (Right) Young integral to integrate with respect to h . Under the conditions of Theorem 2.22, the Left (Right) Young integral exists and equals to the Left (Right) Cauchy λ -integral for any $\lambda \in \Lambda[a, b]$ such that the right (left) discontinuity points of h are accessible by λ . This type of independence on a sequence of partitions λ makes the Left (Right) Young integral preferable whenever it exists. Moreover, any result concerning the Left (Right) Cauchy λ -integral with respect to such a function is an *extension* of the corresponding result concerning the Left (Right) Young integral. Since a (λ, p) -decomposition of an integrator may be non-unique we have to insure that the sum of the two integrals with respect to different decompositions give the same value.

Definition 22 Let F be a regulated function on $[a, b]$, let $f \in \mathcal{D}_{\lambda, p}[a, b]$ for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, and let $A \subset D_{\lambda, p}(f)$.

I. We say that the *Left λ -integral* $(L) \int F d_\lambda f$ on $[a, b]$ is defined with respect to A if for each $(g, h) \in A$ the sum $(LC) \int F d_\lambda g + (LY) \int F dh$ is defined on $[a, b]$ and its values for each $(u, v) \in S[a, b]$ do not depend on $(g, h) \in A$. If the Left λ -integral $(L) \int F d_\lambda f$ is defined on $[a, b]$ with respect to A then let

$$(L) \int_u^v F d_\lambda f := (LC) \int_u^v F d_\lambda g + (LY) \int_u^v F dh$$

for $(u, v) \in S[a, b]$ and for any $(g, h) \in A$.

II. We say that the *Right λ -integral* $(R) \int F d_\lambda f$ on $[a, b]$ is defined with respect to A if for each $(g, h) \in A$ the sum $(RC) \int F d_\lambda g + (RY) \int F dh$ is defined on $[a, b]$ and its values for each $(u, v) \in S[a, b]$ do not depend on $(g, h) \in A$. If the Right λ -integral $(R) \int F d_\lambda f$ is defined on $[a, b]$ with respect to A then let

$$(R) \int_u^v F d_\lambda f := (RC) \int_u^v F d_\lambda g + (RY) \int_u^v F dh$$

for $(u, v) \in S[a, b]$ and for any $(g, h) \in A$.

Under the assumptions of the following proposition, the Left and Right λ -integrals with respect to a set A are defined whenever they are defined for a single element of A .

Proposition 23 For $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, let f be a (λ, p) -decomposable function on $[a, b]$, and let $F \in \text{dual}(\mathcal{W}_p)[a, b]$.

- (a) If $(LC) \int F d_\lambda g$ is defined on $[a, b]$ for some $(g, h) \in D_{\lambda, p}^+(f)$ then $(L) \int F d_\lambda f$ is defined on $[a, b]$ with respect to $D_{\lambda, p}^+(f)$.
- (b) If $(RC) \int F d_\lambda g$ is defined on $[a, b]$ for some $(g, h) \in D_{\lambda, p}^-(f)$ then $(R) \int F d_\lambda f$ is defined on $[a, b]$ with respect to $D_{\lambda, p}^-(f)$.

Proof. We prove only statement (a) because a proof of statement (b) is symmetric. If the set $D_{\lambda, p}^+(f)$ has a single element then there is nothing to prove. Suppose that there are two different elements (g_1, h_1) and (g_2, h_2) in $D_{\lambda, p}^+(f)$. By the (λ, p) -decomposability of f , $g_1 - g_2 = -(h_1 - h_2) \in \mathcal{W}_p$. In the case $1 < p < 2$, $F \in \mathcal{W}_q$ with $p^{-1} + q^{-1} > 1$, and the Left Young integral of F with respect to $g_1 - g_2$ exists by Theorem 2.24. While in the case $p = 1$, this integral exists by Lemma 9. Since the λ -integrals $(LC) \int F d_\lambda g_1$ and $(LC) \int F d_\lambda g_2$ are defined, it follows that $(LC) \int F d_\lambda (g_1 - g_2)$ exists and equals to the difference of the two λ -integrals. If $\Delta^+(g_1 - g_2)(x) \neq 0$ for some $x \in [a, b]$ then either $\Delta^+g_1(x) \neq 0$ or $\Delta^+g_2(x) \neq 0$, and so $x \in \cup_m \lambda_m$. By Theorem 2.22, we then have

$$\begin{aligned} (LC) \int F d_\lambda g_1 - (LC) \int F d_\lambda g_2 &= (LC) \int F d_\lambda (g_1 - g_2) \\ &= (LY) \int F d(g_1 - g_2) = (LY) \int F dh_2 - (LY) \int F dh_1. \end{aligned}$$

Thus the sum $(LC) \int F d_\lambda g + (LY) \int F dh$ does not depend on whether $(g, h) = (g_1, h_1)$ or $(g, h) = (g_2, h_2)$. The proof of Proposition 23 is complete. $\square$

Thus existence of the Left and Right λ -integrals essentially reduces to the problem of existence of the Left and Right Cauchy λ -integrals.

Chain rules. Next we turn to establishing sufficient conditions for existence of the λ -integrals $(L) \int (\psi \circ f) d_\lambda f$ and $(R) \int (\psi \circ f) d_\lambda f$ when f has the quadratic λ -variation and ψ is a smooth function. Also, we show that each such λ -integral satisfies a relation to be called a *chain rule*.

Theorem 24 *Let $f \in \mathcal{D}_{\lambda, p}[a, b]$ for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, and let ϕ be a C^2 class function on an open set containing the range of f . Then the following two statements hold:*

- (a) If $D_{\lambda, p}^+(f) \neq \emptyset$ then the Left λ -integral $(L) \int (\phi' \circ f) d_\lambda f$ is defined on $[a, b]$ with respect to $D_{\lambda, p}^+(f)$, and for $a \leq z < y \leq b$,

$$\begin{aligned} (\phi \circ f)(y) &= (\phi \circ f)(z) + (L) \int_z^y (\phi' \circ f) d_\lambda f + \frac{1}{2} (RS) \int_z^y (\phi'' \circ f) d[f]_\lambda^c \\ &\quad + \sum_{(z, y]} \left\{ \Delta^-(\phi \circ f) - (\phi' \circ f)_- \Delta^- f \right\} + \sum_{[z, y)} \left\{ \Delta^+(\phi \circ f) - (\phi' \circ f)_+ \Delta^+ f \right\}, \end{aligned} \quad (3.34)$$

where the two sums converge unconditionally.

- (b) If $D_{\lambda, p}^-(f) \neq \emptyset$, then the Right λ -integral $(R) \int (\phi' \circ f) d_\lambda f$ is defined on $[a, b]$ with respect to $D_{\lambda, p}^-(f)$, and for $a \leq z < y \leq b$,

$$\begin{aligned} (\phi \circ f)(y) &= (\phi \circ f)(z) + (R) \int_z^y (\phi' \circ f) d_\lambda f - \frac{1}{2} (RS) \int_z^y (\phi'' \circ f) d[f]_\lambda^c \\ &\quad + \sum_{(z, y]} \left\{ \Delta^-(\phi \circ f) - (\phi' \circ f)_- \Delta^- f \right\} + \sum_{[z, y)} \left\{ \Delta^+(\phi \circ f) - (\phi' \circ f)_+ \Delta^+ f \right\}, \end{aligned} \quad (3.35)$$

where the two sums converge unconditionally.

Proof. We prove statement (a) and indicate necessary changes needed to prove statement (b). Let $f \equiv C + g + h$ for some $(g, h) \in D_{\lambda, p}^+(f)$. Then g has the quadratic λ -variation and has bounded q -variation for some $2 < q < p/(p-1)$ if $1 < p < 2$. In the case $1 < p < 2$, the function f has the same q -variation property as the function g because $p < 2 < q$. Since ϕ' is Lipschitz function on the range of f , the composition $\phi' \circ f \in \mathcal{W}_q[a, b]$ if $1 < p < 2$, and it is a regulated function on $[a, b]$ if $p = 1$. Thus the Left Young integral $(LY) \int_a^b (\phi' \circ f) dh$ exists by Theorem 2.22 in the case $1 < p < 2$ and by Lemma 9 in the case $p = 1$. By Proposition 23 with $A = D_{\lambda, p}^+(f)$, it is enough to prove that the Left Cauchy λ -integral $(LC) \int (\phi' \circ f) d\lambda g$ on $[a, b]$ is defined and relation (3.34) holds for each $a \leq z < y \leq b$. In fact, it is enough to prove that for each $a \leq z < y \leq b$, the limit

$$(LC) \int_z^y (\phi' \circ f) d\lambda g := \lim_{m \rightarrow \infty} S_{LC}(\phi' \circ f, g; \lambda_m \cap [z, y]) \quad (3.36)$$

exists and relation (3.34) holds. Indeed, let $\Phi(y) := (LC) \int_a^y (\phi' \circ f) d\lambda g$ for $a \leq y \leq b$. The Left Young integral and the Riemann-Stieltjes integral are additive over adjacent intervals (Theorem 2.25). While the unconditional sums are additive over disjoint intervals (Theorem A.7). Therefore using (3.34), it follows that $\Phi(y) - \Phi(z) = (LC) \int_z^y (\phi' \circ f) d\lambda g$ for $a \leq z < y \leq b$, that is, condition (3.31) of Definition 17 holds. To show that condition (3.32) holds, for $a \leq y < x \leq b$, we have

$$\begin{aligned} \Phi(x) - \Phi(y) &= (LC) \int_y^x (\phi' \circ f) d\lambda g = [(\phi \circ f)(x) - (\phi \circ f)(y)] \\ &\quad - (LY) \int_x^y (\phi' \circ f) dh - \frac{1}{2} (RS) \int_x^y (\phi'' \circ f) d[f]_{\lambda}^c \\ &\quad - \sum_{(x, y]} \left\{ \Delta^-(\phi \circ f) - (\phi' \circ f)_- \Delta^- f \right\} - \sum_{[x, y)} \left\{ \Delta^+(\phi \circ f) - (\phi' \circ f) \Delta^+ f \right\} \\ &\stackrel{y \uparrow x}{=} \Delta^-(\phi \circ f)(x) - \{(\phi' \circ f)_- \Delta^- h\}(x) - \{\Delta^-(\phi \circ f) - (\phi' \circ f)_- \Delta^- f\}(x) \\ &= (\phi' \circ f)_-(x) \Delta^- g(x). \end{aligned} \quad (3.37)$$

For the last step we used Proposition 2.27. Likewise for $a \leq x < y \leq b$, we have

$$\begin{aligned} \Phi(y) - \Phi(x) &\stackrel{y \downarrow x}{=} \Delta^+(\phi \circ f)(x) - \{(\phi' \circ f) \Delta^+ h\}(x) - \{\Delta^+(\phi \circ f) - (\phi' \circ f) \Delta^+ f\}(x) \\ &= (\phi' \circ f)(x) \Delta^+ g(x). \end{aligned}$$

Therefore Φ is a regulated function on $[a, b]$, is equal zero at a , and has jumps satisfying the relations

$$\Delta^- \Phi = (\phi' \circ f)_- \Delta^- g \quad \text{and} \quad \Delta^+ \Phi = (\phi' \circ f) \Delta^+ g. \quad (3.38)$$

That is, (3.32) of Definition 17 holds. Hence statement (a) of the theorem will be proved once we show that for each $a \leq z < y \leq b$, the limit (3.36) exists and relation (3.34) holds.

For notation simplicity, the existence of the limit (3.36) and relation (3.34) will be proved when $z = a$ and $y = b$. Let $A_1 := (LY) \int_a^b (\phi' \circ f) dh$. By Lemma 9, there exists the Riemann-Stieltjes integral $A_2 := (RS) \int_a^b (\phi'' \circ f) d[f]_{\lambda}^c$. By the mean value theorem, for any $y \in (a, b]$,

$$|\Delta^-(\phi \circ f)(y) - (\phi' \circ f)(y-) \Delta^- f(y)| \leq K[\Delta^- f(y)]^2,$$

where K is the Lipschitz constant of ϕ' on the range of f . Thus the first sum in (3.34) converges unconditionally to a value A_3 , because $\sum_{(a,b]} [\Delta^- f]^2 < \infty$. Likewise, it follows that the second sum in (3.34) converges unconditionally to a value A_4 . For each $m \geq 1$, let $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$, and let $\Delta_i^m \rho := \rho(x_i^m) - \rho(x_{i-1}^m)$, $i = 1, \dots, n(m)$, for a function ρ on $[a, b]$. Let $\epsilon > 0$. Since $(g, h) \in D_{\lambda, p}^+(f)$, and so $\{x \in (a, b): \Delta^+ h(x) \neq 0\} \subset \cup_m \lambda_m$, by Theorem 2.22, there exists an integer m_1 such that for all $m \geq m_1$,

$$\left| \sum_{i=1}^{n(m)} \phi'(f(x_{i-1}^m)) \Delta_i^m h - A_1 \right| < \epsilon. \quad (3.39)$$

Since $\{x \in (a, b): \Delta^+ g(x) \neq 0\} \subset \cup_m \lambda_m$, by Theorem 11 and Lemma 9, there exists an integer m_2 such that for all $m \geq m_2$,

$$\left| \sum_{i=1}^{n(m)} \phi''(f(x_{i-1}^m)) \{\Delta_i^m g\}^2 - A_2 - A_5 - A_6 \right| < \epsilon, \quad (3.40)$$

where

$$A_5 := \sum_{(a,b]} (\phi'' \circ f)_- \{\Delta^- g\}^2 \quad \text{and} \quad A_6 := \sum_{[a,b)} (\phi'' \circ f) \{\Delta^+ g\}^2.$$

Since ϕ'' is uniformly continuous on the range of f , there is a $\delta > 0$ such that $|\phi''(u) - \phi''(v)| < \epsilon$ whenever $|u - v| < \delta$ and $|u| \vee |v| \leq \|f\|_\infty$. Therefore and recalling Lemma 2.9, we conclude that there exists a partition $\mu = \{y_j: j = 0, \dots, k\}$ of $[a, b]$ such that for any $\nu \supset \mu$,

$$\left| \sum_{\nu \setminus \{a\}} \left\{ \Delta^-(\phi \circ f) - (\phi' \circ f)_- \Delta^- f \right\} - A_3 \right| \vee \left| \sum_{\nu \setminus \{a\}} (\phi'' \circ f)_- \{\Delta^- g\}^2 - A_5 \right| < \epsilon, \quad (3.41)$$

$$\left| \sum_{\nu \setminus \{b\}} \left\{ \Delta^+(\phi \circ f) - (\phi' \circ f) \Delta^+ f \right\} - A_4 \right| \vee \left| \sum_{\nu \setminus \{b\}} (\phi'' \circ f) \{\Delta^+ g\}^2 - A_6 \right| < \epsilon, \quad (3.42)$$

$$\max_{1 \leq j \leq k} \text{Osc}(f; (y_{j-1}, y_j)) < \delta \quad \text{and} \quad \sum_{j=1}^k v_2(h; (y_{j-1}, y_j)) < \epsilon. \quad (3.43)$$

Let m_0 be the minimal integer such that $\mu \subset \lambda_{m_0}$ and each pair $\{y_{j-1}, y_j\}$ is separated by at least two other points from λ_{m_0} . For each $j = 1, \dots, k-1$ and $m \geq m_0$, there exists an integer $i(j) = i_m(j) \in \{1, \dots, n(m) - 1\}$ such that $x_{i(j)}^m = y_j$. For each $m \geq m_0$, let $I_1(m) := \{x_1^m, x_{n(m)}^m, x_{i(j)}^m, x_{i(j)+1}^m: j = 1, \dots, k-1\}$ and $I_2(m) := \{1, \dots, n(m)\} \setminus I_1(m)$. By (3.41) and (3.42), there exists an integer $m_3 \geq m_0$ such that for all $m \geq m_3$

$$\left| \sum_{i \in I_1(m)} \left\{ \Delta_i^m(\phi \circ f) - (\phi' \circ f)(x_{i-1}^m) \Delta_i^m f \right\} - A_3 - A_4 \right| < 4\epsilon \quad (3.44)$$

and

$$\left| \sum_{i \in I_2(m)} \phi''(f(x_{i-1}^m)) \{\Delta_i^m g\}^2 - A_5 - A_6 \right| < 4\epsilon. \quad (3.45)$$

By Taylor's theorem with Lagrange's form of the remainder: for real numbers u, v , there is a $\theta = \theta(u, v) \in (0, 1)$ such that

$$\phi(v) = \phi(u) + \phi'(u)(v - u) + \frac{1}{2} \phi''(u + \theta(v - u)) [v - u]^2. \quad (3.46)$$

Then by a telescoping sum, for $m \geq m_0$, we have

$$\begin{aligned}
(\phi \circ f)(b) - (\phi \circ f)(a) &= \sum_{i \in I_2(m)} \left\{ \phi'(f(x_{i-1}^m)) \Delta_i^m f + \frac{1}{2} \phi''(y_i^m) \{\Delta_i^m f\}^2 \right\} + \sum_{i \in I_1(m)} \Delta_i^m (\phi \circ f) \\
&= \sum_{i=1}^{n(m)} \phi'(f(x_{i-1}^m)) \{\Delta_i^m g + \Delta_i^m h\} + \frac{1}{2} \sum_{i=1}^{n(m)} \phi''(f(x_{i-1}^m)) \{\Delta_i^m g\}^2 \\
&\quad - \frac{1}{2} \sum_{i \in I_1(m)} \phi''(f(x_{i-1}^m)) \{\Delta_i^m g\}^2 + \sum_{i \in I_1(m)} \left\{ \Delta_i^m (\phi \circ f) - (\phi' \circ f)(x_{i-1}^m) \Delta_i^m f \right\} + R_m,
\end{aligned} \tag{3.47}$$

where $y_i^m := f(x_{i-1}^m) + \theta_i^m \Delta_i^m f$ with $\theta_i^m \in (0, 1)$ for $i = 1, \dots, n(m)$, and

$$R_m := \frac{1}{2} \sum_{i \in I_2(m)} \left\{ \phi''(y_i^m) [2\Delta_i^m g \Delta_i^m h + \{\Delta_i^m h\}^2] + [\phi''(y_i^m) - \phi''(f(x_{i-1}^m))] \{\Delta_i^m g\}^2 \right\}.$$

Let $A := (\phi \circ f)(b) - (\phi \circ f)(a) - A_1 - \frac{1}{2}A_2 - A_3 - A_4$. Then by (3.39), (3.40), (3.44) and (3.45), for all $m \geq m_1 \vee m_2 \vee m_3$,

$$\begin{aligned}
|S_{LC}(\phi' \circ f, g; \lambda_m) - A| &\leq \left| \sum_{i=1}^{n(m)} \phi'(f(x_{i-1}^m)) \Delta_i^m h - A_1 \right| \\
&+ \frac{1}{2} \left| \sum_{i=1}^{n(m)} \phi''(f(x_{i-1}^m)) \{\Delta_i^m g\}^2 - (A_2 + A_5 + A_6) \right| + \frac{1}{2} \left| \sum_{i \in I_1(m)} \phi''(f(x_{i-1}^m)) \{\Delta_i^m g\}^2 - (A_5 + A_6) \right| \\
&+ \left| \sum_{i \in I_1(m)} \left\{ \Delta_i^m (\phi \circ f) - (\phi' \circ f)(x_{i-1}^m) \Delta_i^m f \right\} - (A_3 + A_4) \right| + |R_m| \leq \frac{15}{2} \epsilon + |R_m|.
\end{aligned}$$

To bound $|R_m|$ notice that for each $i \in I_2(m)$ with $m \geq m_0$, $[x_{i-1}^m, x_i^m] \subset (y_{j-1}, y_j)$ for some $j = 1, \dots, k$. By (3.43) and Hölder's inequality, it then follows that

$$\begin{aligned}
\sup_{m \geq m_0} |R_m| &\leq \frac{\|\phi'' \circ f\|_\infty}{2} \left\{ 2 \left([g]_\lambda(b) \sum_{j=1}^k v_2(h; (y_{j-1}, y_j)) \right)^{1/2} + \sum_{j=1}^k v_2(h; (y_{j-1}, y_j)) \right\} + \epsilon [g]_\lambda(b) \\
&\leq \sqrt{\epsilon} \{ (\|\phi'' \circ f\|_\infty / 2) (2\sqrt{[g]_\lambda(b)} + 1) + [g]_\lambda(b) \}.
\end{aligned}$$

Since $\epsilon \in (0, 1)$ is arbitrary, the limit (3.36) exists and relation (3.34) holds for $z = a$ and $y = b$. The proof of statement (a) is complete.

The proof of statement (b) is similar. Instead of (3.47), in this case we use the relations

$$\begin{aligned}
(\phi \circ f)(b) - (\phi \circ f)(a) &= \sum_{i \in I_2(m)} \left\{ \phi'(f(x_i^m)) \Delta_i^m f - \frac{1}{2} \phi''(y_i^m) \{\Delta_i^m f\}^2 \right\} + \sum_{i \in I_1(m)} \Delta_i^m (\phi \circ f) \\
&= \sum_{i=1}^{n(m)} \phi'(f(x_i^m)) \{\Delta_i^m g + \Delta_i^m h\} - \frac{1}{2} \sum_{i=1}^{n(m)} \phi''(f(x_i^m)) \{\Delta_i^m g\}^2 \\
&\quad + \frac{1}{2} \sum_{i \in I_1(m)} \phi''(f(x_i^m)) \{\Delta_i^m g\}^2 + \sum_{i \in I_1(m)} \left\{ \Delta_i^m (\phi \circ f) - (\phi' \circ f)(x_i^m) \Delta_i^m f \right\} + R_m,
\end{aligned}$$

valid for $m \geq m_0$, where $y_i^m := f(x_i^m) - \theta_i^m \Delta_i^m f$ with $\theta_i^m \in (0, 1)$ for $i = 1, \dots, n(m)$, and

$$R_m := -\frac{1}{2} \sum_{i \in I_2(m)} \left\{ \phi''(y_i^m) [2\Delta_i^m g \Delta_i^m h + \{\Delta_i^m h\}^2] + [\phi''(y_i^m) - \phi''(f(x_i^m))] \{\Delta_i^m g\}^2 \right\}.$$

Here we used the rearranged Taylor's theorem (3.46). The other steps of the proof are the same as in the proof of statement (a) except that now we approximate the integrals by Right Cauchy sums. The proof of Theorem 24 is complete. $\square$

The preceding chain rules for (λ, p) -decomposable functions are used to characterize functions having the quadratic λ -variation in terms of the existence of the λ -integrals of a special form. A statement of this result for the Left Cauchy λ -integral is given in the introductory section. The case of the Right Cauchy λ -integral is symmetric and is treated next.

Proposition 25 *Let f be a regulated function on $[a, b]$, and let $\lambda \in \Lambda[a, b]$ be such that the left discontinuity points of f are accessible by λ . The following three statements about f are equivalent:*

- (a) *f has the quadratic λ -variation on $[a, b]$;*
- (b) *for a C^1 class function ψ , the composition $\psi \circ f$ is Right Cauchy λ -integrable on $[a, b]$ with respect to f , and for any $a \leq u < v \leq b$,*

$$\begin{aligned} (\Psi \circ f)(v) &= (\Psi \circ f)(u) + (RC) \int_u^v (\psi \circ f) d_\lambda f - \frac{1}{2} (RS) \int_u^v (\psi' \circ f) d[f]_\lambda^c \\ &\quad + \sum_{(u,v)} \left\{ \Delta^-(\Psi \circ f) - (\psi \circ f) \Delta^- f \right\} + \sum_{[u,v)} \left\{ \Delta^+(\Psi \circ f) - (\psi \circ f)_+ \Delta^+ f \right\}, \end{aligned}$$

where the two sums converge unconditionally and $\Psi(u) = \Psi(0) + \int_0^u \psi(x) dx$, $u \in \mathbb{R}$;

- (c) *f is Right Cauchy λ -integrable on $[a, b]$ with respect to itself.*

If any of the three statements holds, then for $a \leq x \leq b$,

$$(RC) \int_a^x f d_\lambda f = \frac{1}{2} \left\{ f^2(x) - f^2(a) + [f]_\lambda(x) \right\}. \quad (3.48)$$

Proof. Since f is $(\lambda, 1)$ -decomposable and $(f - f(a), 0) \in D_{\lambda,1}^+(f)$, implication (a) $\Rightarrow$ (b) follows from statement (b) of Theorem 24. Taking $\psi(u) = u$ in statement (b) we get statement (c). To prove (c) $\Rightarrow$ (a), let $H(x) := -f^2(x) + f^2(a) + 2(RC) \int_a^x f d_\lambda f$ for $x \in [a, b]$, and notice that for each $a \leq u < v \leq b$ and any partition κ of $[a, b]$, we have

$$s_2(f; \kappa \cap [u, v]) = -f^2(v) + f^2(u) + 2S_{RC}(f, f; \kappa \cap [u, v]).$$

This identity also yields relation (3.48). The proof of Proposition 25 is complete. $\square$

Let f be a (λ, p) -decomposable function on $[a, b]$ for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$ with a decomposition $f \equiv C + g + h$. By Theorem 24, if ϕ is a function of the class C^2 , and if $(g, h) \in D_{\lambda,p}^+(f)$, then the composition $\phi \circ f$ has the decomposition

$$\phi \circ f(x) = \phi \circ f(a) + \Phi(x) + U(x), \quad a \leq x \leq b, \quad (3.49)$$

where

$$\Phi(x) := (LC) \int_a^x (\phi' \circ f) d_\lambda g \quad (3.50)$$

and

$$\begin{aligned} U(x) := & (LY) \int_a^x (\phi' \circ f) dh + \frac{1}{2} (RS) \int_a^x (\phi'' \circ f) d[g]_\lambda^c \\ & + \sum_{(a,x]} \left\{ \Delta^-(\phi \circ f) - (\phi' \circ f)_- \Delta^- f \right\} + \sum_{[a,x)} \left\{ \Delta^+(\phi \circ f) - (\phi' \circ f)_+ \Delta^+ f \right\}. \end{aligned} \quad (3.51)$$

Moreover, for the same function ϕ , if $(g, h) \in D_{\lambda,p}^-(f)$ then the composition $\phi \circ f$ has the decomposition

$$\phi \circ f(x) = \phi \circ f(a) + \Psi(x) + V(x), \quad a \leq x \leq b, \quad (3.52)$$

where

$$\Psi(x) := (RC) \int_a^x (\phi' \circ f) d_\lambda g \quad (3.53)$$

and

$$\begin{aligned} V(x) := & (RY) \int_a^x (\phi' \circ f) dh - \frac{1}{2} (RS) \int_a^x (\phi'' \circ f) d[g]_\lambda^c \\ & + \sum_{(a,x]} \left\{ \Delta^-(\phi \circ f) - (\phi' \circ f) \Delta^- f \right\} + \sum_{[a,x)} \left\{ \Delta^+(\phi \circ f) - (\phi' \circ f)_+ \Delta^+ f \right\}. \end{aligned} \quad (3.54)$$

Using Theorem 2.24, one can show that the functions U and V have bounded p -variation. Also, the functions Φ and Ψ have the same p -variation property as the function g . The latter follows from the corresponding chain rule formulas. Next we show that Φ and Ψ have the quadratic λ -variation. Thus the composition $\phi \circ f$ is a function of the class $\mathcal{D}_{\lambda,p}[a, b]$ by Corollary 3 below. This result is reminiscent of the important fact of Stochastic Analysis that the class of semimartingales is closed under taking a composition with a C^2 class function, which is a consequence of Itô's formula and the fact that the stochastic integral with respect to a martingale is also a martingale.

Theorem 26 *For $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, let $f \equiv C + g + h$ with $(g, h) \in D_{\lambda,p}(f)$, and let ϕ be a C^2 class function. Then the following two statements hold:*

- (a) *If $(g, h) \in D_{\lambda,p}^+(f)$ then the indefinite Left Cauchy λ -integral Φ defined by (3.50) has the quadratic λ -variation with the bracket function*

$$[\Phi]_\lambda(x) = (LY) \int_a^x (\phi' \circ f)^2 d[g]_\lambda, \quad x \in [a, b], \quad (3.55)$$

and the 2-vector function (Φ, g) has the quadratic λ -covariation with the bracket function

$$[\Phi, g]_\lambda(x) = (LY) \int_a^x (\phi' \circ f) d[g]_\lambda, \quad x \in [a, b]. \quad (3.56)$$

- (b) *If $(g, h) \in D_{\lambda,p}^-(f)$ then the indefinite Right Cauchy λ -integral Ψ defined by (3.53) has the quadratic λ -variation with the bracket function*

$$[\Psi]_\lambda(x) = (RY) \int_a^x (\phi' \circ f)^2 d[g]_\lambda, \quad x \in [a, b],$$

and the 2-vector function (Ψ, g) has the quadratic λ -covariation with the bracket function

$$[\Psi, g]_\lambda(x) = (RY) \int_a^x (\phi' \circ f) d[g]_\lambda, \quad x \in [a, b].$$

Proof. We prove only statement (a) because the proof of statement (b) is symmetric. Let $(g, h) \in D_{\lambda, p}^+(f)$ be such that $f \equiv C + g + h$. By Definition 17, the indefinite Left Cauchy λ -integral Φ is a regulated function on $[a, b]$. Statement (a) will be proved once we show that for $a \leq u < v \leq b$,

$$\lim_{m \rightarrow \infty} s_2(\Phi; \lambda_m \cap [u, v]) = (LY) \int_u^v (\phi' \circ f)^2 d[g]_\lambda \quad (3.57)$$

and

$$\lim_{m \rightarrow \infty} C(\Phi, g; \lambda_m \cap [u, v]) = (LY) \int_u^v (\phi' \circ f) d[g]_\lambda. \quad (3.58)$$

Indeed, let $[\Phi]_\lambda(x) := (LY) \int_a^x (\phi' \circ g)^2 d[g]_\lambda$ for $a \leq x \leq b$. Then by statement (a) of Proposition 2.27, by (3.4) with $H = [g]_\lambda$, and by (3.32), we have

$$\Delta^-[\Phi]_\lambda = (\phi' \circ f)_-^2 \Delta^-[g]_\lambda = \{\Delta^- \Phi\}^2 \quad \text{and} \quad \Delta^+[\Phi]_\lambda = (\phi' \circ f)^2 \Delta^+[g]_\lambda = \{\Delta^+ \Phi\}^2.$$

Thus the indefinite Left Cauchy integral Φ has the quadratic λ -variation with the bracket function $[\Phi]_\lambda$ satisfying (3.55) provided (3.57) holds. Likewise, let $[\Phi, g]_\lambda(x) := (LY) \int_a^x (\phi' \circ g) d[g]_\lambda$ for $a \leq x \leq b$. By the same arguments, we have

$$\Delta^-[\Phi, g]_\lambda = (\phi' \circ f)_- \Delta^-[g]_\lambda = \Delta^- \Phi \Delta^- g \quad \text{and} \quad \Delta^+[\Phi, g]_\lambda = (\phi' \circ f) \Delta^+[g]_\lambda = \Delta^+ \Phi \Delta^+ g.$$

Thus the 2-vector function (Φ, g) has the quadratic λ -covariation with the bracket function $[\Phi, g]_\lambda$ satisfying (3.56) provided (3.58) holds.

Let $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$ for $m = 1, 2, \dots$. For notation simplicity, we prove (3.57) and (3.58) only for $u = a$ and $v = b$. Using notation (3.51) for U , by the chain rule formula (3.34), $\Phi(x) = (\phi \circ f)(x) - (\phi \circ f)(a) - U(x)$ for $a \leq x \leq b$. For $i = 1, \dots, n(m)$ and $m = 1, 2, \dots$, by the mean value theorem, there is a $\theta_i^m \in (0, 1)$ such that

$$\Delta_i^m \Phi = \Delta_i^m(\phi \circ f) - \Delta_i^m U = (\phi' \circ f)(x_{i-1}^m) \Delta_i^m g + B_i^m,$$

where $\Delta_i^m \rho := \rho(x_i^m) - \rho(x_{i-1}^m)$ for a function ρ on $[a, b]$,

$$B_i^m := (\phi' \circ f)(x_{i-1}^m) \Delta_i^m h + [\phi'(y_i^m) - \phi'(f(x_{i-1}^m))] \Delta_i^m f - \Delta_i^m U$$

and $y_i^m := f(x_{i-1}^m) + \theta_i^m \Delta_i^m f$. Let $\mu = \{z_j: j = 1, \dots, s-1\}$ be a non-empty set of points in the open interval (a, b) . Then either $I(\mu \cap \lambda) := \{j = 1, \dots, s-1: z_j \in \cup_m \lambda_m\} \neq \emptyset$ or $I(\mu \setminus \lambda) := \{j = 1, \dots, s-1\} \setminus I(\mu \cap \lambda) \neq \emptyset$. If $j \in I(\mu \cap \lambda)$ then $z_j = x_{i(j)}^m$ for some index $i(j) = i_m(j) \in \{1, \dots, n(m)-1\}$ and for all sufficiently large m . If $j \in I(\mu \setminus \lambda)$ then $x_{i(j)-1}^m < z_j < x_{i(j)}^m$ for some index $i(j) = i_m(j) \in \{1, \dots, n(m)\}$ and for all m . Let m_0 be the first integer such that in between each pair $\{z_{j-1}, z_j\}$ there are at least two different points from λ_{m_0} and $\{z_j: j \in I(\mu \cap \lambda)\} \subset \lambda_{m_0}$ provided $I(\mu \cap \lambda) \neq \emptyset$. For $m \geq m_0$, let

$$I_1(m) := \{x_1^m, x_{n(m)}^m\} \cup \{x_{i(j)}^m, x_{i(j)+1}^m: j \in I(\mu \cap \lambda)\} \cup \{x_{i(j)}^m: j \in I(\mu \setminus \lambda)\} \quad (3.59)$$

and let $I_2(m) := \{1, \dots, n(m)\} \setminus I_1(m)$. Then we have

$$\begin{aligned} s_2(\Phi; \lambda_m) &= \sum_{i \in I_1(m)} \{\Delta_i^m \Phi\}^2 + \sum_{i \in I_2(m)} \{(\phi' \circ f)(x_{i-1}^m) \Delta_i^m g + B_i^m\}^2 \\ &= \sum_{i \in I_1(m)} \left[\{\Delta_i^m \Phi\}^2 - \{(\phi' \circ f)(x_{i-1}^m) \Delta_i^m g\}^2 \right] + \sum_{i=1}^{n(m)} \{(\phi' \circ f)(x_{i-1}^m) \Delta_i^m g\}^2 + R_m^{(1)} \end{aligned}$$

for each $m \geq m_0$, where

$$R_m^{(1)} := 2 \sum_{i \in I_2(m)} \{(\phi' \circ f)(x_{i-1}^m) \Delta_i^m g\} B_i^m + \sum_{i \in I_2(m)} \{B_i^m\}^2.$$

Likewise, for $m \geq m_0$, we have

$$\begin{aligned} C(\Phi, g; \lambda_m) &= \sum_{i \in I_1(m)} \Delta_i^m \Phi \Delta_i^m g + \sum_{i \in I_2(m)} \{(\phi' \circ f)(x_{i-1}^m) \Delta_i^m g + B_i^m\} \Delta_i^m g \\ &= \sum_{i \in I_1(m)} \left[\Delta_i^m \Phi \Delta_i^m g - (\phi' \circ f)(x_{i-1}^m) \{\Delta_i^m g\}^2 \right] + \sum_{i=1}^{n(m)} (\phi' \circ f)(x_{i-1}^m) \{\Delta_i^m g\}^2 + R_m^{(2)}, \end{aligned}$$

where $R_m^{(2)} := \sum_{i \in I_2(m)} B_i^m \Delta_i^m g$. Suppose that given $\epsilon > 0$ one can find a set μ for (3.59) such that $|R_m^{(1)}| \vee |R_m^{(2)}| < C\epsilon$ for all sufficiently large m . Since $(g, h) \in D_{\lambda, p}^+(f)$, and so $\Delta^+ g(z_j) = 0$ for $j \in I(\mu \setminus \lambda)$, we have

$$\begin{aligned} &\lim_{m \rightarrow \infty} \sum_{i \in I_1(m)} \left[(\Delta_i^m \Phi)^2 - ((\phi' \circ f)(x_{i-1}^m) \Delta_i^m g)^2 \right] \\ &= \sum_{\mu \cup \{b\}} \left\{ (\Delta^- \Phi)^2 - ((\phi' \circ f)_- (\Delta^- g))^2 \right\} + \sum_{\{a\} \cup \mu} \left\{ (\Delta^+ \Phi)^2 - ((\phi' \circ f) \Delta^+ g)^2 \right\} = 0 \end{aligned}$$

by condition (3.32) of Definition 17 for any given μ . Likewise for any given μ , we have

$$\begin{aligned} &\lim_{m \rightarrow \infty} \sum_{i \in I_1(m)} \left[\Delta_i^m \Phi \Delta_i^m g - (\phi' \circ f)(x_{i-1}^m) (\Delta_i^m g)^2 \right] \\ &= \sum_{\mu \cup \{b\}} \left\{ \Delta^- \Phi \Delta^- g - (\phi' \circ f)_- (\Delta^- g)^2 \right\} + \sum_{\{a\} \cup \mu} \left\{ \Delta^+ \Phi \Delta^+ g - (\phi' \circ f) (\Delta^+ g)^2 \right\} = 0. \end{aligned}$$

Since $\{x \in (a, b): \Delta^+ g(x) \neq 0\} \subset \cup_m \lambda_m$, we can and do apply Theorem 11 which then yields

$$\lim_{m \rightarrow \infty} s_2(\Phi; \lambda_m) = \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} (\phi' \circ f)^2(x_{i-1}^m) (\Delta_i^m g)^2 = (LY) \int_a^b (\phi' \circ f)^2 d[g]_\lambda$$

and

$$\lim_{m \rightarrow \infty} C(\Phi, g; \lambda_m) = \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} (\phi' \circ f)(x_{i-1}^m) (\Delta_i^m g)^2 = (LY) \int_a^b (\phi' \circ f) d[g]_\lambda.$$

Thus (3.57) and (3.58) hold for $u = a$ and $u = b$ provided given $\epsilon \in (0, 1)$ one can find a set $\mu \subset (a, b)$ for (3.59) such that $|R_m^{(1)}| \vee |R_m^{(2)}| < C\epsilon$ for all sufficiently large m . We show

this for the remainder $R_m^{(1)}$ because the bound of the second remainder $R_m^{(2)}$ is similar. Let $\epsilon \in (0, 1)$. Since ϕ' is uniformly continuous on the range of f , there exists a $\delta > 0$ such that $|\phi'(u) - \phi'(v)| < \epsilon$ whenever $|u| \vee |v| \leq \|f\|_\infty$ and $|u - v| < \delta$. In the case $p \in (1, 2)$ there is a $q < p/(p-1)$ such that $f \in \mathcal{W}_q^*$. Let $z_0 := a$, $z_s := b$, $\Delta^- S := \Delta^-(\phi \circ f) - (\phi' \circ f)_- \Delta^- f$ and $\Delta^+ S := \Delta^+(\phi \circ f) - (\phi' \circ f) \Delta^+ f$. Then choose a set $\mu = \{z_j : j = 1, \dots, s-1\} \subset (a, b)$ such that

$$\max_{1 \leq j \leq s} \text{Osc}(f; (z_{j-1}, z_j)) < \delta, \quad \max_{1 \leq j \leq s} \text{Osc}(g; (z_{j-1}, z_j)) < \epsilon,$$

$$\max_{1 \leq j \leq s} \sup_{z_{j-1} \leq u < v \leq z_j} (RS) \int_u^v (\phi'' \circ f) d[g]^c < \epsilon, \quad \sum_{(a,b) \setminus \mu} \{|\Delta^- S| + |\Delta^+ S|\} < \epsilon,$$

and in addition for the case $p \in (1, 2)$,

$$\sum_{j=1}^s v_q(f; (z_{j-1}, z_j)) < \epsilon \quad \text{by Lemma 2.9, and} \quad \sum_{(a,b) \setminus \mu} |\Delta^- f|^q < \epsilon.$$

To bound $|R_m^{(1)}|$, recall that

$$\begin{aligned} B_i^m &= [\phi'(y_i^m) - \phi'(f(x_{i-1}^m))] \Delta_i^m f - \left[(LY) \int_{x_{i-1}^m}^{x_i^m} (\phi' \circ f) dh - (\phi' \circ f)(x_{i-1}^m) \Delta_i^m h \right] \\ &\quad - \frac{1}{2} (RS) \int_{x_{i-1}^m}^{x_i^m} (\phi'' \circ f) d[g]_\lambda^c - \sum_{(x_{i-1}^m, x_i^m]} \Delta^- S - \sum_{[x_{i-1}^m, x_i^m)} \Delta^+ S \end{aligned}$$

for $i = 1, \dots, n(m)$. By Definition 21 of the LY integral, the absolute value of the second term on the right side is equal to

$$D_i^m := \left| (RRS) \int_{x_{i-1}^m}^{x_i^m} \left\{ (\phi' \circ f)_-^{(x_{i-1}^m)} - (\phi' \circ f)(x_{i-1}^m) \right\} dh_+^{(x_i^m)} + \sum_{(x_{i-1}^m, x_i^m)} \Delta^- (\phi' \circ f) \Delta^+ h \right|.$$

In the case $p \in (1, 2)$, by inequality (2.23) and Hölder's inequality, we have the bound

$$\begin{aligned} D_i^m &\leq KV_q((\phi' \circ f)_-^{(x_{i-1}^m)}; [x_{i-1}^m, x_i^m]) V_p(h_+^{(x_i^m)}; [x_{i-1}^m, x_i^m]) \\ &\quad + \left(\sum_{(x_{i-1}^m, x_i^m)} |\Delta^- (\phi' \circ f)|^q \right)^{1/q} \left(\sum_{(x_{i-1}^m, x_i^m)} |\Delta^+ h|^p \right)^{1/p} \\ &\leq K_1 V_q(f; [x_{i-1}^m, x_i^m]) V_p(h; [x_{i-1}^m, x_i^m]) + K_2 \left(\sum_{(x_{i-1}^m, x_i^m)} |\Delta^- f|^q \right)^{1/q} \left(\sum_{(x_{i-1}^m, x_i^m)} |\Delta^+ h|^p \right)^{1/p} \end{aligned}$$

for some finite constants K_1 and K_2 . In the case $p = 1$, again by inequality (2.23), we have the bound

$$D_i^m \leq K_3 \text{Osc}(f; [x_{i-1}^m, x_i^m]) \left\{ V_1(h; [x_{i-1}^m, x_i^m]) + \sum_{(x_{i-1}^m, x_i^m)} |\Delta^+ h| \right\}$$

for some finite constant K_3 . Let m_0 be the integer defined in the first part of the proof. Then in the case $p \in (1, 2)$, for all $m \geq m_0$, we have

$$|R_m^{(1)}| \leq 2 \|\phi' \circ f\|_\infty \sum_{i \in I_2(m)} |\Delta_i^m g| |B_i^m| + \sum_{i \in I_2(m)} \{B_i^m\}^2$$

$$\begin{aligned}
&\leq \|\phi' \circ f\|_\infty \left\{ \max_{i \in I_2(m)} |\phi'(y_i^m) - \phi'(f(x_{i-1}^m))| (3[g]_\lambda(b) + v_2(h)) \right. \\
&\quad + 2K_1 V_p(h) V_q(g) \left(\sum_{j=1}^s v_q(f; (z_{j-1}, z_j)) \right)^{1/q} + 2K_2 \sigma_p(h)^{1/p} V_q(g) \left(\sum_{(a,b) \setminus \mu} |\Delta^- f|^q \right)^{1/q} \\
&\quad + \max_{i \in I_2(m)} |\Delta_i^m g| \left((RS) \int_a^b \phi'' \circ f d[g]_\lambda^c + 2 \sum_{(a,b)} \{|\Delta^- S| + |\Delta^+ S|\} \right) \Big\} \\
&\quad + 4 \max_{i \in I_2(m)} [\phi'(y_i^m) - \phi'(f(x_{i-1}^m))]^2 ([g]_\lambda(b) + v_2(h)) \\
&\quad + 2K_1^2 V_p(h)^2 \left(\sum_{j=1}^s v_q(f; (z_{j-1}, z_j)) \right)^{2/q} + 2K_2 \sigma_p(h)^{2/p} \left(\sum_{(a,b) \setminus \mu} |\Delta^- f|^q \right)^{2/q} \\
&\quad + \frac{1}{2} (RS) \int_a^b \phi'' \circ f d[g]_\lambda^c \max_{i \in I_2(m)} (RS) \int_{x_{i-1}^m}^{x_i^m} \phi'' \circ f d[g]_\lambda^c + 2 \left(\sum_{(a,b) \setminus \mu} \{|\Delta^- S| + |\Delta^+ S|\} \right)^2 \\
&\leq C\epsilon
\end{aligned}$$

for a constant C independent of m by the choice of the set μ . The same bound with a different constant C follows in the case $p = 1$. The proof of Theorem 26 is complete. $\square$

Finally we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. Suppose that $(g, h) \in D_{\lambda, p}^+(f)$ is such that $f \equiv C + g + h$. By Theorem 24, decomposition (3.49) holds. By the preceding Theorem 26, Φ has the quadratic λ -variation, and by Theorem 2.24, the function U has bounded p -variation. If $1 < p < 2$ then since ϕ is uniformly Lipschitz function on the range of f , it follows that Φ has bounded q -variation for some $q < p/(p-1)$, and so $g \in \text{dual}(\mathcal{W}_p)[a, b]$. Therefore (3.49) is a (λ, p) -decomposition of $\phi \circ f$. If $(g, h) \in D_{\lambda, p}^-(f)$ then it follows similarly that (3.52) is a (λ, p) -decomposition of $\phi \circ f$. The proof is complete. $\square$

3.3 Vector valued functions

Definition 27 Let $f_k, k = 1, \dots, d, d \geq 2$, be regulated functions on $[a, b]$, and let $\lambda \in \Lambda[a, b]$. We say that the d -vector function $\mathbf{f} = (f_1, \dots, f_d)$ has the *quadratic λ -variation on $[a, b]$* if each 2-vector function $(f_k, f_l), k, l = 1, \dots, d$ has the quadratic λ -covariation on $[a, b]$. The matrix valued function $[\mathbf{f}]_\lambda := ([f_k, f_l]_\lambda)_{k, l=1, \dots, d}$ is called the bracket function of $\mathbf{f}$.

In the case $d = 2$, a 2-vector function $\mathbf{f} = (f_1, f_2)$ has the quadratic λ -variation if f_1, f_2 each has the quadratic λ -variation and the 2-vector function (f_1, f_2) has the quadratic λ -covariation, and then the bracket function of $\mathbf{f}$ is given by 2×2 -matrix function

$$[\mathbf{f}]_\lambda = \begin{pmatrix} [f_1]_\lambda & [f_1, f_2]_\lambda \\ [f_2, f_1]_\lambda & [f_2]_\lambda \end{pmatrix}.$$

To illustrate the preceding definition we reformulate Theorem 26 as follows. Let $f \equiv C + g + h$ with $(g, h) \in D_{\lambda, p}^+(f)$ for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, let ϕ be a function of the class C^2 and

let Φ be the indefinite Left Cauchy λ -integral given by (3.50). Then the 2-vector function (Φ, g) has the quadratic λ -variation with the bracket function

$$[(\Phi, g)]_\lambda(x) = \begin{pmatrix} [\Phi]_\lambda & [\Phi, g]_\lambda \\ [g, \Phi]_\lambda & [g]_\lambda \end{pmatrix}(x) = (LY) \int_a^x \begin{pmatrix} (\phi' \circ f)^2 & \phi' \circ f \\ \phi' \circ f & 1 \end{pmatrix} d[g]_\lambda$$

for $a \leq x \leq b$. Likewise for the indefinite Right Cauchy integral Ψ given by (3.53), if $(g, h) \in D_{\lambda, p}^-(f)$ then the 2-vector function (Ψ, g) has the quadratic λ -variation with the bracket function

$$[(\Psi, g)]_\lambda(x) = \begin{pmatrix} [\Psi]_\lambda & [\Psi, g]_\lambda \\ [g, \Psi]_\lambda & [g]_\lambda \end{pmatrix}(x) = (RY) \int_a^x \begin{pmatrix} (\phi' \circ f)^2 & \phi' \circ f \\ \phi' \circ f & 1 \end{pmatrix} d[g]_\lambda$$

for $a \leq x \leq b$.

Proposition 28 *Let $f_k \in \mathcal{R}[a, b]$, $k = 1, \dots, d$, and let $\lambda \in \Lambda[a, b]$. The following two statements are equivalent:*

- (a) *the d -vector function $(f_1, \dots, f_d)$ has the quadratic λ -variation on $[a, b]$;*
- (b) *for each pair (k, l) of indices $k, l = 1, \dots, d$, the functions $f_k + f_l$ and $f_k - f_l$ have the quadratic λ -variation on $[a, b]$.*

If either of the two statements hold, then for each pair (k, l) , $k, l = 1, \dots, d$,

$$[f_k, f_l]_\lambda = \frac{1}{4} \{ [f_k + f_l]_\lambda - [f_k - f_l]_\lambda \}. \quad (3.60)$$

Proof. (a) $\Rightarrow$ (b): Let (k, l) be a pair of indices $k, l \in \{1, \dots, d\}$ such that $k \neq l$. By Definition 27, there exist the bracket functions $[f_k]_\lambda$, $[f_l]_\lambda$ and $[f_k, f_l]_\lambda$. Let $[f_k + f_l]_\lambda := [f_k]_\lambda + 2[f_k, f_l]_\lambda + [f_l]_\lambda$ and $[f_k - f_l]_\lambda := [f_k]_\lambda - 2[f_k, f_l]_\lambda + [f_l]_\lambda$. It is easy to check that $[f_k + f_l]_\lambda$ and $[f_k - f_l]_\lambda$ are the bracket functions of $f_k + f_l$ and $f_k - f_l$, respectively.

(b) $\Rightarrow$ (a): Given a pair (k, l) of indices, let $[f_k, f_l]_\lambda := (1/4)\{[f_k + f_l]_\lambda - [f_k - f_l]_\lambda\}$. It is then easy to check that the 2-vector function (f_k, f_l) has the quadratic λ -covariation with the bracket function $[f_k, f_l]_\lambda$. The proof of Proposition 28 is complete. $\square$

Suppose that a 2-vector function (f_1, f_2) on $[a, b]$ has the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$. Let

$$[f_1, f_2]_\lambda^c := \frac{1}{4} \{ [f_1 + f_2]_\lambda^c - [f_1 - f_2]_\lambda^c \}.$$

Then the following is true.

Proposition 29 *Let (f_1, f_2) be a 2-vector function on $[a, b]$ having the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$. Then for each $a \leq x \leq b$,*

$$[f_1, f_2]_\lambda(x) = [f_1, f_2]_\lambda^c(x) + \sum_{(a, x]} \Delta^- f_1 \Delta^- f_2 + \sum_{[a, x)} \Delta^+ f_1 \Delta^+ f_2. \quad (3.61)$$

Proof. Since $[f_1 + f_2]_\lambda^c$ and $[f_1 - f_2]_\lambda^c$ are continuous parts of the bracket functions $[f_1 + f_2]_\lambda$ and $[f_1 - f_2]_\lambda$, respectively, it follows that for each $a \leq x \leq b$,

$$\frac{1}{4} \{ [f_1 + f_2]_\lambda^c(x) - [f_1 - f_2]_\lambda^c(x) \} = \frac{1}{4} \{ [f_1 + f_2]_\lambda(x) - [f_1 - f_2]_\lambda(x) \}$$

$$\begin{aligned}
& -\frac{1}{4}\left\{\sum_{(a,x)}\{\Delta^-(f_1+f_2)\}^2-\sum_{(a,x)}\{\Delta^-(f_1-f_2)\}^2\right\}-\frac{1}{4}\left\{\sum_{[a,x)}\{\Delta^+(f_1+f_2)\}^2-\sum_{[a,x)}\{\Delta^+(f_1-f_2)\}^2\right\} \\
& = [f_1, f_2]_\lambda(x) - \sum_{(a,x)} \Delta^- f_1 \Delta^- f_2 - \sum_{[a,x)} \Delta^+ f_1 \Delta^+ f_2,
\end{aligned}$$

where the last equality follows by linearity of the unconditional sums and by the preceding Proposition 28. Therefore (3.61) holds by the definition of $[f_1, f_2]_\lambda^c$. $\square$

Proposition 30 *For $\lambda \in \Lambda[a, b]$, $1 \leq p < 2$ and $k = 1, 2$, let f_k be a function on $[a, b]$ which is (λ, p) -decomposable by $(g_k, h_k) \in D_{\lambda, p}^+(f_k) \cup D_{\lambda, p}^-(f_k)$. The 2-vector function (f_1, f_2) has the quadratic λ -covariation if and only if the 2-vector function (g_1, g_2) has the quadratic λ -covariation. If both 2-vector functions have the quadratic λ -covariation then for $a \leq x \leq b$,*

$$[f_1, f_2]_\lambda(x) = [g_1, g_2]_\lambda^c(x) + \sum_{(a,x)} \Delta^- f_1 \Delta^- f_2 + \sum_{[a,x)} \Delta^+ f_1 \Delta^+ f_2. \quad (3.62)$$

Proof. For $k = 1, 2$, let $(g_k, h_k) \in D_{\lambda, p}^+(f_k)$. By Proposition 14, the three 2-vector functions (g_1, h_2) , (g_2, h_1) and (h_1, h_2) have the quadratic λ -covariations, and for $a \leq x \leq b$,

$$\begin{aligned}
[g_1, h_2]_\lambda(x) &= \sum_{(a,x)} \Delta^- g_1 \Delta^- h_2 + \sum_{[a,x)} \Delta^+ g_1 \Delta^+ h_2, \\
[g_2, h_1]_\lambda(x) &= \sum_{(a,x)} \Delta^- g_2 \Delta^- h_1 + \sum_{[a,x)} \Delta^+ g_2 \Delta^+ h_1, \\
[h_1, h_2]_\lambda(x) &= \sum_{(a,x)} \Delta^- h_1 \Delta^- h_2 + \sum_{[a,x)} \Delta^+ h_1 \Delta^+ h_2.
\end{aligned}$$

If $(g_k, h_k) \in D_{\lambda, p}^-(f_k)$ for $k = 1, 2$, then the same conclusion holds. Let $\lambda = \{\lambda_m; m \geq 1\}$. Then for each $m \geq 1$, we have

$$C(f_1, f_2; \lambda_m) - C(g_1, g_2; \lambda_m) = C(g_1, h_2; \lambda_m) + C(g_2, h_1; \lambda_m) + C(h_1, h_2; \lambda_m).$$

Since the right side has a limit as $m \rightarrow \infty$, the first part of the conclusion holds. By linearity of the unconditional sums, it follows that for $a \leq x \leq b$,

$$\begin{aligned}
[f_1, f_2]_\lambda(x) &= [g_1, g_2]_\lambda(x) + [g_1, h_2]_\lambda(x) + [g_2, h_1]_\lambda(x) + [h_1, h_2]_\lambda(x) \\
&= [g_1, g_2]_\lambda^c(x) + \sum_{(a,x)} \left\{ \Delta^- g_1 \Delta^- g_2 + \Delta^- g_1 \Delta^- h_2 + \Delta^- g_2 \Delta^- h_1 + \Delta^- h_1 \Delta^- h_2 \right\} \\
&\quad + \sum_{[a,x)} \left\{ \Delta^+ g_1 \Delta^+ g_2 + \Delta^+ g_1 \Delta^+ h_2 + \Delta^+ g_2 \Delta^+ h_1 + \Delta^+ h_1 \Delta^+ h_2 \right\},
\end{aligned}$$

proving (3.62). The proof is complete. $\square$

The next statement extends Theorem 11 to bracket functions of the quadratic λ -covariation, and hence to bracket functions of the quadratic λ -variation of vector functions.

Lemma 31 *Let $(f_1, \dots, f_d)$ have the quadratic λ -variation on $[a, b]$ for some sequence $\lambda = \{\lambda_m; m \geq 1\} \in \Lambda[a, b]$ of partitions $\lambda_m = \{x_i^m; i = 0, \dots, n(m)\}$, and let $h \in \mathcal{R}[a, b]$. Then the following two statements hold:*

(a) If $\cup_k \{x \in (a, b): \Delta^+ f_k(x) \neq 0\} \subset \cup_m \lambda_m$, then

$$\lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} h(x_{i-1}^m) [f_k(x_i^m) - f_k(x_{i-1}^m)] [f_l(x_i^m) - f_l(x_{i-1}^m)] = (LY) \int_a^b h d[f_k, f_l]_\lambda. \quad (3.63)$$

(b) If $\cup_k \{x \in (a, b): \Delta^- f_k(x) \neq 0\} \subset \cup_m \lambda_m$, then

$$\lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} h(x_i^m) [f_k(x_i^m) - f_k(x_{i-1}^m)] [f_l(x_i^m) - f_l(x_{i-1}^m)] = (RY) \int_a^b h d[f_k, f_l]_\lambda. \quad (3.64)$$

Proof. Since $[f_k, f_l]_\lambda$, $k, l = 1, \dots, d$, are functions of bounded variation on $[a, b]$, the Left Young and Right Young integrals in (3.63) and (3.64), respectively, exist by Lemma 9. For (a) suppose that $\cup_k \{x \in (a, b): \Delta^+ f_k(x) \neq 0\} \subset \cup_m \lambda_m$. If $k = l$ then (3.63) holds by Theorem 11. If $k \neq l$ then $\{x \in (a, b): \Delta^+(f_k + f_l)(x) \neq 0\} \subset \cup_k \{x \in (a, b): \Delta^+ f_k(x) \neq 0\}$ and $\{x \in (a, b): \Delta^+(f_k - f_l)(x) \neq 0\} \subset \cup_k \{x \in (a, b): \Delta^+ f_k(x) \neq 0\}$. Letting $\Delta_i^k \rho := \rho(x_i^k) - \rho(x_{i-1}^k)$ for a function ρ on $[a, b]$, by (3.60), by Theorem 11 and by the bilinearity of the Left Young integral, we have

$$\begin{aligned} \sum_{i=1}^{n(m)} h(x_{i-1}^m) \Delta_i^m f_k \Delta_i^m f_l &= \frac{1}{4} \sum_{i=1}^{n(m)} h(x_{i-1}^m) \left\{ (\Delta_i^m (f_k + f_l))^2 - (\Delta_i^m (f_k - f_l))^2 \right\} \\ &\xrightarrow{m \rightarrow \infty} \frac{1}{4} \left\{ (LY) \int_a^b h d[f_k + f_l]_\lambda - (LY) \int_a^b h d[f_k - f_l]_\lambda \right\} = (LY) \int_a^b h d[f_k, f_l]_\lambda, \end{aligned}$$

proving (3.63). The proof of statement (b) is analogous and is omitted. $\square$

Next we extend Definitions 17 and 18 of the two λ -integrals for vector-valued functions. Recall the notation of the sums S_{LC} and S_{RC} given before Definition 17.

Definition 32 Let $\mathbf{F} = (F_1, \dots, F_d)$ and $\mathbf{f} = (f_1, \dots, f_d)$ be two d -vector functions such that for each $k = 1, \dots, d$ $f_k \in \mathcal{R}[a, b]$, $F_k(t-)$ exists at $t \in (a, b]$ whenever $\Delta^- f_k(t) \neq 0$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. We say that the *Left Cauchy λ -integral (LC) $\int \langle \mathbf{F}, d_\lambda \mathbf{f} \rangle$ is defined on $[a, b]$* if there exists a regulated function Φ on $[a, b]$ such that $\Phi(a) = 0$ and for each $a \leq u < v \leq b$,

$$\Phi(v) - \Phi(u) = \lim_{m \rightarrow \infty} \sum_{k=1}^d S_{LC}(F_k, f_k; \lambda_m \cap [u, v]), \quad (3.65)$$

$$\Delta^- \Phi(v) = \sum_{k=1}^d F_k(v-) \Delta^- f_k(v) \quad \text{and} \quad \Delta^+ \Phi(u) = \sum_{k=1}^d F_k(u) \Delta^+ f_k(u), \quad (3.66)$$

where $F_k(v-) := 0$ whenever $\Delta^- f_k(v) = 0$. For $a \leq u < v \leq b$, let $(LC) \int_u^v \langle \mathbf{F}, d_\lambda \mathbf{f} \rangle := \Phi(v) - \Phi(u)$. If a regulated function Φ on $[a, b]$ exists and satisfies the same conditions except that (3.65) and (3.66) do not hold for $v = b$, then we say that the *Left Cauchy λ -integral (LC) $\int \langle \mathbf{F}, d_\lambda \mathbf{f} \rangle$ is defined on $[a, b)$* .

Definition 33 Let $\mathbf{F} = (F_1, \dots, F_d)$ and $\mathbf{f} = (f_1, \dots, f_d)$ be two d -vector functions such that for each $k = 1, \dots, d$ $f_k \in \mathcal{R}[a, b]$, $F_k(t+)$ exists at $t \in [a, b]$ whenever $\Delta^+ f_k(t) \neq 0$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. We say that the *Right Cauchy λ -integral* $(RC) \int \langle \mathbf{F}, d_\lambda \mathbf{f} \rangle$ is defined on $[a, b]$ if there exists a regulated function Ψ on $[a, b]$ such that $\Psi(a) = 0$ and for each $a \leq u < v \leq b$,

$$\Psi(y) - \Psi(z) = \lim_{m \rightarrow \infty} \sum_{k=1}^d S_{RC}(F_k, f_k; \lambda \cap [u, v]),$$

$$\Delta^- \Psi(v) = \sum_{k=1}^d F_k(v) \Delta^- f_k(v) \quad \text{and} \quad \Delta^+ \Psi(u) = \sum_{k=1}^d F_k(u+) \Delta^+ f_k(u),$$

where $F_k(u+) := 0$ whenever $\Delta^+ f_k(u) = 0$. For $a \leq u < v \leq b$, let $(RC) \int_u^v \langle \mathbf{F}, d_\lambda \mathbf{f} \rangle := \Psi(v) - \Psi(u)$.

Next chain rule formulas (3.34) and (3.35) of Theorem 24 are extended in two directions. First, the increment of the composition on the left side of each formula is replaced by an integral with respect to the composition. Second, the real valued functions f, g, h are replaced by d -vector functions $\mathbf{f} = (f_1, \dots, f_d)$, $\mathbf{g} = (g_1, \dots, g_d)$ and $\mathbf{h} = (h_1, \dots, h_d)$ so that for each $k = 1, \dots, d$, f_k, g_k and h_k satisfy the conditions of the case $d = 1$. For $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, we say that $\mathbf{f}$ is (λ, p) -decomposable by $(\mathbf{g}, \mathbf{h})$ if $\mathbf{f} = \mathbf{C} + \mathbf{g} + \mathbf{h}$ for some constant $\mathbf{C}$ and for each $k = 1, \dots, d$, (g_k, h_k) is a (λ, p) -dual pair. If in addition (cf. (1.17)),

$$\cup_{k=1}^d \left(N_{(a,b)}(\Delta^+ g_k) \cup N_{(a,b)}(\Delta^+ h_k) \right) \subset \cup \lambda \quad (3.67)$$

then we write $(\mathbf{g}, \mathbf{h}) \in D_{\lambda,p}^+(\mathbf{f})$. Likewise, if in addition, (3.67) holds with Δ^+ replaced by Δ^- then we write $(\mathbf{g}, \mathbf{h}) \in D_{\lambda,p}^-(\mathbf{f})$.

Theorem 34 Let $\mathbf{f} = (f_1, \dots, f_d)$, $\mathbf{g} = (g_1, \dots, g_d)$, $\mathbf{h} = (h_1, \dots, h_d)$, $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$ be such that $\mathbf{f}$ is (λ, p) -decomposable by $(\mathbf{g}, \mathbf{h})$. Let $\phi: \mathbb{R}^d \mapsto \mathbb{R}$ be a function of the class C^2 , and let $\psi \in \text{dual}(\mathcal{W}_p)[a, b]$. Then the following two statements hold:

- (a) If $(\mathbf{g}, \mathbf{h}) \in D_{\lambda,p}^+(\mathbf{f})$ then the Left Cauchy λ -integral $(LC) \int \langle (\psi, -\psi(\nabla \phi \circ \mathbf{f})), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle$ is defined on $[a, b]$, and for each $a \leq u < v \leq b$,

$$\begin{aligned} (LC) \int_u^v \langle (\psi, -\psi(\nabla \phi \circ \mathbf{f})), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle & \quad (3.68) \\ &= \sum_{k=1}^d (LY) \int_u^v \psi(\phi'_k \circ \mathbf{f}) dh_k + \frac{1}{2} \sum_{l,k=1}^d (RS) \int_u^v \psi(\phi''_{lk} \circ \mathbf{f}) d[g_l, g_k]_\lambda^c \\ &+ \sum_{(u,v)} \psi_- \left\{ \Delta^-(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f})_- \Delta^- f_k \right\} + \sum_{(u,v)} \psi \left\{ \Delta^+(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f}) \Delta^+ f_k \right\}, \end{aligned}$$

where the two sums converge unconditionally.

- (b) If $(\mathbf{g}, \mathbf{h}) \in D_{\lambda,p}^-(\mathbf{f})$ then the Right Cauchy λ -integral $(RC) \int \langle (\psi, -\psi(\nabla \phi \circ \mathbf{f})), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle$ is defined on $[a, b]$, and for each $a \leq u < v \leq b$,

$$(RC) \int_u^v \langle (\psi, -\psi(\nabla \phi \circ \mathbf{f})), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle \quad (3.69)$$

$$\begin{aligned}
&= \sum_{k=1}^d (RY) \int_u^v \psi(\phi'_k \circ \mathbf{f}) dh_k - \frac{1}{2} \sum_{l,k=1}^d (RS) \int_u^v \psi(\phi''_{lk} \circ \mathbf{f}) d[g_l, g_k]_\lambda^c \\
&\quad + \sum_{(u,v]} \psi \left\{ \Delta^-(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f}) \Delta^- f_k \right\} + \sum_{[u,v)} \psi_+ \left\{ \Delta^+(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f})_+ \Delta^+ f_k \right\},
\end{aligned}$$

where the two sums converge unconditionally.

Proof. We prove statement (a) and indicate necessary changes needed to prove statement (b). Suppose that (3.67) holds. In the case $1 < p < 2$ there exists $2 < q < p/(p-1)$ such that g_k has bounded q -variation for $k = 1, \dots, d$. We can and do assume that in this case ψ has bounded q -variation as well, and ψ is a regulated function in the case $p = 1$. Let $\phi: \mathbb{R}^d \mapsto \mathbb{R}$ be a function of the class C^2 . Statement (a) will be proved once we show that the limit

$$\begin{aligned}
&(LC) \int_u^v \langle (\psi, -\psi(\nabla \phi \circ \mathbf{f})), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle \\
&:= \lim_{m \rightarrow \infty} \left[S_{LC}(\psi, \phi \circ \mathbf{f}; \lambda \cap [u, v]) - \sum_{k=1}^d S_{LC}(\psi(\phi'_k \circ \mathbf{f}), g_k; \lambda \cap [u, v]) \right]
\end{aligned} \tag{3.70}$$

exists for each $a \leq u < v \leq b$ and satisfies relation (3.68). Indeed, let

$$\Phi(x) := (LC) \int_a^x \langle (\psi, -\psi(\nabla \phi \circ \mathbf{f})), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle, \quad a \leq x \leq b.$$

Clearly, condition (3.65) holds by definition of Φ , while condition (3.66) follows from relation (3.68) as in the proof of Theorem 24.

For notation simplicity, we prove that (3.70) exists and (3.68) holds for $u = a$ and $v = b$. If $1 < p < 2$ then $f_k \in \mathcal{W}_q$ for $2 < q < p/(p-1)$ and $k = 1, \dots, d$. Since ϕ' is Lipschitz function on the range of $\mathbf{f}$, and since $\mathcal{W}_q$ is a Banach algebra, $\psi(\phi'_k \circ \mathbf{f}) \in \mathcal{W}_q$ for $k = 1, \dots, d$. By Theorem 2.24 if $1 < p < 2$ and by Lemma 9 if $p = 1$, it then follows that there exist the Left Young integrals $A_k := (LY) \int_a^b \psi(\phi'_k \circ \mathbf{f}) dh_k$ for $k = 1, \dots, d$. The Riemann-Stieltjes integrals $B_{l,k} := (RS) \int_a^b \psi(\phi''_{lk} \circ \mathbf{f}) d[g_l, g_k]_\lambda^c$ for $l, k = 1, \dots, d$ exist by Lemma 9. The mean value theorem assure that the two sums in (3.68) converge absolutely because $\max_k \sigma_2(f_k) < \infty$, and thus converge unconditionally to values, say C^- and C^+ , respectively. Let $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$ for $m = 1, 2, \dots$. For each $m \geq 1$ and $i = 1, \dots, n(m)$, let $\Delta_i^m \rho := \rho(x_i^m) - \rho(x_{i-1}^m)$ for a function ρ on $[a, b]$. Let $\epsilon > 0$. Since $\cup_k \{x \in (a, b): \Delta^+ h_k(x) \neq 0\} \subset \cup_m \lambda_m$, by Theorem 2.22, there exists an integer m_1 such that for all $m \geq m_1$,

$$|A_k - S_{LC}(\psi(\phi'_k \circ \mathbf{f}), h_k; \lambda_m)| < \epsilon \tag{3.71}$$

for $k = 1, \dots, d$. Since $\cup_k \{x \in (a, b): \Delta^+ g_k(x) \neq 0\} \subset \cup_m \lambda_m$, by Lemmas 31 and 9, there exists an integer m_2 such that for all $m \geq m_2$,

$$\left| B_{l,k} + B_{l,k}^- + B_{l,k}^+ - \sum_{i=1}^{n(m)} (\psi(\phi''_{lk} \circ \mathbf{f}))(x_{i-1}^m) \Delta_i^m g_l \Delta_i^m g_k \right| < \epsilon \tag{3.72}$$

for $l, k = 1, \dots, d$, where

$$B_{l,k}^- := \sum_{(a,b]} \psi_-(\phi''_{lk} \circ \mathbf{f})_- \Delta^- g_l \Delta^- g_k \quad \text{and} \quad B_{l,k}^+ := \sum_{[a,b)} \psi(\phi''_{lk} \circ \mathbf{f}) \Delta^+ g_l \Delta^+ g_k.$$

Since ϕ''_{lk} is uniformly continuous on the range of $\mathbf{f}$, there is a $\delta > 0$ such that $|\phi''_{lk}(u) - \phi''_{lk}(v)| < \epsilon$ for $l, k = 1, \dots, d$ whenever $\max_r |u_r - v_r| < \delta$ and $|u_r| \vee |v_r| \leq \|f_r\|_\infty$ for $r = 1, \dots, d$ and $u = (u_1, \dots, u_d)$, $v = (v_1, \dots, v_d)$. Therefore recalling Lemma 2.9, we conclude that there exists a finite set $\mu = \{z_j: j = 1, \dots, s-1\} \subset (a, b)$ such that for each $l, k = 1, \dots, d$,

$$\left| C^- - \sum_{\mu \cup \{b\}} \psi_- \left\{ \Delta^-(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f})_- \Delta^- f_k \right\} \right| \vee \left| B_{lk}^- - \sum_{\mu \cup \{b\}} \psi_- (\phi''_{lk} \circ \mathbf{f})_- \Delta^- g_l \Delta^- g_k \right| < \epsilon, \quad (3.73)$$

$$\left| C^+ - \sum_{\{a\} \cup \mu} \psi \left\{ \Delta^+(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f}) \Delta^+ f_k \right\} \right| \vee \left| B_{lk}^+ - \sum_{\{a\} \cup \mu} \psi (\phi''_{lk} \circ \mathbf{f}) \Delta^+ g_l \Delta^+ g_k \right| < \epsilon, \quad (3.74)$$

$$\max_{1 \leq k \leq d} \max_{1 \leq j \leq s} \text{Osc}(f_k; (z_{j-1}, z_j)) < \delta \quad \text{and} \quad \max_{1 \leq k \leq d} \sum_{j=1}^s v_2(h_k; (z_{j-1}, z_j)) < \epsilon, \quad (3.75)$$

where $z_0 := a$ and $z_s := b$. Let $I(\mu \cap \lambda) := \{j = 1, \dots, s-1: z_j \in \cup_m \lambda_m\}$. If $j \in I(\mu \cap \lambda)$ then $z_j = x_{i(j)}^m$ for some index $i(j) \in \{1, \dots, n(m)-1\}$ and for all sufficiently large m . If $j \in I(\mu \setminus \lambda) := \{j = 1, \dots, s-1\} \setminus I(\mu \cap \lambda)$ then $x_{i(j)-1}^m < z_j < x_{i(j)}^m$ for some index $i(j) \in \{1, \dots, n(m)\}$ and for all m . Let m_0 be the first integer such that $\{z_j: j \in I(\mu \cap \lambda)\} \subset \lambda_{m_0}$ and in between each pair (z_{j-1}, z_j) there are at least two different points from λ_{m_0} . For $m \geq m_0$, let $I_1(m) := \{x_1^m, x_{n(m)}^m\} \cup \{x_{i(j)}^m, x_{i(j)+1}^m: j \in I(\mu \cap \lambda)\} \cup \{x_{i(j)}^m: j \in I(\mu \setminus \lambda)\}$ and let $I_2(m) := \{1, \dots, n(m)\} \setminus I_1(m)$. Since $\Delta^+ g_k(z_j) = \Delta^+ h_k(z_j) = 0$ for $z_j \notin \cup_m \lambda_m$ and $k = 1, \dots, d$, we have that

$$\begin{aligned} & \lim_{m \rightarrow \infty} \sum_{i \in I_1(m)} \psi(x_{i-1}^m) \left\{ \Delta_i^m \phi \circ \mathbf{f} - \sum_{k=1}^d (\phi'_k \circ \mathbf{f})(x_{i-1}^m) \Delta_i^m f_k \right\} \\ &= \sum_{\mu \cup \{b\}} \psi_- \left\{ \Delta^-(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f})_- \Delta^- f_k \right\} + \sum_{\{a\} \cup \mu} \psi \left\{ \Delta^+(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f}) \Delta^+ f_k \right\} \end{aligned}$$

and

$$\begin{aligned} & \lim_{m \rightarrow \infty} \sum_{i \in I_1(m)} (\psi(\phi''_{lk} \circ \mathbf{f}))(x_{i-1}^m) \Delta_i^m g_l \Delta_i^m g_k \\ &= \sum_{\mu \cup \{b\}} \psi_- (\phi''_{lk} \circ \mathbf{f})_- \Delta^- g_l \Delta^- g_k + \sum_{\{a\} \cup \mu} \psi (\phi''_{lk} \circ \mathbf{f}) \Delta^+ g_l \Delta^+ g_k. \end{aligned}$$

Therefore by (3.73) and (3.74), there exists an integer $m_3 \geq m_0$ such that for all $m \geq m_3$

$$\left| \sum_{i \in I_1(m)} \psi(x_{i-1}^m) \left\{ \Delta_i^m \phi \circ \mathbf{f} - \sum_{k=1}^d (\phi'_k \circ \mathbf{f})(x_{i-1}^m) \Delta_i^m f_k \right\} - C^- - C^+ \right| < 4\epsilon \quad (3.76)$$

and

$$\left| \sum_{i \in I_1(m)} (\psi(\phi''_{lk} \circ \mathbf{f}))(x_{i-1}^m) \Delta_i^m g_l \Delta_i^m g_k - B_{lk}^- - B_{lk}^+ \right| < 4\epsilon \quad (3.77)$$

for each $l, k = 1, \dots, d$. We use Taylor's theorem with Lagrange's form of the remainder: for vectors $u = (u_1, \dots, u_d)$, $v = (v_1, \dots, v_d)$ there is a $\theta = \theta(u, v) \in (0, 1)$ such that

$$\phi(v) = \phi(u) + \sum_{k=1}^d \phi'_k(u)[v_k - u_k] + \frac{1}{2} \sum_{l,k=1}^d \phi''_{lk}(u + \theta(v - u))[v_l - u_l][v_k - u_k],$$

where $\phi'_l(y) = \frac{\partial \phi}{\partial y_l}(y)$ and $\phi''_{lk}(y) = \frac{\partial^2 \phi}{\partial y_l \partial y_k}(y)$ for $y = (y_1, \dots, y_d)$. Then by a telescoping sum, for $m \geq m_0$, it follows that

$$\begin{aligned}
S_{LC}(\psi, \phi \circ \mathbf{f}; \lambda_m) &= \sum_{i \in I_1(m)} \psi(x_{i-1}^m) \Delta_i^m(\phi \circ \mathbf{f}) \\
&+ \sum_{i \in I_2(m)} \psi(x_{i-1}^m) \left\{ \sum_{k=1}^d (\phi'_k \circ \mathbf{f})(x_{i-1}^m) \Delta_i^m f_k + \frac{1}{2} \sum_{l,k=1}^d \phi''_{lk}(y_i^m) \Delta_i^m f_l \Delta_i^m f_k \right\} \\
&= \sum_{k=1}^d S_{LC}(\psi(\phi'_k \circ \mathbf{f}), g_k; \lambda_m) + \sum_{k=1}^d S_{LC}(\psi(\phi''_k \circ \mathbf{f}), h_k; \lambda_m) \\
&+ \frac{1}{2} \sum_{l,k=1}^d \left(\sum_{i=1}^{n(m)} - \sum_{i \in I_1(m)} \right) (\psi(\phi''_{lk} \circ \mathbf{f}))(x_{i-1}^m) \Delta_i^m g_l \Delta_i^m g_k \\
&+ \sum_{i \in I_1(m)} \psi(x_{i-1}^m) \left\{ \Delta_i^m(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f})(x_{i-1}^m) \Delta_i^m f_k \right\} + R_m,
\end{aligned} \tag{3.78}$$

where $y_i^m := \mathbf{f}(x_{i-1}^m) + \theta_i^m \Delta_i^m \mathbf{f}$, $\theta_i^m \in (0, 1)$ for $i = 1, \dots, n(m)$, and

$$\begin{aligned}
R_m &:= \frac{1}{2} \sum_{l,k=1}^d \sum_{i \in I_2(m)} \left\{ \psi(x_{i-1}^m) \phi''_{lk}(y_i^m) [\Delta_i^m g_l \Delta_i^m h_k + \Delta_i^m h_l \Delta_i^m g_k + \Delta_i^m h_l \Delta_i^m h_k] \right. \\
&\quad \left. + \psi(x_{i-1}^m) [\phi''_{lk}(y_i^m) - \phi''_{lk}(\mathbf{f}(x_{i-1}^m))] \Delta_i^m g_k \Delta_i^m g_l \right\}.
\end{aligned}$$

Then by (3.71), (3.72), (3.76) and (3.77), for all $m \geq m_1 \vee m_1 \vee m_3$,

$$\begin{aligned}
&\left| S_{LC}(\psi, \phi \circ \mathbf{f}; \lambda_m) - \sum_{k=1}^d S_{LC}(\psi(\phi'_k \circ \mathbf{f}), g_k; \lambda_m) - \sum_{k=1}^d A_k - \frac{1}{2} \sum_{l,k=1}^d B_{l,k} - C^- - C^+ \right| \\
&\leq \sum_{k=1}^d \left| A_k - S_{LC}(\psi(\phi'_k \circ \mathbf{f}), h_k; \lambda_m) \right| + \frac{1}{2} \sum_{l,k=1}^d \left| B_{l,k} - \sum_{i \in I_2(m)} (\psi(\phi''_{lk} \circ \mathbf{f}))(x_{i-1}^m) \Delta_i^m g_l \Delta_i^m g_k \right| \\
&+ \left| C^- + C^+ - \sum_{i \in I_1(m)} \psi(x_{i-1}^m) \left\{ \Delta_i^m(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f})(x_{i-1}^m) \Delta_i^m f_k \right\} \right| + |R_m| \leq d\epsilon + (5/2)d^2\epsilon + 4\epsilon + |R_m|.
\end{aligned}$$

To bound $|R_m|$ notice that for each $i \in I_2(m)$, $[x_{i-1}^m, x_i^m] \subset (z_{j-1}, z_j)$ for some $j = 1, \dots, s$. By (3.75) and Hölder's inequality, it then follows

$$\limsup_{m \rightarrow \infty} |R_m| \leq \frac{\sqrt{\epsilon}}{2} \sum_{l,k=1}^d \|\phi''_{lk} \circ \mathbf{f}\|_{\infty} (\sqrt{[g_l](b)} + \sqrt{[g_k](b)} + \sqrt{\epsilon}) + \frac{\epsilon}{2} \sum_{l,k=1}^d \sqrt{[g_k](b)[g_l](b)}.$$

Since $\epsilon > 0$ is arbitrary (3.70) exists and (3.68) holds with $z = a$ and $y = b$. The proof of statement (a) is complete.

The proof of statement (b) is analogous to the proof of statement (a). Instead of representation (3.78), in this case we have a telescoping representation

$$\begin{aligned}
S_{RC}(\psi, \phi \circ \mathbf{f}; \lambda_m) &= \sum_{i \in I_1(m)} \psi(x_i^m) \Delta_i^m(\phi \circ \mathbf{f}) \\
&+ \sum_{i \in I_2(m)} \psi(x_{i-1}^m) \left\{ \sum_{k=1}^d (\phi'_k \circ \mathbf{f})(x_i^m) \Delta_i^m f_k - \frac{1}{2} \sum_{l,k=1}^d \phi''_{lk}(y_i^m) \Delta_i^m f_l \Delta_i^m f_k \right\} \\
&= \sum_{k=1}^d S_{RC}(\psi(\phi'_k \circ \mathbf{f}), g_k; \lambda_m) + \sum_{k=1}^d S_{RC}(\psi(\phi'_k \circ \mathbf{f}), h_k; \lambda_m) \\
&- \frac{1}{2} \sum_{l,k=1}^d \left(\sum_{i=1}^{n(m)} - \sum_{i \in I_1(m)} \right) (\psi(\phi''_{lk} \circ \mathbf{f}))(x_i^m) \Delta_i^m g_l \Delta_i^m g_k \\
&+ \sum_{i \in I_1(m)} \psi(x_i^m) \left\{ \Delta_i^m(\phi \circ \mathbf{f}) - \sum_{k=1}^d (\phi'_k \circ \mathbf{f})(x_i^m) \Delta_i^m f_k \right\} + R_m
\end{aligned}$$

for $m \geq m_0$, where $y_i^m := \mathbf{f}(x_i^m) - \theta_i^m \Delta_i^m \mathbf{f}$, $\theta_i^m \in (0, 1)$ for $i = 1, \dots, n(m)$, and

$$\begin{aligned}
R_m &:= -\frac{1}{2} \sum_{l,k=1}^d \sum_{i \in I_2(m)} \left\{ \psi(x_i^m) \phi''_{lk}(y_i^m) [\Delta_i^m g_l \Delta_i^m h_k + \Delta_i^m h_l \Delta_i^m g_k + \Delta_i^m h_l \Delta_i^m h_k] \right. \\
&\quad \left. + \psi(x_i^m) [\phi''_{lk}(y_i^m) - \phi''_{lk}(\mathbf{f}(x_i^m))] \Delta_i^m g_k \Delta_i^m g_l \right\}.
\end{aligned}$$

This representation allows to prove the existence of the Right Cauchy λ -integral in (3.69) and relation (3.69). The proof of Theorem 34 is complete. $\square$

We finish this section with a chain rule which is used in Chapter 5 to prove the Black-Scholes option pricing formula. Let $U \subset \mathbb{R}$ be an open set and let $\phi: [a, b] \times U \mapsto \mathbb{R}$ be a continuous function. We say that ϕ is a C^2 class function on $[a, b] \times U$ if it is a C^2 class function on $(a, b) \times U$ and has continuous one-sided first and second derivatives at a . For such function ϕ and a function $f: [a, b] \mapsto U$, let

$$\Phi f(t) := \phi(t, f(t)), \quad t \in [a, b], \quad \Phi'_t f(t) := \begin{cases} \frac{\partial \phi}{\partial t}(t, f(t)) & \text{if } t \in [a, b), \\ 0 & \text{if } t = b, \end{cases} \quad (3.79)$$

$$\Phi'_x f(t) := \begin{cases} \frac{\partial \phi}{\partial x}(t, f(t)) & \text{if } t \in [a, b), \\ 0 & \text{if } t = b, \end{cases} \quad \Phi''_{xx} f(t) := \begin{cases} \frac{\partial^2 \phi}{\partial x^2}(t, f(t)) & \text{if } t \in [a, b), \\ 0 & \text{if } t = b. \end{cases} \quad (3.80)$$

The functions $\Phi'_t f$, $\Phi'_x f$ and $\Phi''_{xx} f$ so defined on $[a, b]$, may be discontinuous at b and unbounded.

Proposition 35 *For an open set $U \subset \mathbb{R}$ and $\lambda \in \Lambda[a, b]$, let $f: [a, b] \mapsto U$ be a continuous function having the quadratic λ -variation on $[a, b]$, and let $\phi: [a, b] \times U \mapsto \mathbb{R}$ be a continuous function such that its restriction to $[a, b) \times U$ is a C^2 class function. Suppose that there exist the Henstock-Kurzweil integrals $(HK) \int_a^b \Phi'_t f$ and $(HK) \int_a^b \Phi''_{xx} f d[f]_\lambda$. Then the Left Cauchy λ -integral $(LC) \int \Phi'_x f d_\lambda f$ is defined on $[a, b)$ and there exists the limit*

$$\lim_{u \uparrow b} (LC) \int_a^u \Phi'_x f d_\lambda f = \Phi f(b) - \Phi f(a) - (HK) \int_a^b \Phi'_t f - \frac{1}{2} (HK) \int_a^b \Phi''_{xx} f d[f]_\lambda. \quad (3.81)$$

Proof. By the proof of Theorem 34 with $d = 2$, $\psi \equiv 1$, $g_1 \equiv 0$, $h_1(x) \equiv x$, $g_2 \equiv f$ and $h_2 \equiv 0$, it follows that the Left Cauchy λ -integral $(LC) \int \Phi'_x f d_\lambda f$ is defined on $[a, b]$, and for each $a \leq u < b$,

$$\Phi f(u) - \Phi f(a) = (R) \int_a^u \Phi'_t f + (LC) \int_a^u \Phi'_x f d_\lambda f + \frac{1}{2} (RS) \int_a^u \Phi''_{xx} f d[f]_\lambda. \quad (3.82)$$

By the Hake theorem (see e.g. Section 7.3 in [77]), for a continuous function h with bounded variation on $[a, b]$, a function g is Henstock-Kurzweil integrable over $[a, b]$ with respect to h if and only if for each $a \leq u < b$, g is Henstock-Kurzweil integrable over $[a, u]$ with respect to h and there exists the limit $\lim_{u \uparrow b} (HK) \int_a^u g dh$. In this case,

$$\lim_{u \uparrow b} (HK) \int_a^u g dh = (HK) \int_a^b g dh.$$

Since the Henstock-Kurzweil integral extends the Riemann-Stieltjes integral, the second part of the conclusion follows. The proof is complete. $\square$

At this writing we are not able to prove under the same conditions that the Left Cauchy λ -integral in Proposition 35 is defined on the whole interval $[a, b]$. The method explored in this paper requires a convergence theorem for the Henstock-Kurzweil integral similar to the convergence theorem of Lemma 10, which cannot hold by Example 10 in [77, p. 213].

3.4 The product λ -integral

Let f be a real-valued function on $[a, b]$. Recall that the product integral on $[a, b]$ with respect to f is defined to be the limit

$$\mathfrak{P}_a^b(1 + df) := \lim_{\kappa, \mathfrak{P}} P(f; [a, b], \kappa)$$

provided it exists, where $P(f; [a, b], \kappa)$ is defined by (1.11). As it is mentioned earlier, by Theorem 4.4 of [23, Part II], if the product integral with respect to f exists then $f \in \mathcal{W}_2^*$. In this section we extend the definition of the product integral, so that the extended product integral exists with respect to functions having the quadratic λ -variation (Definition 38 below). But first we give a new proof of Theorem 4.4 of [23, Part II] for regulated functions which is based on a new characterization of the Wiener class $\mathcal{W}_2^*$ (Theorem 2.8 above).

The product integral. The product integral is more often used for non-commutative vector valued functions when no explicit formulas can be given. The product integral with respect to a *real-valued* function, or more generally with respect to a function with values in a commutative algebra, if exists, can be computed explicitly. Let f be a regulated function on $[a, b]$, and let $(u, v) \in S[[a, b]]$ be such that $u < v$. For each $\delta > 0$, let $D(u, v; \delta) := D_-(u, v; \delta) \cup D_+(u, v; \delta)$, where

$$\begin{cases} D_-(u, v; \delta) := \{x \in (a, b]: u < x \leq v, |\Delta^- f(x)| > \delta\} \\ D_+(u, v; \delta) := \{x \in [a, b): u \leq x < v, |\Delta^+ f(x)| > \delta\}. \end{cases}$$

Here we use the order on the extended interval defined in Section 2.1. Also, let

$$\prod_{D(u, v; \delta)} (1 + \Delta f) e^{-\Delta f} := \prod_{D_-(u, v; \delta)} (1 + \Delta^- f) e^{-\Delta^- f} \prod_{D_+(u, v; \delta)} (1 + \Delta^+ f) e^{-\Delta^+ f}.$$

Define the function $\gamma(f)$ on $S[[a, b]]$ by

$$\gamma(f; u, v) := \prod_{(u, v)} (1 + \Delta f) e^{-\Delta f} := \lim_{\delta \downarrow 0} \prod_{D(u, v; \delta)} (1 + \Delta f) e^{-\Delta f} \quad (3.83)$$

provided $(u, v) \in S[[a, b]]$, $u < v$, and the infinite product converges absolutely. If $u = v$ then let $\gamma(f; u, u) := 1$. Now the necessary and sufficient conditions for the existence of the product integral can be formulated as follows:

Theorem 36 *The following two statements about a regulated function f on $[a, b]$ are equivalent:*

- (a) *the product integral $\mathfrak{P}_a^b(1 + df)$ exists and is non-zero;*
- (b) *$f \in \mathcal{W}_2^*[a, b]$ and*

$$\Delta^- f(x) \neq -1 \neq \Delta^+ f(y) \quad \text{for each } a \leq y < x \leq b. \quad (3.84)$$

Moreover, if any one of the two statements holds then the product integral extends to the function

$$S[[a, b]] \ni (u, v) \mapsto \mathfrak{P}_u^v(1 + df) = \exp\{f(v) - f(u)\} \prod_{(u, v)} (1 + \Delta f) e^{-\Delta f}, \quad (3.85)$$

which is nondegenerate, bounded, multiplicative and upper continuous function on $S[[a, b]]$.

The function $\gamma(f)$ defined by (3.83) is a discontinuous part of values (3.85) and it exists under the weaker condition $\sigma_2(f) < \infty$ (see (2.7) for its definition) than above condition (b).

Lemma 37 *Let f be a regulated function on $[a, b]$ such that $\sigma_2(f) < \infty$. The function $\gamma(f)$ on $S[[a, b]]$ is well defined, multiplicative, bounded, and for any $(u, v) \in S[[a, b]]$ such that f has no jumps on $[[u, v]]$ exceeding $1/2$,*

$$|\gamma(f; u, v) - 1| \leq 2e^{2\sigma_2(f)} \sum_{[[u, v]]} [\Delta f]^2. \quad (3.86)$$

Moreover, $\gamma(f)$ is nondegenerate if and only if (3.84) holds. If $\gamma(f)$ is nondegenerate then it is upper continuous.

Proof. First we show that $\gamma(f)$ is well defined. By a Taylor series expansion with remainder, we have that for $|u| \leq 1/2$,

$$\xi(u) := (1 + u)e^{-u} = 1 - \frac{\theta(u)}{2}u^2, \quad (3.87)$$

where $1/(2\sqrt{e}) \leq \theta(u) \leq 3\sqrt{e}/2$. Thus

$$\sum_{[a, b]} |1 - \xi(\Delta^- f)| 1_{\{|\Delta^- f| \leq 1/2\}} + \sum_{(a, b]} |1 - \xi(\Delta^+ f)| 1_{\{|\Delta^+ f| \leq 1/2\}} \leq \frac{3\sqrt{e}}{4} \sigma_2(f)^2 < \infty.$$

Since f is regulated there exists a partition $\{z_k: k = 0, \dots, l\}$ of $[a, b]$ such that

$$\text{Osc}(f; (z_{k-1}, z_k)) \leq 1/2 \quad \text{for } k = 1, \dots, l. \quad (3.88)$$

Let $(u, v) \in S[[a, b]]$. By the preceding bounds, each product $\gamma(f; z_{k-1}+, z_k-)$, $k = 1, \dots, l$, converges absolutely, and hence so does the product $\gamma(f; u, v)$. Therefore the function $\gamma(f)$ is well defined.

For each $\delta > 0$, the map

$$S[[a, b]] \ni (u, v) \mapsto \prod_{D(u, v; \delta)} (1 + \Delta f) e^{-\Delta f}$$

is multiplicative. Thus the function $\gamma(f)$ is multiplicative as a limit of multiplicative functions. Next we show that $\gamma(f)$ is bounded. A Taylor series expansion with remainder gives for $|u| \leq 1/2$,

$$\log(1 + u) = u - \frac{\theta(u)}{2} u^2, \quad (3.89)$$

where $4/9 \leq \theta(u) \leq 4$. Let $(u, v) \in S[[a, b]]$ be such that jumps of f on $[[u, v]]$ do not exceed $1/2$, and let $0 < \delta \leq 1/2$. By (3.89), we have

$$\begin{aligned} \prod_{D(u, v; \delta)} (1 + \Delta f) e^{-\Delta f} &= \exp \left\{ -\frac{1}{2} \sum_{D(u, v; \delta)} [\theta(\Delta^- f)(\Delta^- f)^2 + \theta(\Delta^+ f)(\Delta^+ f)^2] \right\} \\ &\leq \exp \left\{ -\frac{2}{9} \sum_{D(u, v; \delta)} [(\Delta^- f)^2 + (\Delta^+ f)^2] \right\}. \end{aligned} \quad (3.90)$$

Letting $\delta \downarrow 0$ on both sides, we obtain the upper bound

$$\gamma(f; u, v) \leq \exp \left\{ -\frac{2}{9} \sum_{[[u, v]]} [(\Delta^- f)^2 + (\Delta^+ f)^2] \right\} \leq 1.$$

For any $(u, v) \in S[[a, b]]$, applying these bounds to the intersections $[[u, v]] \cap [[z_{k-1}+, z_k-]]$, where $\{z_k: k = 0, \dots, l\}$ is defined by (3.88), and using notation (3.87) we get

$$\|\gamma(f)\|_\infty \leq \prod_{k=1}^l \left[1 \vee |\xi(\Delta^+ f(z_{k-1}))| \right] \left[1 \vee |\xi(\Delta^- f(z_k))| \right] < +\infty,$$

proving that $\gamma(f)$ is bounded. Let $(u, v) \in S[[a, b]]$ be such that jumps of f on $[[u, v]]$ do not exceed $1/2$, and let $0 < \delta \leq 1/2$. An application of the inequality $|e^x - 1| \leq |x|e^{|x|}$ for $x \in \mathbb{R}$ and (3.89) imply that

$$\begin{aligned} \left| \prod_{D(u, v; \delta)} (1 + \Delta f) e^{-\Delta f} - 1 \right| &= \left| \exp \left\{ \sum_{D(u, v; \delta)} [\log(1 + \Delta f) - \Delta f] \right\} - 1 \right| \\ &\leq 2e^{2\sigma_2(f)^2} \sum_{[[u, v]]} [(\Delta^- f)^2 + (\Delta^+ f)^2]. \end{aligned}$$

Letting $\delta \downarrow 0$ on the left side, we obtain that (3.86) holds.

Second we show that $\gamma(f)$ is nondegenerate on $[a, b]$ if and only if (3.84) holds. Clearly if $\gamma(f)$ is nondegenerate on $[a, b]$ then (3.84) holds. To show the converse let $(x, y) \subset [a, b]$ be an open interval such that f have no jumps on (x, y) bigger than $1/2$, and let $0 < \delta \leq 1/2$. In the

left side of (3.90) using the upper bound for θ (see (3.89)), and then letting $\delta \downarrow 0$ on both sides of the resulting inequality, we get the lower bound

$$\gamma(f; x+, y-) \geq \exp \left\{ -2 \sum_{(x,y)} [(\Delta^- f)^2 + (\Delta^+ f)^2] \right\}$$

for any open interval (x, y) which does not contain discontinuities of f with jumps bigger than $1/2$. Applying this bound to each interval (z_{k-1}, z_k) satisfying (3.88), and using notation (3.87) we get

$$|\gamma(f; a, b)| \geq \left| \prod_{k=1}^l \xi(\Delta^+ f(z_{k-1})) \xi(\Delta^- f(z_k)) \right| \exp \left\{ -2 \sum_{k=1}^l \sigma_2(f; (z_{k-1}, z_k))^2 \right\}.$$

Since $\sigma_2(f) < \infty$ and (3.84) holds, $\gamma(f; a, b) \neq 0$.

Finally we show that $\gamma(f)$ is upper continuous provided it is nondegenerate. To this aim we show that statement (vii) of Theorem 2.4 holds. For $a \leq y < x \leq b$ with $|\Delta^- f(x)| \leq 1/2$ and $|\Delta^+ f(y)| \leq 1/2$, by (3.87),

$$|\gamma(f; x-, x) - 1| \leq \frac{3\sqrt{e}}{8} (\Delta^- f(x))^2 \quad \text{and} \quad |\gamma(f; y, y+) - 1| \leq \frac{3\sqrt{e}}{8} (\Delta^+ f(y))^2.$$

Thus (2.5) holds for $\mu = \gamma(f)$ by the analogous property for the regulated function f . Suppose that extended open intervals $\llbracket x_k+, y_k- \rrbracket \downarrow \llbracket u, u \rrbracket$ as $k \rightarrow \infty$, for some $u \in \llbracket a, b \rrbracket$. Then either $y_k = y$ and $x_k \uparrow y$ for some $y \in (a, b]$ and all sufficiently large k , or $x_k = x$ and $y_k \downarrow x$ for some $x \in [a, b)$ and all sufficiently large k . In either case for all sufficiently large k , $\llbracket x_k+, y_k- \rrbracket$ does not contain discontinuity points with jumps bigger than $1/2$. Since $\sigma_2(f) < \infty$, by (3.86), it then follows that

$$\limsup_{k \rightarrow \infty} |\gamma(f; x_k+, y_k-) - 1| \leq 2e^{2\sigma_2(f)^2} \limsup_{k \rightarrow \infty} \sum_{(x_k, y_k)} [(\Delta^- f)^2 + (\Delta^+ f)^2] = 0.$$

Thus statement (vii) of Theorem 2.4 holds for $\mu = \gamma(f)$, and hence $\gamma(f)$ is upper continuous. The proof of Lemma 37 is now complete. $\square$

Proof of Theorem 36. We start with a useful representation of the product $P(f; \kappa)$ defined by (1.11). Let $\zeta = \{z_j: j = 0, \dots, k\}$ be a partition of $[a, b]$ to be chosen later, and let $\kappa = \{x_i: i = 0, \dots, n\}$ be a refinement of ζ such that each intersection $(z_{j-1}, z_j) \cap \kappa$, $j = 1, \dots, k$, contains at least two different points. For $j = 0, \dots, k$, let $i(j) \in \{0, \dots, n\}$ be such that $x_{i(j)} = z_j$. Then let

$$I_1(\zeta, \kappa) := \{i(j-1) + 1, i(j): j = 1, \dots, k\} \subset \{1, \dots, n\} \quad \text{and} \quad I_2(\zeta, \kappa) := \{1, \dots, n\} \setminus I_1(\zeta, \kappa).$$

For a function f on $[a, b]$, let $\Delta_i f := f(x_i) - f(x_{i-1})$, $i = 1, \dots, n$. If $|\Delta_i f| \leq 1/2$ for each $i \in I_2(\zeta, \kappa)$, then by (3.89),

$$\begin{aligned} P(f; \kappa) &= \exp \left\{ \sum_{i \in I_2(\zeta, \kappa)} \Delta_i f - \frac{1}{2} \sum_{i \in I_2(\zeta, \kappa)} \theta_i (\Delta_i f)^2 \right\} \prod_{i \in I_1(\zeta, \kappa)} (1 + \Delta_i f) \\ &= \exp \left\{ f(b) - f(a) - \frac{1}{2} \sum_{i \in I_2(\zeta, \kappa)} \theta_i (\Delta_i f)^2 \right\} \prod_{i \in I_1(\zeta, \kappa)} \left\{ (1 + \Delta_i f) e^{-\Delta_i f} \right\}, \end{aligned} \quad (3.91)$$

where $\theta_i := \theta(\Delta_i f)$ for $i \in I_2(\zeta, \kappa)$ and $\theta(u) = (1+v)^{-2}$ for some v between u and 0.

Suppose that statement (a) holds. First we show that (3.84) holds. Since $\mathfrak{Y}_a^b(1+df) \neq 0$, there exist $\epsilon > 0$ and a partition $\zeta = \{z_j: j = 0, \dots, k\}$ of $[a, b]$ such that $D(a, b; 1/2) \subset \zeta$ and $|P(f; \kappa)| \geq \epsilon$ for each refinement κ of ζ . Suppose that $\Delta^- f(x) = -1$ for some $x \in (a, b]$. Then $x = z_j \in \zeta$ for some $j \in \{1, \dots, k\}$. Let κ be a refinement of ζ such that $z_{j-1} < x_{i(j)-1} < z_j (= x_{i(j)})$. Letting $x_{i(j)-1} \uparrow z_j$ and keeping other $x_i \in \kappa$ unchanged, we have that $(1 + \Delta_{i(j)-1} f)(1 + \Delta_{i(j)} f) \rightarrow 0$. Thus $P(f; \kappa) \rightarrow 0$, a contradiction. Similarly, the assumption that $\Delta^+ f(y) = -1$ for some $y \in [a, b)$ leads to a contradiction. Therefore (3.84) must hold.

Next we show that $v_2(f) < \infty$. This is so if $\sum_{j=1}^k v_2(f; (z_{j-1}, z_j)) < \infty$ for some partition $\{z_j: j = 0, \dots, k\}$ of $[a, b]$. Again, since f is regulated and $\mathfrak{Y}_a^b(1+df) \neq 0$, there exist $\epsilon > 0$ and a partition $\zeta = \{z_j: j = 0, \dots, k\}$ of $[a, b]$ such that $D(a, b; 1/2) \subset \zeta$, $\text{Osc}(f; (z_{j-1}, z_j)) \leq 1/2$ for $j = 1, \dots, k$, and $|P(f; \kappa)| \geq \epsilon$ for each refinement κ of ζ . Since $|f|$ is bounded by a finite constant M , by (3.91), we have that

$$0 < \epsilon \leq \exp \left\{ f(b) - f(a) - \frac{2}{9} \sum_{i \in I_2(\zeta, \kappa)} (\Delta_i f)^2 \right\} \left[(1 + 2M)e^{2M} \right]^{2k}$$

for all refinements κ of ζ . If $\sum_{j=1}^k v_2(f; (z_{j-1}, z_j)) = +\infty$ then the right side of the preceding bound can be made arbitrarily small. The contradiction yields that $v_2(f) < \infty$.

To prove that $f \in \mathcal{W}_2^*[a, b]$, let $\epsilon \in (0, 1/2]$. Since f is regulated and since the product integral with respect to f is defined, there exists a partition $\kappa_0 = \{y_r: r = 0, \dots, l\}$ of $[a, b]$ such that

$$\max_{1 \leq r \leq l} \text{Osc}(f; (y_{r-1}, y_r)) < \epsilon \quad \text{and} \quad \left| \log \mathfrak{Y}_a^b(1+df) - \log P(f; \kappa) \right| < \epsilon$$

for each refinement κ of κ_0 . By Lemma 2.7, $\sigma_2(f) \leq v_2(f) < +\infty$. Thus by Lemma 37, there exists a $\delta_0 \in (0, 1/2]$ such that

$$\left| \log \gamma(f; a, b) - \log \left\{ \prod_{D(a, b; \delta)} (1 + \Delta f) e^{-\Delta f} \right\} \right| < \epsilon \quad (3.92)$$

for each $0 < \delta \leq \delta_0$. Let δ be the minimum of the subset of non-zero values in the set

$$\{\delta_0, |\Delta^+ f(a)|, |\Delta^- f(b)|, |\Delta^- f(x)|, |\Delta^+ f(x)|: x \in (a, b) \cap \kappa_0\}, \quad (3.93)$$

and let $\zeta = \{z_j: j = 0, \dots, k\} := \kappa_0 \cup D(a, b; \delta)$. Then choose $\{u_{j-1}, v_j: j = 1, \dots, k\} \subset (a, b)$ such that for $j = 1, \dots, k$, $z_{j-1} < u_{j-1} < v_j < z_j$,

$$\text{Osc}([f - f(z_{j-1})]^2; (z_{j-1}, u_{j-1})) < \epsilon/2k, \quad \text{Osc}([f(z_j) - f]^2; (v_j, z_j)) < \epsilon/2k,$$

and for each refinement κ of $\kappa_1 := \zeta \cup \{u_{j-1}, v_j: j = 1, \dots, k\}$,

$$\left| \log \left\{ \prod_{D(a, b; \delta)} (1 + \Delta f) e^{-\Delta f} \right\} - \log \left\{ \prod_{i \in I_1(\zeta, \kappa)} (1 + \Delta_i f) e^{-\Delta_i f} \right\} \right| < \epsilon. \quad (3.94)$$

Here we use the notation introduced at the beginning of the proof. Letting

$$A := f(b) - f(a) + \log \gamma(f; a, b) - \log \mathfrak{Y}_a^b(1+df) \quad (3.95)$$

and using (3.91), we then have the bound

$$\begin{aligned}
& \left| A - \frac{1}{2} \sum_{i \in I_2(\zeta, \kappa)} \theta_i(\Delta_i f)^2 \right| \\
&= \left| A - [f(b) - f(a) + \log \left\{ \prod_{i \in I_1(\zeta, \kappa)} (1 + \Delta_i f) e^{-\Delta_i f} \right\} - \log P(f; \kappa)] \right| \\
&\leq \left| \log \gamma(f; [a, b]) - \log \left\{ \prod_{i \in I_1(\zeta, \kappa)} (1 + \Delta_i f) e^{-\Delta_i f} \right\} \right| + \left| \log \mathfrak{Y}_a^b(1 + df) - \log P(f; \kappa) \right| < 3\epsilon
\end{aligned}$$

for each refinement κ of κ_1 . Since $|\Delta_i f| < \epsilon$ for $i \in I_2(\zeta, \kappa)$, it then follows that

$$2(1 - \epsilon)^2(A - 3\epsilon) < \sum_{i \in I_2(\zeta, \kappa)} (\Delta_i f)^2 < 2(1 + \epsilon)^2(A + 3\epsilon)$$

for each refinement κ of κ_1 . Thus for any two refinements $\kappa' = \{x'_i\}$ and $\kappa'' = \{x''_i\}$ of κ_1 ,

$$\begin{aligned}
& |s_2(f; \kappa') - s_2(f; \kappa'')| \leq \sum_{j=1}^k \left| [f(x'_{i(j-1)+1}) - f(z_{j-1})]^2 - [f(x''_{i(j-1)+1}) - f(z_{j-1})]^2 \right| \\
&+ \sum_{j=1}^k \left| [f(z_j) - f(x'_{i(j)-1})]^2 - [f(z_j) - f(x''_{i(j)-1})]^2 \right| + \left| \sum_{i \in I_2(\zeta, \kappa')} (\Delta'_i f)^2 - \sum_{i \in I_2(\zeta, \kappa'')} (\Delta''_i f)^2 \right| \\
&< \epsilon + 8\epsilon A + 12\epsilon(1 + \epsilon^2) \leq \epsilon + 8\epsilon(3 + A),
\end{aligned}$$

where $\Delta'_i f := f(x'_i) - f(x'_{i-1})$ and $\Delta''_i f := f(x''_i) - f(x''_{i-1})$. Hence

$$0 \leq v_2^*(f)^2 - \sigma_2^*(f)^2 \leq \sup_{\kappa \supset \kappa_1} s_2(f; \kappa) - \inf_{\kappa \supset \kappa_1} s_2(f; \kappa) \leq \epsilon + 8\epsilon(3 + A).$$

Since ϵ is arbitrary, $f \in \mathcal{W}_2^*[a, b]$. Thus (b) follows from (a).

Now suppose that statement (b) holds. The following proof of existence of the product integral is a variation of the preceding argument showing that A defined by (3.95) actually is zero. Let $\epsilon \in (0, 1/4]$. By Lemma 2.9, there exists a partition $\kappa_0 = \{y_r: r = 0, \dots, l\}$ of $[a, b]$ such that

$$\sum_{r=1}^l v_2(f; (y_{r-1}, y_r)) < \epsilon.$$

In particular, we have that $\text{Osc}(f; (y_{r-1}, y_r)) < \sqrt{\epsilon} \leq 1/2$ for each $r = 1, \dots, l$. By Lemma 37, $\gamma(f; a, b) \neq 0$. Using Lemma 37 again we find a $\delta_0 > 0$ such that (3.92) holds, and then choose a $\delta \in (0, \delta_0]$ to be the minimum of the subset of non-zero values in the set (3.93). As before, let $\zeta := \{z_j: j = 0, \dots, k\} := \kappa_0 \cup D(a, b; \delta)$. Then choose $\{u_{j-1}, v_j: j = 1, \dots, k\} \subset (a, b)$ such that $z_{j-1} < u_{j-1} < v_j < z_j$ for $j = 1, \dots, k$, and (3.94) holds for each refinement κ of $\kappa_1 := \zeta \cup \{u_{j-1}, v_j: j = 1, \dots, k\}$. By (3.91), it then follows that

$$\begin{aligned}
& \left| \log P(f; \kappa) - [f(b) - f(a) + \log \gamma(f; a, b)] \right| \\
&\leq \left| \log \left\{ \prod_{i \in I_1(\zeta, \kappa)} (1 + \Delta_i f) e^{-\Delta_i f} \right\} - \log \gamma(f; a, b) \right| + \frac{2}{9} \sum_{i \in I_2(\zeta, \kappa)} (\Delta_i f)^2 < 2\epsilon + \frac{2}{9}\epsilon
\end{aligned}$$

for each refinement of κ_1 . Thus $\mathfrak{Y}_a^b(1 + df)$ exists and equals to the right side of (3.85) with $[u, v] = [a, b]$. Since $\gamma(f; a, b) \neq 0$, (a) follows from (b).

Since the interval function $\mu(f)$ on $S[[a, b]]$ defined by (2.2) is additive and upper continuous, by Lemma 37, the map

$$S[[a, b]] \ni (u, v) \mapsto \exp \{ \mu(f; u, v) \} \gamma(f; u, v) =: \pi(f; u, v)$$

is nondegenerate, bounded, multiplicative and upper continuous function on $S[[a, b]]$. The second part of the theorem will follow once we show that for each $(u, v) \in S[[a, b]]$, the product integral $\mathfrak{Y}_u^v(1 + df)$ exists and equals $\pi(f; u, v)$. For the intervals $[[u, v] = [x-, x]]$ and $[[u, v] = [y, y+]]$, we have

$$\mathfrak{Y}_{x-}^x(1 + df) = 1 + \Delta^- f(x) = \pi(f; x-, x) \quad \text{and} \quad \mathfrak{Y}_y^{y+}(1 + df) = 1 + \Delta^+ f(y) = \pi(f; y, y+).$$

Due to multiplicativity of the product integral it is enough to prove the existence of $\mathfrak{Y}_u^v(1 + df)$ when $[[u, v] = [x+, y-]]$ for some $a \leq x < y \leq b$. The proof in this case is the same as for $[u, v] = [a, b]$ except that a partition $\zeta = \{z_j: j = 0, \dots, k\}$ is such that $z_0 > x$ if $u = x+$ and $z_k < y$ if $v = y-$. Also, κ_1 is the same as before except that in the cases $x < z_0$ and $z_k < y$ we include additional points $u_{-1} \in (x, z_0)$ and $v_{k+1} \in (z_k, y)$, respectively. By the proof it also follows that the product integral $\mathfrak{Y}_u^v(1 + df)$ has the value $\pi(f; u, v)$. Thus $\mathfrak{Y}(1 + df)$ is the interval function on $S[[a, b]]$ which agree with the interval function $\pi(f)$. The proof of Theorem 36 is now complete. $\square$

The product λ -integral. By Theorem 36, if the product integral with respect to a regulated function f exists then $f \in \mathcal{W}_2^*$. Thus Example 6 shows that the product integral may not exist with respect to a function having the quadratic λ -variation. Next we extend the product integral so as to accommodate a product integration with such a situation. Recall that $P(f; \kappa)$ is defined by (1.11) and the trace partition $\kappa \pitchfork [[u, v]]$ is defined in Notation 2.

Definition 38 Let f be a regulated function on $[a, b]$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. We say that the *product λ -integral* $\mathfrak{Y}(1 + d_\lambda f)$ is defined on $[a, b]$ if there exists a nondegenerate, bounded, multiplicative and upper continuous function $\pi_\lambda(f)$ on $S[[a, b]]$ such that for each $(u, v) \in S[[a, b]]$,

$$\pi_\lambda(f; u, v) = \lim_{m \rightarrow \infty} P(f; \lambda_m \pitchfork [[u, v]]). \quad (3.96)$$

The right distribution function of $\pi_\lambda(f)$ restricted to $[a, b]$, that is, $\mathcal{P}_\lambda f(x) := \pi(f; a, x)$ for $a \leq x \leq b$, is called the *indefinite product λ -integral*.

Alternatively the product λ -integral can be defined solely in terms of a regulated function on $[a, b]$ as follows:

Proposition 39 Let f be a regulated function on $[a, b]$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. The product λ -integral $\mathfrak{Y}(1 + d_\lambda f)$ is defined on $[a, b]$ if and only if there exists a regulated function G on $[a, b]$ such that $G(a) = 1$, $|G| \gg 0$ and for $a \leq s < t \leq b$,

$$G(t)/G(s) = \lim_{m \rightarrow \infty} P(f; \lambda_m \pitchfork [s, t]), \quad (3.97)$$

$$G(t)/G(t-) = 1 + (\Delta^- f)(t) \quad \text{and} \quad G(s+)/G(s) = 1 + (\Delta^+ f)(s). \quad (3.98)$$

If the two equivalent assertions hold then $\pi_\lambda(f; u, v) = G(v)/G(u)$ for each $(u, v) \in S[[a, b]]$.

Proof. First suppose that there exists a regulated function G on $[a, b]$ equal 1 at a , $|G| \gg 0$, and which satisfy both (3.97) and (3.98). Define a function $\pi_\lambda(f)$ on $S[[a, b]]$ by $\pi_\lambda(f; u, v) := G(v)/G(u)$ for $(u, v) \in S[[a, b]]$. Then $\pi_\lambda(f)$ is nondegenerate, bounded, multiplicative and upper continuous function on $[[a, b]]$ by Theorem 2.4. We prove (3.96) only for the cases $(u, v) = (a, t-)$ and $(u, v) = (a, t+)$ because the proofs for the other cases are similar. To this aim let $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$, $m = 1, 2, \dots$, and first let $t \in (a, b]$. For each $m \geq 1$, there is an index $i(t) = i_m(t) \in \{1, \dots, n(m) - 1\}$ such that $x_{i(t)}^m < t \leq x_{i(t)+1}^m$. Then by (3.97) and (3.98), we have

$$\begin{aligned} \lim_{m \rightarrow \infty} P(f; \lambda_m \cap [[a, t-]]) &= \lim_{m \rightarrow \infty} \left\{ P(f; \lambda_m \cap [a, t]) \frac{[1 + f(t-) - f(x_{i(t)}^m)]}{[1 + f(t) - f(x_{i(t)}^m)]} \right\} \\ &= \frac{G(t)}{1 + (\Delta^- f)(t)} = G(t-) = \pi_\lambda(f; a, t-), \end{aligned}$$

proving (3.96) for $(u, v) = (a, t-)$. Now let $t \in [a, b)$. By the definition of the trace partition, we have for each $m \geq 1$,

$$P(f; \lambda_m \cap [[a, t+]]) = P(f; \lambda_m \cap [a, t])(1 + \Delta^+ f(t)),$$

proving (3.96) for $(u, v) = (a, t+)$.

To prove the converse implication let $G(t) := \pi_\lambda(f; a, t)$ for $t \in [a, b]$. By the statement (iv) of Theorem 2.4 and (3.96), G is regulated on $[a, b]$, it is 1 at a , $|G(t)| \geq C$ with $C := \inf\{|\pi_\lambda(f; a, t)|: t \in [a, b]\} > 0$, and satisfies (3.97) and (3.98). The proof of Proposition 39 is complete. $\square$

Next we show that the product λ -integral extends the product integral for $\lambda \in \Lambda$ such that (1.19) holds.

Proposition 40 *Let f be a regulated function on $[a, b]$, and let $\lambda \in \Lambda[a, b]$ be such that the two-sided discontinuity points of f are accessible by λ . If the product integral $\mathfrak{P}(1 + df)$ exists and is non-zero on $[a, b]$ then the product λ -integral $\mathfrak{P}(1 + d_\lambda f)$ is defined on $[a, b]$, and both have the same values on $S[[a, b]]$.*

Proof. By the second part of Theorem 36 and by Proposition 39, it is enough to prove that for each $(s, t) \in S[a, b]$,

$$\mathfrak{P}_s^t(1 + df) = \lim_{m \rightarrow \infty} P(f; \lambda_m \cap [s, t]).$$

Given $(s, t) \in S[a, b]$, let $\epsilon > 0$. Then there is a partition κ_0 of $[s, t]$ such that

$$\left| \mathfrak{P}_s^t(1 + df) - P(f; \kappa) \right| < \epsilon$$

for each refinement κ of κ_0 . If $\kappa_0 \subset \cup_m \lambda_m$ then $P(f; \lambda_m \cap [s, t])$ is within ϵ from $\mathfrak{P}_s^t(1 + df)$ for all sufficiently large m . Suppose that $\zeta = \{z_j: j = 1, \dots, k\} := \kappa_0 \setminus (\cup_m \lambda_m \cup \{s, t\})$ is non-empty. For $j = 1, \dots, k$ and all m such that each $(z_{j-1}, z_j) \cap \lambda_m$ is non-empty, let $i(j) = i(j, m) \in \{1, \dots, n(m)\}$ be such that $x_{i(j)-1}^m < z_j < x_{i(j)}^m$. Also let $\Delta_{i(j)}^m f := f(x_{i(j)}^m) - f(x_{i(j)-1}^m)$, $\Delta_{i(j)}^{m,1} f := f(z_j) - f(x_{i(j)-1}^m)$ and $\Delta_{i(j)}^{m,2} f := f(x_{i(j)}^m) - f(z_j)$. By a telescoping sum

$$\prod_{j=1}^k a_j - \prod_{j=1}^k b_j = \sum_{j=1}^k \left(\prod_{i=j+1}^k a_i \right) (a_j - b_j) \left(\prod_{i=1}^{j-1} b_i \right),$$

we have the relation

$$P(f; (\lambda_m \mathbin{\frown} [s, t]) \cup \zeta) - P(f; \lambda_m \mathbin{\frown} [s, t]) \\ = P(f; (\lambda_m \mathbin{\frown} [s, t]) \cup \zeta) \sum_{j=1}^k \left\{ \frac{\Delta_{i(j)}^{m,1} f \Delta_{i(j)}^{m,2} f}{[1 + \Delta_{i(j)}^{m,1} f][1 + \Delta_{i(j)}^{m,2} f]} \prod_{l=1}^{j-1} \frac{1 + \Delta_{i(l)}^m f}{[1 + \Delta_{i(l)}^{m,1} f][1 + \Delta_{i(l)}^{m,2} f]} \right\}$$

for all sufficiently large m , where the products over the empty set of indices are equal to 1. By Theorem 36(b), $\Delta^- f(z_j) \neq -1 \neq \Delta^+ f(z_j)$. By (1.19), $\Delta_{i(j)}^{m,1} f \Delta_{i(j)}^{m,2} f \rightarrow 0$ as $m \rightarrow \infty$. Thus each term of the sum on the right side tends to 0 as $m \rightarrow \infty$. Therefore since for all sufficiently large m , $P(f; (\lambda_m \mathbin{\frown} [s, t]) \cup \zeta)$ is within ϵ from the value of the product integral over $[s, t]$,

$$\limsup_{m \rightarrow \infty} \left| \mathfrak{Y}_s^t(1 + df) - P(f; \lambda_m \mathbin{\frown} [s, t]) \right| \leq \epsilon,$$

proving the proposition. $\square$

Suppose that a function f on $[a, b]$ has the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$. Thus by Definition 3, there is an additive and upper continuous function $\alpha_\lambda(f)$ on $S[[a, b]]$ such that $\alpha_\lambda(f; u, v) = [f]_\lambda(v) - [f]_\lambda(u)$ for $(u, v) \in S[[a, b]]$. Define the additive upper continuous function $\alpha_\lambda^c(f)$ by

$$\alpha_\lambda^c(f; u, v) := [f]_\lambda^c(v) - [f]_\lambda^c(u), \quad (u, v) \in S[[a, b]]. \quad (3.99)$$

Then by (3.6), for $(u, v) \in S[a, b]$, we have the decomposition

$$\alpha_\lambda(f; u, v) = \alpha_\lambda^c(f; u, v) + \sum_{(u, v]} \{\Delta^- f\}^2 + \sum_{[u, v)} \{\Delta^+ f\}^2. \quad (3.100)$$

Define the function $\beta_\lambda(f)$ on $S[[a, b]]$ by

$$\beta_\lambda(f; u, v) := \exp \{ \mu(f; u, v) - \frac{1}{2} \alpha_\lambda^c(f; u, v) \} \gamma(f; u, v), \quad (u, v) \in S[[a, b]], \quad (3.101)$$

where $\mu(f)$, $\alpha_\lambda^c(f)$ and $\gamma(f)$ are defined by (2.2), (3.99) and (3.83), respectively. By the definition of $\gamma(f)$, it follows that for $a \leq s < t \leq b$,

$$\beta_\lambda(f; t-, t) = 1 + \Delta^- f(t) \quad \text{and} \quad \beta_\lambda(f; s, s+) = 1 + \Delta^+ f(s). \quad (3.102)$$

The following is a consequence of Lemma 37.

Corollary 41 *If f has the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$ and (3.84) holds, then $\beta_\lambda(f)$ is nondegenerate, bounded, multiplicative and upper continuous function on $S[[a, b]]$.*

Theorem 42 *Let f be a regulated function on $[a, b]$ having the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$ such that the two-sided discontinuity points of f are accessible by λ . If also (3.84) holds then the product λ -integral $\mathfrak{Y}(1 + d_\lambda f)$ is defined on $[a, b]$, and equals $\beta_\lambda(f)$.*

Proof. For $x \in [a, b]$, let $G(x) := \beta_\lambda(f; a, x)$. By Theorem 2.4 and Corollary 41, G is a regulated function on $[a, b]$ such that $G(a) = 1$ and (3.98) holds for each $a \leq x < y \leq b$. Thus by Proposition 39, it is enough to prove that for each $a \leq x < y \leq b$,

$$\lim_{m \rightarrow \infty} \log P(f; \lambda_m \mathbin{\frown} [x, y]) = \mu(f; x, y) - \frac{1}{2} \alpha_\lambda^c(f; x, y) + \log \gamma(f; x, y). \quad (3.103)$$

To begin the proof, let $a \leq x < y \leq b$ and let $\epsilon > 0$. Since f is regulated there exists a partition $\kappa = \{t_r: r = 0, \dots, l\}$ of $[x, y]$ such that

$$\max_{1 \leq r \leq l} \text{Osc}(f; (t_{r-1}, t_r)) < \epsilon. \quad (3.104)$$

Since f has the quadratic λ -variation, by Lemma 37, there exists $\delta_0 > 0$ such that for each $0 < \delta \leq \delta_0$ and $\sigma := D(x, y; \delta)$,

$$\left| \log \gamma(f; x, y) - \log \left\{ \prod_{\sigma} (1 + \Delta f) e^{-\Delta f} \right\} \right| < \epsilon \quad \text{and} \quad \left| \sum_{[x, y]} (\Delta f)^2 - \sum_{\sigma} (\Delta f)^2 \right| < \epsilon. \quad (3.105)$$

Let δ be the minimal number among the positive values in the set $\{\delta_0, |\Delta^- f(t_r)|, |\Delta^+ f(t_{r-1})|: r = 1, \dots, l\}$, and let $\zeta = \{z_j: j = 0, \dots, k\} := \kappa \cup D(x, y; \delta)$. Thus if $z_j \in \zeta$ and f has a jump at z_j then $z_j \in D(x, y; \delta)$. Let $\zeta_0 := \zeta \setminus \cup_m \lambda_m$, and let m_0 be the minimal integer such that $\zeta \setminus \zeta_0 \subset \lambda_{m_0}$ and each $(z_{j-1}, z_j) \cap \lambda_{m_0}$ contains at least two points. By (1.19), if $z_j \in \zeta_0$ then f cannot have jumps from the two sides at z_j . Suppose that each $\lambda_m \cap [x, y] = \{t_i^m: i = i_m(x) - 1, \dots, i_m(y)\}$, and let $\Delta_i^m f := f(t_i^m) - f(t_{i-1}^m)$ for each $i \in I(m) := \{i_m(x), \dots, i_m(y)\}$ and $m \geq m_0$. For each $m \geq m_0$, if $z_j \in \zeta_0$ then there is $i(j) \in I(m)$ such that $t_{i(j)-1}^m < z_j < t_{i(j)}^m$, and if $z_j \in \zeta \setminus \zeta_0$ then there is $i(j) \in I(m)$ such that $z_j = t_{i(j)}^m$. For $m \geq m_0$, let $I_1(m)$ be the set of all indices $i(j)$ with j such that $z_j \in \zeta_0$, and all pairs of indices $i(j), i(j) + 1$ with j such that $z_j \in \zeta \setminus \zeta_0$, and let $I_2(m) := I(m) \setminus I_1(m)$. By (1.19), if $z_j \in \zeta_0$ then $\lim_{m \rightarrow \infty} \Delta_{i(j)}^m f$ exists and equals either to $\Delta^- f(z_j)$ or to $\Delta^+ f(z_j)$. Otherwise if $z_j \in \zeta \setminus \zeta_0$ then $\lim_{m \rightarrow \infty} \Delta_{i(j)}^m f = \Delta^- f(z_j)$ and $\lim_{m \rightarrow \infty} \Delta_{i(j)+1}^m f = \Delta^+ f(z_j)$. Again since f is regulated and has the quadratic λ -variation, there is an $m_1 \geq m_0$ such that for each $m \geq m_1$,

$$|s_2(f; \lambda_m \cap [x, y]) - \alpha_{\lambda}(f; x, y)| < \epsilon, \quad \left| \sum_{i \in I_1(m)} (\Delta_i^m f)^2 - \sum_{\sigma} (\Delta f)^2 \right| < \epsilon \quad (3.106)$$

and

$$\left| \log \left\{ \prod_{i \in I_1(m)} (1 + \Delta_i^m f) e^{-\Delta_i^m f} \right\} - \log \left\{ \prod_{\sigma} (1 + \Delta f) e^{-\Delta f} \right\} \right| < \epsilon, \quad (3.107)$$

where $\sigma = D(x, y; \delta)$. By Taylor's theorem with remainder, for $|z| \leq 1/2$,

$$\log(1 + z) = z - z^2/2 + 3\theta z^3, \quad (3.108)$$

where $|\theta| = |\theta(z)| \leq 1$. Thus for each $m \geq m_1$, we have

$$\begin{aligned} & \log P(f; \lambda_m \cap [x, y]) \\ &= \log \prod_{i \in I_1(m)} (1 + \Delta_i^m f) + \sum_{i \in I_2(m)} \Delta_i^m f - \frac{1}{2} \sum_{i \in I_2(m)} (\Delta_i^m f)^2 + 3 \sum_{i \in I_2(m)} \theta_i (\Delta_i^m f)^3 \\ &= \log \left\{ \prod_{i \in I_1(m)} (1 + \Delta_i^m f) e^{-\Delta_i^m f} \right\} + \mu(f; x, y) \\ & \quad - \frac{1}{2} \left\{ s_2(f; \lambda_m \cap [x, y]) - \sum_{i \in I_1(m)} (\Delta_i^m f)^2 \right\} + 3 \sum_{i \in I_2(m)} \theta_i (\Delta_i^m f)^3, \end{aligned}$$

where $\theta_i := \theta(\Delta_i^m f)$. By (3.100), (3.106) and by the second inequality in (3.105), we have for each $m \geq m_1$,

$$\begin{aligned} & \left| s_2(f; \lambda_m \mathbin{\mathbb{M}} [x, y]) - \sum_{i \in I_1(m)} (\Delta_i^m f)^2 - \alpha_\lambda^c(f; x, y) \right| \\ & \leq |s_2(f; \lambda_m \mathbin{\mathbb{M}} [x, y]) - \alpha_\lambda(f; x, y)| + \left| \sum_{[x, y]} (\Delta f)^2 - \sum_{i \in I_1(m)} (\Delta_i^m f)^2 \right| < 3\epsilon. \end{aligned}$$

By (3.107) and by the first inequality in (3.105), we have for each $m \geq m_1$,

$$\left| \log \left\{ \prod_{i \in I_1(m)} (1 + \Delta_i^m f) e^{-\Delta_i^m f} \right\} - \log \gamma(f; x, y) \right| < 2\epsilon.$$

This in conjunction with (3.104) yields that the bound

$$\begin{aligned} & \left| \log P(f; \lambda_m \mathbin{\mathbb{M}} [x, y]) - \log \gamma(f; x, y) - \mu(f; x, y) + \frac{1}{2} \alpha_\lambda^c(f; x, y) \right| \\ & \leq \left| \log \left\{ \prod_{i \in I_1(m)} (1 + \Delta_i^m f) e^{-\Delta_i^m f} \right\} - \log \gamma(f; x, y) \right| \\ & \quad + \frac{1}{2} \left| s_2(f; \lambda_m \mathbin{\mathbb{M}} [x, y]) - \sum_{i \in I_1(m)} (\Delta_i^m f)^2 - \alpha_\lambda^c(f; x, y) \right| + 3 \sum_{i \in I_2(m)} |\theta_i| |\Delta_i^m f|^3 \\ & \leq 2\epsilon + (3/2)\epsilon + 3\epsilon s_2(f; \lambda_m \mathbin{\mathbb{M}} [x, y]) \end{aligned}$$

holds for all $m \geq m_1$. Since $s_2(f; \lambda_m \mathbin{\mathbb{M}} [x, y])$ is bounded in m and since ϵ is arbitrary, (3.103) holds for any $a \leq x < y \leq b$. The proof of Theorem 42 is complete. $\square$

3.5 Extended Doléans exponentials

Let f be a regulated function on $[a, b]$ having the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$. Define the forward *Doléans exponential* $\mathcal{E}_a(f) \equiv \mathcal{E}_{\lambda, a}(f)$ on $[a, b]$ by

$$\mathcal{E}_a(f; x) := \begin{cases} \exp \{f(x) - f(a) - \frac{1}{2}[f]_\lambda^c(x)\} \prod_{[a, x]} (1 + \Delta f) e^{-\Delta f} & \text{if } x \in (a, b], \\ 1 & \text{if } x = a. \end{cases} \quad (3.109)$$

Here the product is defined by (3.83) and it exists by Lemma 37. Similarly, define the backward *Doléans exponential* $\mathcal{E}_b(f) \equiv \mathcal{E}_{\lambda, b}(f)$ on $[a, b]$ by

$$\mathcal{E}_b(f; x) := \begin{cases} \exp \{f(b) - f(x) - \frac{1}{2}[f]_\lambda^c(b) + \frac{1}{2}[f]_\lambda^c(x)\} \prod_{[x, b]} (1 + \Delta f) e^{-\Delta f} & \text{if } x \in [a, b), \\ 1 & \text{if } x = b. \end{cases}$$

The forward Doléans exponential is a pathwise variant of the stochastic Doléans exponential which is the unique solution to the linear stochastic differential equation (Doléans-Dade [18]). We show in the next section that the two Doléans exponentials are unique solutions to the augmented forward and backward linear λ -integral equations, respectively. If a real valued function f is in the class $\mathcal{W}_2^*[a, b]$, so that $[f]_\lambda^c \equiv 0$, then the two Doléans exponentials agree with the values of the product integral with respect to f over the intervals $[a, x]$ and $[x, b]$,

respectively (see Theorem 36). In this section we show that the 2-vector functions $(\mathcal{E}_{\lambda,a}(f), f)$ and $(\mathcal{E}_{\lambda,b}(f), f)$ have the quadratic λ -variations.

It is easy to see that the Doléans exponentials $\mathcal{E}_a(f; x) = \beta_\lambda(f; a, x)$ and $\mathcal{E}_b(f; x) = \beta_\lambda(f; x, b)$ for $x \in [a, b]$, where $\beta_\lambda(f)$ is defined by (3.101). The two Doléans exponentials are regulated functions on $[a, b]$. Jumps of $\mathcal{E}_a(f)$ satisfy the relations:

$$\begin{aligned} (\Delta^- \mathcal{E}_a(f))(x) &= \beta_\lambda(f; a, x) - \beta_\lambda(f; a, x-) = \mathcal{E}_a(f; x-)[\beta_\lambda(f; x-, x) - 1] \\ &= \mathcal{E}_a(f; x-) \Delta^- f(x), \quad x \in (a, b], \end{aligned} \quad (3.110)$$

and

$$\begin{aligned} (\Delta^+ \mathcal{E}_a(f))(x) &= \beta_\lambda(f; a, x+) - \beta_\lambda(f; a, x) = \mathcal{E}_a(f; x)[\beta_\lambda(f; x, x+) - 1] \\ &= \mathcal{E}_a(f; x) \Delta^+ f(x), \quad x \in [a, b). \end{aligned} \quad (3.111)$$

While jumps of $\mathcal{E}_b(f)$ satisfy the relations:

$$\begin{aligned} (\Delta^- \mathcal{E}_b(f))(x) &= \beta_\lambda(f; x, b) - \beta_\lambda(f; x-, b) = \mathcal{E}_b(f; x)[1 - \beta_\lambda(f; x-, x)] \\ &= -\mathcal{E}_b(f; x) \Delta^- f(x), \quad x \in (a, b], \end{aligned} \quad (3.112)$$

and

$$\begin{aligned} (\Delta^+ \mathcal{E}_b(f))(x) &= \beta_\lambda(f; x+, b) - \beta_\lambda(f; x, b) = \mathcal{E}_b(f; x+)[1 - \beta_\lambda(f; x, x+)] \\ &= -\mathcal{E}_b(f; x+) \Delta^+ f(x), \quad x \in [a, b). \end{aligned} \quad (3.113)$$

Next we show that the distribution functions $R_{\gamma(f)}$ and $L_{\gamma(f)}$ of the function $\gamma(f)$ restricted to $[a, b]$ (see (2.4)) are pure jump functions of bounded variation. Therefore let

$$V_a(x) := \gamma(f; a, x) \text{ and } V_b(x) := \gamma(f; x, b) \text{ for } x \in [a, b].$$

By Theorem 2.4, if f is a regulated function such that $\sigma_2(f) < \infty$ and (3.84) holds, then $V_a(x-) = \gamma(f; a, x-)$, $V_b(x-) = \gamma(f; x-, b)$ for $x \in (a, b]$ and $V_a(x+) = \gamma(f; a, x+)$, $V_b(x+) = \gamma(f; x+, b)$ for $x \in [a, b)$. The multiplicativity of $\gamma(f)$ yields the following relations: for $x \in (a, b]$,

$$V_a(x) = \gamma(f; a, x) = \gamma(f; a, x-) \gamma(f; x-, x) = V_a(x-) [1 + \Delta^- f(x)] e^{-\Delta^- f(x)}, \quad (3.114)$$

$$V_b(x-) = \gamma(f; x-, b) = \gamma(f; x-, x) \gamma(f; x, b) = V_b(x) [1 + \Delta^- f(x)] e^{-\Delta^- f(x)}, \quad (3.115)$$

and for $x \in [a, b)$,

$$V_a(x+) = \gamma(f; a, x+) = \gamma(f; a, x) \gamma(f; x, x+) = V_a(x) [1 + \Delta^+ f(x)] e^{-\Delta^+ f(x)}, \quad (3.116)$$

$$V_b(x) = \gamma(f; x, b) = \gamma(f; x, x+) \gamma(f; x+, b) = V_b(x+) [1 + \Delta^+ f(x)] e^{-\Delta^+ f(x)}.$$

Lemma 43 *The functions V_a and V_b are pure jump functions on $[a, b]$ of bounded variation.*

Proof. We prove that $V := V_a$ is a pure jump function of bounded variation. The proof of the same property for V_b is similar, and hence is omitted. If $\Delta^-f(x) = -1$ for some $x \in (a, b]$ then $V(y) = 0$ for all $y \geq x$, and if $\Delta^+f(x) = -1$ for some $x \in [a, b)$ then $V(y) = 0$ for all $y > x$. In either case the claim of the lemma holds for V if it holds for its restriction to the interval $[a, x)$ if $\Delta^-f(x) = -1$, or for its restriction to $[a, x]$ if $\Delta^+f(x) = 1$. For notation simplicity we assume that (3.84) holds. By (3.114) and (3.87), for any $a < x \leq b$, we have

$$|\Delta^-V(x)| = |V(x-)| \left| (1 + \Delta^-f(x))e^{-\Delta^-f(x)} - 1 \right| \leq \frac{3\sqrt{e}}{2} \|\gamma(f)\|_\infty [\Delta^-f(x)]^2$$

provided $|\Delta^-f(x)| \leq 1/2$. The same bound holds for $|\Delta^+V(y)|$, $a \leq y < b$, with Δ^-f replaced by Δ^+f , provided $|\Delta^+f(y)| \leq 1/2$. Therefore $\sigma_1(V; (a, b)) < \infty$, and hence there exists the limit

$$\sum_{(a,x)} \{\Delta^-V + \Delta^+V\} = \lim_{\kappa, \mathfrak{P}} \sum_{\kappa \cap (a,x)} \{\Delta^-V + \Delta^+V\}$$

for each $a < x \leq b$. Let $a < x \leq b$. For a partition $a = x_0 < x_1 < \dots < x_n = x$ of $[a, x]$, by a telescoping sum, we have

$$V(x) - 1 = \Delta^+V(a) + \sum_{i=1}^n \{V(x_i-) - V(x_{i-1}+)\} + \sum_{i=1}^{n-1} \{\Delta^-V(x_i) + \Delta^+V(x_i)\} + \Delta^-V(x).$$

By multiplicativity of $\gamma(f)$ and by (3.86), for each $i = 1, \dots, n$, we get

$$\begin{aligned} |V(x_i-) - V(x_{i-1}+)| &= |V(x_{i-1}+)| \left| \prod_{(x_{i-1}+, x_i-)} (1 + \Delta f) e^{-\Delta f} - 1 \right| \\ &\leq 2e^{2\sigma_2(f)} \|\gamma(f)\|_\infty \sigma_2(f; (x_{i-1}, x_i)) \end{aligned}$$

provided (x_{i-1}, x_i) does not contain discontinuities of f with jumps bigger than $1/2$. Therefore given $\epsilon > 0$, one can choose a partition of $[a, x]$ such that for each its refinement $\{x_i: i = 0, \dots, n\}$,

$$\left| \sum_{(a,x)} \{\Delta^-V + \Delta^+V\} - \sum_{i=1}^n \{\Delta^-V(x_i) + \Delta^+V(x_i)\} \right| < \epsilon \quad \text{and} \quad \left| \sum_{i=1}^n \{V(x_i-) - V(x_{i-1}+)\} \right| < \epsilon.$$

Since $\epsilon > 0$ is arbitrary, the relation

$$V(x) = 1 + \Delta^+V(a) + \sum_{(a,x)} \{\Delta^-V + \Delta^+V\} + \Delta^-V(x)$$

holds for each $a < x \leq b$. Also, for a partition $\kappa = \{x_i: i = 0, \dots, n\}$ of $[a, b]$, we have

$$s_1(V; \kappa) \leq \sum_{i=1}^n \left\{ |\Delta^+V(x_{i-1})| + \sum_{(x_{i-1}, x_i)} \{|\Delta^-V| + |\Delta^+V|\} + |\Delta^-V(x_i)| \right\} \leq \sigma_1(V; [a, b]) < \infty.$$

Since κ is arbitrary, the bound implies that $V = V_a$ is of bounded variation. The proof of Lemma 43 is now complete. $\square$

Theorem 44 *Let f be a regulated function on $[a, b]$ having the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$ such that right discontinuity points of f are accessible by λ . Then the Doléans exponential $\mathcal{E}_a(f)$ has the quadratic λ -variation on $[a, b]$, and its bracket function is given by*

$$[\mathcal{E}_a(f)]_\lambda(x) = (LY) \int_a^x \mathcal{E}_a(f)^2 d[f]_\lambda, \quad a \leq x \leq b. \quad (3.117)$$

Also, the 2-vector function $(\mathcal{E}_a(f), f)$ has the quadratic λ -covariation on $[a, b]$, and its bracket function is given by

$$[\mathcal{E}_a(f), f]_\lambda(x) = (LY) \int_a^x \mathcal{E}_a(f) d[f]_\lambda, \quad a \leq x \leq b. \quad (3.118)$$

Proof. Since the Doléans exponential $\mathcal{E}_a(f)$ is a regulated function on $[a, b]$ and the bracket function $[f]_\lambda$ is nondecreasing, the two Left Young integrals in (3.117) and (3.118) exist by Lemma 9. Also by Proposition 2.27, the two indefinite Left Young integrals are regulated functions with jumps

$$\{\mathcal{E}_a(f)_- \Delta^- f\}^2, \{\mathcal{E}_a(f) \Delta^+ f\}^2 \quad \text{and} \quad \mathcal{E}_a(f)_- \{\Delta^- f\}^2, \mathcal{E}_a(f) \{\Delta^+ f\}^2,$$

respectively. Hence the jumps of both sides of (3.117) and (3.118) agree by (3.110) and (3.111). By Propositions 4 and 13, it is enough to prove the relations

$$\lim_{m \rightarrow \infty} C(\mathcal{E}_a(f), f; \lambda_m \mathbin{\mathbb{M}} [a, x]) = (LY) \int_a^x \mathcal{E}_a(f) d[f]_\lambda \quad (3.119)$$

and

$$\lim_{m \rightarrow \infty} s_2(\mathcal{E}_a(f); \lambda_m \mathbin{\mathbb{M}} [a, x]) = (LY) \int_a^x \mathcal{E}_a(f)^2 d[f]_\lambda \quad (3.120)$$

for $a \leq x \leq b$. For notation simplicity, we prove only that (3.119) and (3.120) hold for $x = b$. The proof for $x < b$ is the same.

Let $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$, $m \geq 1$, be partitions of $[a, b]$ constituting the sequence $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ such that f has the quadratic λ -variation and the right discontinuity points of f are accessible by λ . For a real valued function ρ on $[a, b]$, let $\Delta_i^m \rho := \rho(x_i^m) - \rho(x_{i-1}^m)$ for $i = 1, \dots, n(m)$ and $m = 1, 2, \dots$. Let $\phi(u, v) := e^u v$, $u, v \in \mathbb{R}$, and let $\mathbf{f} := (\bar{f}, V_a)$, where $\bar{f} = f - f(a) - (1/2)[f]_\lambda^c$ and $V_a = \gamma(f; a, \cdot)$ is defined by (3.83). Then $\mathcal{E}_a(f) = \phi \circ \mathbf{f}$. For the 2-vector function $\mathbf{f}$, let $\Delta_i^m \mathbf{f} := (\Delta_i^m \bar{f}, \Delta_i^m V_a)$. By the mean value theorem in the Lagrange form, for each $m \geq 1$ and $i = 1, \dots, n(m)$, there is a $\theta_i^m \in (0, 1)$ such that

$$\Delta_i^m(\mathcal{E}_a(f)) = \Delta_i^m(\phi \circ \mathbf{f}) = \langle \nabla \phi(\mathbf{y}_i^m), \Delta_i^m \mathbf{f} \rangle \quad (3.121)$$

where $\nabla \phi(u, v) = (\frac{\partial \phi}{\partial u}, \frac{\partial \phi}{\partial v})(u, v) = (e^u v, e^u)$, $u, v \in \mathbb{R}$, and $\mathbf{y}_i^m := \mathbf{f}(x_{i-1}^m) + \theta_i^m \Delta_i^m \mathbf{f}$. Given a partition $\mu = \{z_j: j = 0, \dots, m\}$ of $[a, b]$ let $J := \{j = 1, \dots, m-1: z_j \in \cup_m \lambda_m\}$. There exists the minimal integer $m_0 \geq 1$ such that $\{z_j: j \in J\} \subset \lambda_{m_0}$ and $(z_{j-1}, z_j) \cap \lambda_{m_0} \neq \emptyset$ for $j = 1, \dots, m$. For each $m \geq m_0$, let $i(j)$ be the index in $\{1, \dots, n(m)\}$ such that $x_{i(j)-1}^m < z_j \leq x_{i(j)}^m$, $I_1(m) := \{1, i(j): j = 1, \dots, m\} \cup \{i(j) + 1: j \in J\}$ and let $I_2(m) := \{1, \dots, n(m)\} \setminus I_1(m)$.

First consider relation (3.119). For each $m \geq m_0$, we then have

$$C(\mathcal{E}_a(f), f; \lambda_m) = \sum_{i \in I_1(m)} \Delta_i^m \mathcal{E}_a(f) \Delta_i^m f + \sum_{i \in I_2(m)} \Delta_i^m(\phi \circ \mathbf{f}) \Delta_i^m f$$

$$= \sum_{i=1}^{n(m)} \mathcal{E}_a(f)(x_{i-1}^m) \{\Delta_i^m f\}^2 + \sum_{i \in I_1(m)} \left[\Delta_i^m \mathcal{E}_a(f) \Delta_i^m f - \mathcal{E}_a(f)(x_{i-1}^m) \{\Delta_i^m f\}^2 \right] + R_m^{(1)},$$

where

$$R_m^{(1)} := \sum_{i \in I_2(m)} \left\{ \left[\frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) - \frac{\partial \phi}{\partial u}(\mathbf{f}(x_{i-1}^m)) \right] \{\Delta_i^m f\}^2 + \left[\frac{\partial \phi}{\partial v}(\mathbf{y}_i^m) \Delta_i^m V_a - \frac{1}{2} \frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \Delta_i^m [f]_\lambda^c \right] \Delta_i^m f \right\}.$$

Suppose that given $\epsilon > 0$ one can find a partition μ such that $|R_m^{(1)}| < C\epsilon$ for all sufficiently large m . Since $\Delta^+ f(x) = 0$ if $x \notin \cup_m \lambda_m$, the limit

$$\begin{aligned} & \lim_{m \rightarrow \infty} \sum_{i \in I_1(m)} \left[\Delta_i^m \mathcal{E}_a(f) \Delta_i^m f - \mathcal{E}_a(f)(x_{i-1}^m) \{\Delta_i^m f\}^2 \right] \\ &= \sum_{\mu \setminus a} \left[\Delta^- \mathcal{E}_a(f) \Delta^- f - \mathcal{E}_a(f)_- \{\Delta^- f\}^2 \right] + \sum_{\mu \setminus b} \left[\Delta^+ \mathcal{E}_a(f) \Delta^+ f - \mathcal{E}_a(f) \{\Delta^+ f\}^2 \right] = 0 \end{aligned}$$

for any given μ by (3.110) and (3.111). An application of Lemma 11 then yields

$$\lim_{m \rightarrow \infty} C(\mathcal{E}_a(f), f; \lambda_m) = \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} \mathcal{E}_a(f)(x_{i-1}^m) \{\Delta_i^m f\}^2 = (LY) \int_a^b \mathcal{E}_a(f) d[f]_\lambda. \quad (3.122)$$

To choose a partition μ , let $\epsilon \in (0, 1)$ be given. Since $\frac{\partial \phi}{\partial u} = \phi$ is uniformly continuous on the range of $\mathbf{f}$, there exists $\delta > 0$ such that $|\phi(u_1, v_1) - \phi(u_2, v_2)| < \epsilon$ if $|u_1| \vee |u_2| \leq \|\mathbf{f}\|_\infty$, $|v_1| \vee |v_2| \leq \|V_a\|_\infty$ and $|u_1 - u_2| \vee |v_1 - v_2| \leq \delta$. Since f and V_a are regulated, there exists a partition $\mu = \{z_j: j = 0, \dots, m\}$ of $[a, b]$ such that for each $j = 1, \dots, m$,

$$\text{Osc}(\bar{f}; (z_{j-1}, z_j)) < \delta, \quad \text{Osc}(V_a; (z_{j-1}, z_j)) < \delta,$$

$$\text{Osc}(f; (z_{j-1}, z_j)) < \epsilon \quad \text{and} \quad \text{Osc}([f]_\lambda^c; (z_{j-1}, z_j)) < \epsilon.$$

Since $[\Delta_i^m] := [x_{i-1}^m, x_i^m] \subset \cup_j (z_{j-1}, z_j)$ for $i \in I_2(m)$, we then have the bound

$$\begin{aligned} |R_m^{(1)}| &\leq \max_{i \in I_2(m)} \text{Osc}(\phi \circ \mathbf{f}; [\Delta_i^m]) s_2(f; \lambda_m) + \left\| \frac{\partial \phi}{\partial v} \circ \mathbf{f} \right\|_\infty \max_{i \in I_2(m)} \text{Osc}(f; [\Delta_i^m]) s_1(V_a; \lambda_m) \\ &\quad + \frac{1}{2} \|\phi \circ \mathbf{f}\|_\infty \left(\max_{i \in I_2(m)} \text{Osc}([f]_\lambda^c; [\Delta_i^m]) s_1([f]_\lambda^c; \lambda_m) s_2(f; \lambda_m) \right)^{1/2} \leq C\epsilon \end{aligned}$$

for all sufficiently large m . Therefore (3.122) holds.

Now consider relation (3.120). Again using the mean value theorem (3.121) and the notation following it, for each $m \geq m_0$, we have

$$\begin{aligned} s_2(\mathcal{E}_a(f); \lambda_m) &= \sum_{i \in I_1(m)} \{\Delta_i^m \mathcal{E}_a(f)\}^2 + \sum_{i \in I_2(m)} \{\Delta_i^m (\phi \circ \mathbf{f})\}^2 \\ &= \sum_{i=1}^{n(m)} \{\mathcal{E}_a(f)(x_{i-1}^m) \Delta_i^m f\}^2 + \sum_{i \in I_1(m)} \left[\{\Delta_i^m \mathcal{E}_a(f)\}^2 - \{\mathcal{E}_a(f)(x_{i-1}^m) \Delta_i^m f\}^2 \right] + R_m^{(2)}, \end{aligned}$$

where as before letting $\mathbf{y}_i^m := \mathbf{f}(x_{i-1}^m) + \theta_i^m \Delta_i^m \mathbf{f}$,

$$R_m^{(2)} := \sum_{i \in I_2(m)} \left\{ \left[\left(\frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \right)^2 - \left(\frac{\partial \phi}{\partial u}(\mathbf{f}(x_{i-1}^m)) \right)^2 \right] \{\Delta_i^m f\}^2 \right. \\ \left. + 2 \frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \Delta_i^m f \left[\frac{\partial \phi}{\partial v}(\mathbf{y}_i^m) \Delta_i^m V_a - \frac{1}{2} \frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \Delta_i^m [f]^c \right] + \left[\frac{\partial \phi}{\partial v}(\mathbf{y}_i^m) \Delta_i^m V_a - \frac{1}{2} \frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \Delta_i^m [f]^c \right]^2 \right\}.$$

As in the preceding case, given $\epsilon > 0$ there exists a set μ such that $|R_m^{(2)}| < C\epsilon$ for all sufficiently large m . Again since $\Delta^+ f(x) = 0$ if $x \notin \cup_m \lambda_m$, the limit

$$\lim_{m \rightarrow \infty} \sum_{i \in I_1(m)} \left[\{\Delta_i^m \mathcal{E}_a(f)\}^2 - \{\mathcal{E}_a(f)(x_{i-1}^m) \Delta_i^m f\}^2 \right] \\ \sum_{\mu \setminus a} \left[\{\Delta^- \mathcal{E}_a(f)\}^2 - \{\mathcal{E}_a(f) \Delta^- f\}^2 \right] + \sum_{\mu \setminus b} \left[\{\Delta^+ \mathcal{E}_a(f)\}^2 - \{\mathcal{E}_a(f) \Delta^+ f\}^2 \right] = 0$$

by (3.110) and (3.111). An application of Lemma 11 then yields

$$\lim_{m \rightarrow \infty} s_2(\mathcal{E}_a(f); \lambda_m) = \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} \{\mathcal{E}_a(f)(x_{i-1}^m) \Delta_i^m f\}^2 = (LY) \int_a^b \mathcal{E}_a(f)^2 d[f]_\lambda.$$

The proof of Theorem 44 is complete. $\square$

Next we prove analogous theorem with the forward Doléans exponential $\mathcal{E}_a(f)$ replaced by the backward Doléans exponential $\mathcal{E}_b(f)$.

Theorem 45 *Let f be a regulated function on $[a, b]$ having the quadratic λ -variation for some $\lambda \in \Lambda[a, b]$ such that left discontinuity points of f are accessible by λ . Then the Doléans exponential $\mathcal{E}_b(f)$ has the quadratic λ -variation on $[a, b]$, and its bracket function is given by*

$$[\mathcal{E}_b(f)]_\lambda(x) = (RY) \int_a^x \mathcal{E}_b(f)^2 d[f]_\lambda, \quad a \leq x \leq b. \quad (3.123)$$

Also, the 2-vector function $(\mathcal{E}_b(f), f)$ has the quadratic λ -covariation on $[a, b]$, and its bracket function is given by

$$[\mathcal{E}_b(f), f]_\lambda(x) = -(RY) \int_a^x \mathcal{E}_b(f) d[f]_\lambda, \quad a \leq x \leq b. \quad (3.124)$$

Proof. Since the Doléans exponential $\mathcal{E}_b(f)$ is a regulated function on $[a, b]$ and the bracket function $[f]_\lambda$ is nondecreasing, the two Right Young integrals in (3.123) and (3.124) are defined by Lemma 9. Also by Proposition 2.27, the two indefinite Right Young integrals are regulated functions with jumps

$$\{\mathcal{E}_b(f) \Delta^- f\}^2, \{\mathcal{E}_b(f)_+ \Delta^+ f\}^2 \quad \text{and} \quad \mathcal{E}_b(f) \{\Delta^- f\}^2, \mathcal{E}_b(f)_+ \{\Delta^+ f\}^2,$$

respectively. Hence the jumps of both sides of (3.123) and (3.124) agree by (3.112) and (3.113). By Propositions 4 and 13, it is enough to prove the relations

$$\lim_{m \rightarrow \infty} C(\mathcal{E}_b(f), f; \lambda_m \cap [a, x]) = -(RY) \int_a^x \mathcal{E}_b(f) d[f]_\lambda \quad (3.125)$$

and

$$\lim_{m \rightarrow \infty} s_2(\mathcal{E}_b(f); \lambda_m \cap [a, x]) = (RY) \int_a^x \mathcal{E}_b(f)^2 d[f]_\lambda \quad (3.126)$$

for $a \leq x \leq b$. For notation simplicity, we prove only that (3.125) and (3.126) hold for $x = b$.

Let $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$, $m \geq 1$, be partitions of $[a, b]$ constituting the sequence $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ such that f has the quadratic λ -variation and the left discontinuity points of f are accessible by λ . For a real valued function ρ on $[a, b]$, let $\Delta_i^m \rho := \rho(x_i^m) - \rho(x_{i-1}^m)$ for $i = 1, \dots, n(m)$ and $m = 1, 2, \dots$. Let $\phi(u, v) := e^{-u}v$, $u, v \in \mathbb{R}$, and let $\mathbf{f} := (\bar{f}, V_b)$, where $\bar{f} = f - f(b) + (1/2)[f]_\lambda^c(b) - (1/2)[f]_\lambda^c$ and $V_b = \gamma(f; \cdot, b)$ is defined by (3.83). Then $\mathcal{E}_b(f) = \phi \circ \mathbf{f}$. For the 2-vector function $\mathbf{f}$, let $\Delta_i^m \mathbf{f} := (\Delta_i^m \bar{f}, \Delta_i^m V_b)$. By the mean value theorem in the Lagrange form, for each $m \geq 1$ and $i = 1, \dots, n(m)$, there is a $\theta_i^m \in (0, 1)$ such that

$$\Delta_i^m(\mathcal{E}_b(f)) = \Delta_i^m(\phi \circ \mathbf{f}) = \langle \nabla \phi(\mathbf{y}_i^m), \Delta_i^m \mathbf{f} \rangle \quad (3.127)$$

where $\nabla \phi(u, v) = (\frac{\partial \phi}{\partial u}, \frac{\partial \phi}{\partial v})(u, v) = (-e^{-u}v, e^{-u})$, $u, v \in \mathbb{R}$, and $\mathbf{y}_i^m := \mathbf{f}(x_{i-1}^m) + \theta_i^m \Delta_i^m \mathbf{f}$. Given a partition $\mu = \{z_j: j = 0, \dots, m\}$ of $[a, b]$ let $J := \{j = 1, \dots, m-1: z_j \in \cup_m \lambda_m\}$. There exists the minimal integer $m_0 \geq 1$ such that $\{z_j: j \in J\} \subset \lambda_{m_0}$ and $(z_{j-1}, z_j) \cap \lambda_{m_0} \neq \emptyset$ for $j = 1, \dots, m$. For each $m \geq m_0$, let $i(j)$ be the index in $\{1, \dots, n(m)\}$ such that $x_{i(j)-1}^m < z_j \leq x_{i(j)}^m$, $I_1(m) := \{1, i(j): j = 1, \dots, m\} \cup \{i(j) + 1: j \in J\}$ and let $I_2(m) := \{1, \dots, n(m)\} \setminus I_1(m)$.

First consider relation (3.125). For each $m \geq m_0$, we then have

$$\begin{aligned} C(\mathcal{E}_b(f), f; \lambda_m) &= \sum_{i \in I_1(m)} \Delta_i^m \mathcal{E}_b(f) \Delta_i^m f + \sum_{i \in I_2(m)} \Delta_i^m(\phi \circ \mathbf{f}) \Delta_i^m f \\ &= - \sum_{i=1}^{n(m)} \mathcal{E}_b(f)(x_i^m) \{\Delta_i^m f\}^2 + \sum_{i \in I_1(m)} \left[\Delta_i^m \mathcal{E}_b(f) \Delta_i^m f + \mathcal{E}_b(f)(x_i^m) \{\Delta_i^m f\}^2 \right] + R_m^{(1)}, \end{aligned}$$

where

$$R_m^{(1)} := \sum_{i \in I_2(m)} \left\{ \left[\frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) - \frac{\partial \phi}{\partial u}(\mathbf{f}(x_{i-1}^m)) \right] \{\Delta_i^m f\}^2 + \left[\frac{\partial \phi}{\partial v}(\mathbf{y}_i^m) \Delta_i^m V_b - \frac{1}{2} \frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \Delta_i^m [f]_\lambda^c \right] \Delta_i^m f \right\}.$$

As in the proof of Theorem 44, given $\epsilon > 0$ one can find a partition μ such that $|R_m^{(1)}| < C\epsilon$ for all sufficiently large m . Since $\Delta^- f(x) = 0$ if $x \notin \cup_m \lambda_m$, the limit

$$\begin{aligned} &\lim_{m \rightarrow \infty} \sum_{i \in I_1(m)} \left[\Delta_i^m \mathcal{E}_b(f) \Delta_i^m f + \mathcal{E}_b(f)(x_i^m) \{\Delta_i^m f\}^2 \right] \\ &= \sum_{\mu \setminus a} \left[\Delta^- \mathcal{E}_b(f) \Delta^- f + \mathcal{E}_b(f) \{\Delta^- f\}^2 \right] + \sum_{\mu \setminus b} \left[\Delta^+ \mathcal{E}_b(f) \Delta^+ f + \mathcal{E}_b(f) \{\Delta^+ f\}^2 \right] = 0 \end{aligned}$$

for any given μ by (3.112) and (3.113). An application of Lemma 11 then yields

$$\lim_{m \rightarrow \infty} C(\mathcal{E}_b(f), f; \lambda_m) = - \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} \mathcal{E}_b(f)(x_i^m) \{\Delta_i^m f\}^2 = -(RY) \int_a^b \mathcal{E}_b(f) d[f]_\lambda.$$

That is (3.125) holds with $x = b$.

Now consider relation (3.126). Again using the mean value theorem (3.127) and the notation following it, for each $m \geq m_0$, we have

$$\begin{aligned} s_2(\mathcal{E}_b(f); \lambda_m) &= \sum_{i \in I_1(m)} \{\Delta_i^m \mathcal{E}_b(f)\}^2 + \sum_{i \in I_2(m)} \{\Delta_i^m(\phi \circ \mathbf{f})\}^2 \\ &= \sum_{i=1}^{n(m)} \{\mathcal{E}_b(f)(x_i^m) \Delta_i^m f\}^2 + \sum_{i \in I_1(m)} \left[\{\Delta_i^m \mathcal{E}_b(f)\}^2 - \{\mathcal{E}_b(f)(x_i^m) \Delta_i^m f\}^2 \right] + R_m^{(2)}, \end{aligned}$$

where as before letting $\mathbf{y}_i^m := \mathbf{f}(x_{i-1}^m) + \theta_i^m \Delta_i^m \mathbf{f}$,

$$\begin{aligned} R_m^{(2)} &:= \sum_{i \in I_2(m)} \left\{ \left[\left(\frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \right)^2 - \left(\frac{\partial \phi}{\partial u}(\mathbf{f}(x_{i-1}^m)) \right)^2 \right] \{\Delta_i^m f\}^2 \right. \\ &\quad \left. + 2 \frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \Delta_i^m f \left[\frac{\partial \phi}{\partial v}(\mathbf{y}_i^m) \Delta_i^m V_b - \frac{1}{2} \frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \Delta_i^m [f]^c \right] + \left[\frac{\partial \phi}{\partial v}(\mathbf{y}_i^m) \Delta_i^m V_b - \frac{1}{2} \frac{\partial \phi}{\partial u}(\mathbf{y}_i^m) \Delta_i^m [f]^c \right]^2 \right\}. \end{aligned}$$

As in the preceding theorem, given $\epsilon > 0$ there exists a partition μ such that $|R_m^{(2)}| < C\epsilon$ for all sufficiently large m . Again since $\Delta^- f(x) = 0$ if $x \notin \cup_m \lambda_m$, the limit

$$\begin{aligned} &\lim_{m \rightarrow \infty} \sum_{i \in I_1(m)} \left[\{\Delta_i^m \mathcal{E}_b(f)\}^2 - \{\mathcal{E}_b(f)(x_i^m) \Delta_i^m f\}^2 \right] \\ &\sum_{\mu \setminus a} \left[\{\Delta^- \mathcal{E}_b(f)\}^2 - \{\mathcal{E}_b(f) \Delta^- f\}^2 \right] + \sum_{\mu \setminus b} \left[\{\Delta^+ \mathcal{E}_b(f)\}^2 - \{\mathcal{E}_b(f)_+ \Delta^+ f\}^2 \right] = 0 \end{aligned}$$

by (3.112) and (3.113). An application of Lemma 11 then yields

$$\lim_{m \rightarrow \infty} s_2(\mathcal{E}_b(f); \lambda_m) = \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} \{\mathcal{E}_b(f)(x_i^m) \Delta_i^m f\}^2 = (RY) \int_a^b \mathcal{E}_b(f)^2 d[f]_\lambda.$$

The proof of Theorem 45 is complete. $\square$

3.6 Augmented linear λ -integral equations

Let f be a regulated function on $[a, b]$ which is (λ, p) -decomposable for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, and let $D_{\lambda, p}^+(f)$ be nonempty. We consider the forward linear Left λ -integral equation

$$F(y) = 1 + (L) \int_a^y F d_\lambda f, \quad a \leq y \leq b, \quad (3.128)$$

provided the Left λ -integral is defined on $[a, b]$ with respect to $D_{\lambda, p}^+(f)$. It is not hard to check using a chain rule that the forward Doléans exponential $\mathcal{E}_a(f)$ satisfy (3.128). More challenging is the task to find a large enough class of functions F which contains the Doléans exponential $\mathcal{E}_a(f)$ and contains no other solutions of (3.128). For the meantime we show that an augmented forward linear Left λ -integral equation has the Doléans exponential as the unique solution in the class of all functions in $\text{dual}(\mathcal{W}_p)$ which have the quadratic λ -variation.

Definition 46 For $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, let f be a (λ, p) -decomposable function on $[a, b]$, and let $D_{\lambda, p}^+(f)$ be nonempty. We say that a function $F \in \text{dual}(\mathcal{W}_p)[a, b] \cap Q_\lambda[a, b]$ is a *solution of the augmented forward linear Left λ -integral equation*

$$d_\lambda^\rightarrow F = F d_\lambda^\rightarrow f \quad \text{on } [a, b] \quad (3.129)$$

if (a) and (b) hold, where

- (a) for $(g, h) \in D_{\lambda, p}^+(f)$ and $\psi \in \text{dual}(\mathcal{W}_p)[a, b]$, $(LC) \int \langle (\psi, -\psi F), d_\lambda(F, g) \rangle$ is defined on $[a, b]$, and for $a \leq y \leq b$,

$$(LC) \int_a^y \langle (\psi, -\psi F), d_\lambda(F, g) \rangle = (LY) \int_a^y \psi F dh; \quad (3.130)$$

- (b) the 2-vector function (F, f) has the quadratic λ -variation and for $a \leq y \leq b$,

$$\begin{pmatrix} [F]_\lambda & [f, F]_\lambda \\ [F, f]_\lambda & [f]_\lambda \end{pmatrix} (y) = (LY) \int_a^y \begin{pmatrix} F^2 & F \\ F & 1 \end{pmatrix} d[f]_\lambda. \quad (3.131)$$

Let f be a (λ, p) -decomposable function on $[a, b]$ for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, and let $D_{\lambda, p}^+(f)$ be nonempty. Suppose that $F \in \text{dual}(\mathcal{W}_p)$ has the quadratic λ -variation and is a solution of the augmented forward linear Left λ -integral equation (3.129). Then in (a) of Definition 46 taking $\psi \equiv 1$, by Proposition 23, it follows that the Left λ -integral $(L) \int F d_\lambda f$ is defined on $[a, b]$ with respect to $D_{\lambda, p}^+(f)$, and (3.128) holds. The more general relation (3.130) contains a form of the substitution rule for the Left λ -integral, analogous to the associativity property of the stochastic integral. Indeed, assuming the following integrals exist and using (3.130), (3.128), we have

$$\begin{aligned} & (LC) \int_a^b \psi d_\lambda F - (LC) \int_a^b \psi F d_\lambda g \\ &= (LC) \int_a^b \langle (\psi, -\psi F), d_\lambda(F, g) \rangle = (LY) \int_a^b \psi F dh \quad \text{by Theorem 2.29} \\ &= (LY) \int_a^b \psi d \left((LY) \int_a^\cdot F dh \right) = (LC) \int_a^b \psi d_\lambda F - (LC) \int_a^b \psi d_\lambda \left((LC) \int_a^\cdot F d_\lambda g \right). \end{aligned}$$

Comparing the left and right sides of these relations we get the substitution rule for the Left λ -integral:

$$(LC) \int_a^b \psi F d_\lambda g = (LC) \int_a^b \psi d_\lambda \left((LC) \int_a^\cdot F d_\lambda g \right).$$

The linear Left Young integral with respect to a function having bounded p -variation for some $1 \leq p < 2$ was solved by Dudley and Norvaiša [23, Theorem II.5.21]:

Theorem 47 Let $f \in \mathcal{W}_p[a, b]$ for some $1 \leq p < 2$. In the class $\text{dual}(\mathcal{W}_p)[a, b]$ the Doléans exponential $\mathcal{E}_a(f)$ is the unique solution of the forward linear Left Young integral equation

$$F(y) = 1 + (LY) \int_a^y F df, \quad a \leq y \leq b. \quad (3.132)$$

Dudley and Norvaiša considered equation (3.132) with respect to a function f having values in a Banach algebra, and the unique solution of their equation is the indefinite product integral with respect to the function f . In the case f has real values, the indefinite product integral with respect to f has values which agree with the Doléans exponential $\mathcal{E}_a(f)$ by Theorem II.4.4 of Dudley and Norvaiša [23]. The only difference with the cited result of Dudley and Norvaiša is that the uniqueness class in Theorem 47 is larger in the case $p = 1$, which is the class of all regulated functions.

Now we are ready to solve the augmented forward linear Left λ -integral equation (3.129).

Theorem 48 *Let $\lambda \in \Lambda[a, b]$, $1 \leq p < 2$ and let f be a (λ, p) -decomposable function on $[a, b]$. If $D_{\lambda, p}^+(f) \neq \emptyset$ then in the class $\text{dual}(\mathcal{W}_p)[a, b] \cap Q_\lambda[a, b]$ the Doléans exponential $\mathcal{E}_a(f)$ is the unique solution of the augmented forward linear Left λ -integral equation (3.129).*

Proof. Let $(g, h) \in D_{\lambda, p}^+(f)$. By Corollary 16, f has the quadratic λ -variation, and therefore the Doléans exponential $\mathcal{E}_a(f)$ is defined. Next we show that $\mathcal{E}_a(f)$ is a solution of the augmented forward linear Left λ -integral equation (3.129). By Theorem 44, the 2-vector function $(\mathcal{E}_a(f), f)$ has the quadratic λ -variation and (3.131) holds with $F = \mathcal{E}_a(f)$. Let $\bar{f} := f - f(a) - (1/2)[g]_\lambda^c$, and $V_a := \gamma(f; a, \cdot)$, where $\gamma(f)$ is defined by (3.83). Then $\mathcal{E}_a(f) = e^{\bar{f}} V_a$ and V_a is pure jump function of bounded variation by Lemma 43. Since the composition of the function $\bar{f}$ with a smooth function does not change its p -variation, and since $\mathcal{W}_p$ is a Banach algebra, $\mathcal{E}_a(f) \in \text{dual}(\mathcal{W}_p)[a, b]$, and so $\mathcal{E}_a(f)$ satisfies condition (b) of Definition 46. To show that $\mathcal{E}_a(f)$ satisfies condition (a), we apply the chain rule formula of Theorem 34.(a) to the composition $\phi \circ \mathbf{f} = \mathcal{E}_a(f)$, where the 2-vector function $\mathbf{f}$ is defined by

$$\mathbf{f} := (\bar{f}, V_a) = (C - f(a), 0) + (g, 0) + (h - \frac{1}{2}[g]_\lambda^c, V_a) \quad \text{and} \quad \phi(u, v) = e^u v, \quad u, v \in \mathbb{R}. \quad (3.133)$$

Let $\mathbf{g} := (g, 0)$ and $\mathbf{h} := (h - (1/2)[g]_\lambda^c, V_a)$. Then $\mathbf{f}$ is (λ, p) -decomposable by $(\mathbf{g}, \mathbf{h})$. Since all discontinuity points of V_a are the same as discontinuity points of f , the right discontinuity points of $\mathbf{f}$ are accessible by λ . Let $\psi \in \text{dual}(\mathcal{W}_p)[a, b]$. By statement (a) of Theorem 34, the Left Cauchy λ -integral

$$(LC) \int \langle (\psi, -\psi(\nabla \phi \circ \mathbf{f})), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle = (LC) \int \langle (\psi, -\psi \mathcal{E}_a(f)), d_\lambda(\mathcal{E}_a(f), g) \rangle$$

is defined on $[a, b]$, where $\nabla \phi(u, v) = (\frac{\partial \phi}{\partial u}, \frac{\partial \phi}{\partial v})(u, v) = (e^u v, e^u)$ for $u, v \in \mathbb{R}$, and the equalities

$$\begin{aligned} & (LC) \int_a^y \langle (\psi, -\psi \mathcal{E}_a(f)), d_\lambda(\mathcal{E}_a(f), g) \rangle \\ &= (LY) \int_a^y \psi \mathcal{E}_a(f) d(h - (1/2)[g]_\lambda^c) + (LY) \int_a^y \psi e^{\bar{f}} dV_a + \frac{1}{2}(RS) \int_a^y \psi \mathcal{E}_a(f) d[g]_\lambda^c \\ &+ \sum_{(a, y]} \psi_- \left\{ \Delta^- \mathcal{E}_a(f) - \mathcal{E}_a(f) - \Delta^- f - e^{\bar{f}} \Delta^- V_a \right\} + \sum_{(a, y]} \psi \left\{ \Delta^+ \mathcal{E}_a(f) - \mathcal{E}_a(f) \Delta^+ f - e^{\bar{f}} \Delta^+ V_a \right\} \\ &= (LY) \int_a^y \psi \mathcal{E}_a(f) dh + (LY) \int_a^y \psi e^{\bar{f}} dV_a - \left\{ \sum_{(a, y]} \psi_- e^{\bar{f}} \Delta^- V_a + \sum_{(a, y]} \psi e^{\bar{f}} \Delta^+ V_a \right\} \end{aligned} \quad (3.134)$$

hold for $a \leq y \leq b$. The last equality follows by Lemma 9 and relations (3.110), (3.111). Let $\lambda = \{\lambda_m: m \geq 1\}$ with $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$. Since V_a is pure jump function of bounded

variation, and since $\{x \in (a, b): \Delta^+ V_a(x) \neq 0\} \subset \cup_m \lambda_m$ by (3.116), by Theorem 2.22, we have the equalities

$$\begin{aligned} (LY) \int_a^y \psi e^{\bar{f}} dV_a &= \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} \psi(x_{i-1}^m) e^{\bar{f}(x_{i-1}^m)} [V_a(x_i^m \wedge y) - V_a(x_{i-1}^m \wedge y)] \\ &= \sum_{(a,y]} \psi_- e^{\bar{f}} \Delta^- V_a + \sum_{[a,y)} \psi e^{\bar{f}} \Delta^+ V_a \end{aligned}$$

for $a \leq y \leq b$. Thus the last two terms in (3.134) give zero to the equality, and hence the Doléans exponential $\mathcal{E}_a(f)$ solves the augmented forward linear left λ -integral equation (3.129).

Finally, suppose that $F \in \text{dual}(\mathcal{W}_p)$, has the quadratic λ -variation and solves the augmented forward linear Left λ -integral equation (3.129). We then show that $F = \mathcal{E}_a(f)$. To this aim let $V := e^{-\bar{f}} F$, where $\bar{f} := f - f(a) - (1/2)[g]_\lambda^c$. It is enough to prove that V is equal to the pure jump function $V_a = \gamma(f; a, \cdot)$. Let $\mathbf{C} := (C - f(a), 0)$, $\mathbf{g} := (g, F)$, $\mathbf{h} := (h - (1/2)[g]_\lambda^c, 0)$,

$$\mathbf{f} := (\bar{f}, F) = \mathbf{C} + \mathbf{g} + \mathbf{h} \quad \text{and} \quad \phi(u, v) := e^{-u} v \quad \text{for } u, v \in \mathbb{R}. \quad (3.135)$$

Then $V = \phi \circ \mathbf{f}$. We wish to apply a chain rule to show that V is the unique solution to the linear Left Young integral equation with respect to a pure jump function of bounded variation. To this aim we have to show that the 2-vector function $\mathbf{g} = (g, F)$ has the quadratic λ -variation. Recall that by condition (b) of Definition 46, the 2-vector function (f, F) has the quadratic λ -variation.

Lemma 49 *For $\lambda \in \Lambda[a, b]$ and $p \in [1, 2)$, let f be a regulated function on $[a, b]$ which is (λ, p) -decomposable by $(g, h) \in D_{\lambda, p}^+(f)$. Let $F \in \text{dual}(\mathcal{W}_p)[a, b]$ be such that*

$$\{x \in (a, b): \Delta^+ F(x) \neq 0\} \subset \{x \in (a, b): \Delta^+ f(x) \neq 0\}. \quad (3.136)$$

Then the 2-vector function (f, F) has the quadratic λ -covariation if and only if the 2-vector function (g, F) has the quadratic λ -covariation, and if the two functions have the quadratic λ -covariation then $[f, F]_\lambda^c = [g, F]_\lambda^c$.

Proof. To begin with we show that the 2-vector function (h, F) has the quadratic λ -covariation with the bracket function

$$[h, F]_\lambda(y) = \sum_{(a,y]} \Delta^- h \Delta^- F + \sum_{[a,y)} \Delta^+ h \Delta^+ F, \quad a \leq y \leq b. \quad (3.137)$$

To this aim we apply Proposition 14 with $f_1 = h \in \mathcal{W}_p$ and $f_2 = F \in \text{dual}(\mathcal{W}_p)$. If $(\Delta^- F \Delta^+ h + \Delta^- h \Delta^+ F)(x) \neq 0$ for some $x \in (a, b)$, then either $\{x \in (a, b): \Delta^+ h(x) \neq 0\}$ or $\{x \in (a, b): \Delta^+ F(x) \neq 0\}$. Since $(g, h) \in D_{\lambda, p}^+(f)$, by (3.136), (3.25) holds. Therefore by Proposition 14, the function (h, F) has the quadratic λ -covariation with the bracket function (3.137). Letting $\lambda = \{\lambda_m: m \geq 1\}$, the conclusion then follows from the relation

$$C(f, F; \lambda_m \mathbin{\frown} [u, v]) = C(g, F; \lambda_m \mathbin{\frown} [u, v]) + C(h, F; \lambda_m \mathbin{\frown} [u, v]),$$

which holds for each m and $a \leq u < v \leq b$, and since $\Delta f \Delta F = \Delta g \Delta F + \Delta h \Delta F$. The proof of Lemma 49 is complete. $\square$

To apply Lemma 49 to f, g, h and F as before we have to check its assumption (3.136). By condition (a) of Definition 46 with $\psi \equiv 1$ and by Proposition 23, the Left λ -integral $(L) \int F d_\lambda f$

is defined on $[a, b]$ with respect to $D_{\lambda, p}^+(f)$, and (3.128) holds. By Definition 17 of the Left Cauchy λ -integral in conjunction with Proposition 2.27 concerning jumps of the indefinite Left Young integral, we have that

$$\Delta^- F(y) = (F_- \Delta^- f)(y) \quad \text{and} \quad \Delta^+ F(x) = (F \Delta^+ f)(x) \quad (3.138)$$

for $a \leq y < x \leq b$. In particular, it then follows that (3.136) holds. Therefore by Lemma 49, the 2-vector function $\mathbf{g} = (g, F)$ has the quadratic λ -covariation, and thereby has the quadratic λ -variation. Since the pair $(\mathbf{g}, \mathbf{h})$ is (λ, p) -dual and (3.135) holds, $\mathbf{f}$ is (λ, p) -decomposable by $(\mathbf{g}, \mathbf{h})$. By (3.138), it follows that $(\mathbf{g}, \mathbf{h}) \in D_{\lambda, p}^+(\mathbf{f})$. Therefore all assumptions of Theorem 34.(a) are satisfied by $\mathbf{f}$ and ϕ . By the first part of statement (a) of Theorem 34 with $\psi \equiv 1$, the Left Cauchy λ -integrals

$$(LC) \int \langle (1, -\nabla \phi \circ \mathbf{f}), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle = \mu(\phi \circ \mathbf{f}) - (LC) \int \langle \nabla \phi \circ \mathbf{f}, d_\lambda \mathbf{g} \rangle$$

are defined on $[a, b]$, where $\nabla \phi(u, v) = (\frac{\partial \phi}{\partial u}, \frac{\partial \phi}{\partial v})(u, v) = (-e^{-u}v, e^{-u})$ for $u, v \in \mathbb{R}$, and μ is defined by (2.2). Further, by condition (a) of Definition 46 with $\psi = \exp\{-\bar{f}\}$, we have the equality

$$(LC) \int \langle \nabla \phi \circ \mathbf{f}, d_\lambda \mathbf{g} \rangle = (LC) \int \left\langle \left(-e^{-\bar{f}} F, e^{-\bar{f}} \right), d_\lambda(g, F) \right\rangle = (LY) \int e^{-\bar{f}} F dh. \quad (3.139)$$

By condition (b) of Definition 46, by the second part of Lemma 49, and by (3.138), we have

$$(LY) \int_a^y F d[f]_\lambda = [f, F]_\lambda(y) = [g, F]_\lambda^c(y) + \sum_{(a, y]} F_- \{\Delta^- f\}^2 + \sum_{[a, y)} F \{\Delta^+ f\}^2$$

for $a \leq y \leq b$. On the other hand, by the first part of Lemma 3 and since $[f]_\lambda^c = [g]_\lambda^c$, we have

$$(LY) \int_a^y F d[f]_\lambda = (RS) \int_a^y F d[g]_\lambda^c + \sum_{(a, y]} F_- \{\Delta^- f\}^2 + \sum_{[a, y)} F \{\Delta^+ f\}^2$$

for $a \leq y \leq b$. Therefore $[g, F]_\lambda^c(y) = (RS) \int_a^y F d[g]_\lambda^c$ for $a \leq y \leq b$. This in conjunction with the substitution theorem for the Riemann-Stieltjes integral, yields the equality

$$(RS) \int_a^y e^{-\bar{f}} d[g, F]_\lambda^c = (RS) \int_a^y e^{-\bar{f}} F d[g]_\lambda^c \quad (3.140)$$

for $a \leq y \leq b$. By the second part of Theorem 34.(a) with $\psi \equiv 1$ applied to $V = \phi \circ \mathbf{f}$, we then have

$$\begin{aligned} V(y) - V(a) &= (LC) \int_a^y \langle \nabla \phi \circ \mathbf{f}, d_\lambda \mathbf{g} \rangle - (LY) \int_a^y e^{-\bar{f}} F d(h - \frac{1}{2}[g]_\lambda^c) \\ &\quad + \frac{1}{2}(RS) \int_a^y e^{-\bar{f}} F d[g]_\lambda^c - (RS) \int_a^y e^{-\bar{f}} d[g, F]_\lambda^c \\ &\quad + \sum_{(a, y]} \left\{ \Delta^- V - \left(e^{-\bar{f}} \right)_- \Delta^- F + \left(e^{-\bar{f}} F \right)_- \Delta^- f \right\} \\ &\quad + \sum_{[a, y)} \left\{ \Delta^+ V - e^{-\bar{f}} \Delta^+ F + e^{-\bar{f}} F \Delta^+ f \right\} \end{aligned}$$

$$\begin{aligned}
\text{by Lemma 9} &= (LC) \int_a^y \langle \nabla \phi \circ \mathbf{f}, d_\lambda \mathbf{g} \rangle - (LY) \int_a^y e^{-\bar{f}} F dh \\
&+ \frac{1}{2} (RS) \int_a^y e^{-\bar{f}} F d[g]_\lambda^c + \frac{1}{2} (RS) \int_a^y e^{-\bar{f}} F d[g]_\lambda^c - (RS) \int_a^y e^{-\bar{f}} F d[g]_\lambda^c \\
&+ \sum_{(a,y]} \left\{ \Delta^- V - \left(e^{-\bar{f}} \right)_- F_- \Delta^- f + \left(e^{-\bar{f}} \right)_- F_- \Delta^- f \right\} \\
&+ \sum_{[a,y)} \left\{ \Delta^+ V - e^{-\bar{f}} F \Delta^+ f + e^{-\bar{f}} F \Delta^+ f \right\} \\
\text{by (3.139)} &= \sum_{(a,y]} \Delta^- V + \sum_{[a,y)} \Delta^+ V \tag{3.141}
\end{aligned}$$

for all $a < y \leq b$. Therefore V is pure jump function of bounded variation. Further by (3.138), we have

$$\Delta^- V = (F_- + \Delta^- F) \left(e^{-\bar{f}} \right)_- e^{-\Delta^- f} - \left(F e^{-\bar{f}} \right)_- = V_- \left\{ (1 + \Delta^- f) e^{-\Delta^- f} - 1 \right\}$$

and

$$\Delta^+ V = (F + \Delta^+ F) e^{-\bar{f}} e^{-\Delta^+ f} - F e^{-\bar{f}} = V \left\{ (1 + \Delta^+ f) e^{-\Delta^+ f} - 1 \right\}.$$

By Taylor's formula (3.87), the function A defined by

$$A(y) := \sum_{(a,y]} \left\{ (1 + \Delta^- f) e^{-\Delta^- f} - 1 \right\} + \sum_{[a,y)} \left\{ (1 + \Delta^+ f) e^{-\Delta^+ f} - 1 \right\} \tag{3.142}$$

for $a \leq y \leq b$, is pure jump function of bounded variation. By Theorem 2.22, we have

$$(LY) \int_a^y V dA = \lim_{m \rightarrow \infty} S_{LC}(V, A; \lambda_m \mathbb{m}[a, y]) = \sum_{(a,y]} V_- \Delta^- A + \sum_{[a,y)} V \Delta^+ A = \sum_{(a,y]} \Delta^- V + \sum_{[a,y)} \Delta^+ V$$

for $a \leq y \leq b$. Thus by (3.141), the function V is a solution to the linear Left Young integral equation

$$V(y) = 1 + (LY) \int_a^y V dA, \quad a \leq y \leq b.$$

By Theorem 47, the solution is unique in the class of all regulated functions, which certainly contains the class of functions of bounded variation. Therefore

$$V(y) = e^{A(y) - A(a)} \prod_{[a,y]} (1 + \Delta A) e^{-\Delta A} = \prod_{[a,y]} (1 + \Delta f) e^{-\Delta f},$$

for $a \leq y \leq b$. That is $V = V_a = \gamma(f; a, \cdot)$, and hence $F = \mathcal{E}_a(f)$ as desired. The proof of Theorem 48 is complete. $\square$

Again for given $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, let f be a (λ, p) -decomposable function on $[a, b]$. This time we consider the backward linear Right λ -integral equation

$$G(y) = 1 + (R) \int_y^b G d_\lambda f, \quad a \leq y \leq b, \tag{3.143}$$

provided the Right λ -integral is defined on $[a, b]$ with respect to $D_{\lambda,p}^-(f)$. Using a chain rule it is easy to see that the backward Doléans exponential $\mathcal{E}_b(f)$ satisfy (3.143). Similarly to the forward case, we show that this function is the unique solution of an augmented equation.

Definition 50 For $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, let f be a (λ, p) -decomposable function on $[a, b]$, and let $D_{\lambda, p}^-(f)$ be nonempty. We say that a function $G \in \text{dual}(\mathcal{W}_p)[a, b] \cap Q_\lambda[a, b]$ is a *solution of the augmented backward linear Right λ -integral equation*

$$d_\lambda^- G = G d_\lambda^- f \quad \text{on } [a, b] \quad (3.144)$$

if (a) and (b) hold, where

- (a) for $(g, h) \in D_{\lambda, p}^-(f)$ and $\psi \in \text{dual}(\mathcal{W}_p)[a, b]$, $(RC) \int \langle (\psi, \psi G), d_\lambda(G, g) \rangle$ is defined on $[a, b]$, and for $a \leq y \leq b$,

$$(RC) \int_y^b \langle (\psi, \psi G), d_\lambda(G, g) \rangle = -(RY) \int_y^b \psi G dh; \quad (3.145)$$

- (b) the 2-vector function (G, f) has the quadratic λ -variation and for $a \leq y \leq b$,

$$\begin{pmatrix} [G]_\lambda & [f, G]_\lambda \\ [G, f]_\lambda & [f]_\lambda \end{pmatrix} (y) = (RY) \int_a^y \begin{pmatrix} G^2 & -G \\ -G & 1 \end{pmatrix} d[f]_\lambda. \quad (3.146)$$

Let f be a (λ, p) -decomposable function on $[a, b]$ for some $\lambda \in \Lambda[a, b]$ and $1 \leq p < 2$, and let $D_{\lambda, p}^-(f)$ be nonempty. Suppose that $G \in \text{dual}(\mathcal{W}_p)$, has the quadratic λ -variation and is a solution of the augmented backward linear Right λ -integral equation (3.144). Then in (a) of Definition 50 taking $\psi \equiv 1$, by Proposition 23, it follows that the Right λ -integral $(R) \int G d_\lambda f$ is defined on $[a, b]$ with respect to $D_{\lambda, p}^-(f)$, and (3.143) holds.

Theorem 51 Let $\lambda \in \Lambda[a, b]$, $1 \leq p < 2$ and let f be a (λ, p) -decomposable function on $[a, b]$. If $D_{\lambda, p}^-(f) \neq \emptyset$ then in the class $\text{dual}(\mathcal{W}_p) \cap Q_\lambda[a, b]$ the Doléans exponential $\mathcal{E}_b(f)$ is the unique solution of the augmented backward linear Right λ -integral equation (3.144).

Proof. Let $(g, h) \in D_{\lambda, p}^-(f)$. By Corollary 16, f has the quadratic λ -variation, and therefore the Doléans exponential $\mathcal{E}_b(f)$ is defined. Next we show that $\mathcal{E}_b(f)$ is a solution of the augmented backward linear Right λ -integral equation (3.144). By Theorem 45, the 2-vector function $(\mathcal{E}_b(f), f)$ has the quadratic λ -variation and (3.146) holds with $G = \mathcal{E}_b(f)$. Let $\bar{f} := f - f(b) + (1/2)[g]_\lambda^c(b) - (1/2)[g]_\lambda^c$, and $V_b := \gamma(f; \cdot, b)$, where $\gamma(f)$ is defined by (3.83). Then $\mathcal{E}_b(f) = e^{-\bar{f}} V_b$ and V_b is pure jump function of bounded variation by Lemma 43. Since the composition of the function $\bar{f}$ with a smooth function does not change its p -variation, and since $\mathcal{W}_p$ is a Banach algebra, $\mathcal{E}_b(f) \in \text{dual}(\mathcal{W}_p)[a, b]$, and so $\mathcal{E}_b(f)$ satisfies condition (b) of Definition 50. To show that $\mathcal{E}_b(f)$ satisfies condition (a), we apply the chain rule formula of Theorem 34.(b) to the composition $\phi \circ \mathbf{f} = \mathcal{E}_b(f)$, where the 2-vector function $\mathbf{f}$ is defined by

$$\mathbf{f} := (\bar{f}, V^b) = (-g(b) - h(b) + \frac{1}{2}[g]_\lambda^c(b), 0) + (g, 0) + (h - \frac{1}{2}[g]_\lambda^c, V^b) \quad \text{and} \quad \phi(u, v) = e^{-u}v$$

for $u, v \in \mathbb{R}$. Let $\mathbf{g} := (g, 0)$ and $\mathbf{h} := (h - (1/2)[g]_\lambda^c, V_b)$. Then $\mathbf{f}$ is (λ, p) -decomposable by $(\mathbf{g}, \mathbf{h})$. Since all discontinuity points of V_b are the same as discontinuity points of f , the left discontinuity points of $\mathbf{f}$ are accessible by λ . Let $\psi \in \text{dual}(\mathcal{W}_p)[a, b]$. By statement (b) of Theorem 34, the Right Cauchy λ -integral

$$(RC) \int \langle (\psi, -\psi(\nabla \phi \circ \mathbf{f})), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle = (RC) \int \langle (\psi, \psi \mathcal{E}_b(f)), d_\lambda(\mathcal{E}_b(f), g) \rangle$$

is defined on $[a, b]$, where $\nabla\phi(u, v) = (\frac{\partial\phi}{\partial u}, \frac{\partial\phi}{\partial v})(u, v) = (-e^{-u}v, e^{-u})$ for $u, v \in \mathbb{R}$, and the equalities

$$\begin{aligned}
(RC) \int_y^b \langle (\psi, \psi \mathcal{E}_b(f)), d\lambda(\mathcal{E}_b(f), g) \rangle \\
&= -(RY) \int_y^b \psi \mathcal{E}_b(f) d(h - (1/2)[g]^c) + (RY) \int_y^b \psi e^{-\bar{f}} dV^b - \frac{1}{2}(RS) \int_y^b \psi \mathcal{E}_b(f) d[g]^c \\
&+ \sum_{(y, b]} \psi \left\{ \Delta^- \mathcal{E}_b(f) + \mathcal{E}_b(f) \Delta^- f - e^{-\bar{f}} \Delta^- V^b \right\} \\
&+ \sum_{[y, b)} \psi_+ \left\{ \Delta^+ \mathcal{E}_b(f) + \mathcal{E}_b(f)_+ \Delta^+ f - e^{-\bar{f}_+} \Delta^+ V^b \right\} = (RY) \int_y^b \psi \mathcal{E}_b(f) dh \\
&+ (RY) \int_y^b \psi e^{-\bar{f}} dV^b - \left\{ \sum_{(y, b]} \psi e^{-\bar{f}} \Delta^- V^b + \sum_{[y, b)} \psi_+ e^{-\bar{f}_+} \Delta^+ V^b \right\} \tag{3.147}
\end{aligned}$$

hold for $a \leq y \leq b$. The last equality follows by Lemma 9 and relations (3.112), (3.113). Let $\lambda = \{\lambda_m: m \geq 1\}$ with $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$. Since V_b is pure jump function of bounded variation, and since $\{x \in (a, b): \Delta^- V_b(x) \neq 0\} \subset \cup_m \lambda_m$ by (3.115), by Theorem 2.22, we have the equalities

$$\begin{aligned}
(RY) \int_y^b \psi e^{-\bar{f}} dV^b &= \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} \psi(x_i^m) e^{-\bar{f}(x_i^m)} [V^b(x_i^m \vee y) - V^b(x_{i-1}^m \vee y)] \\
&= \sum_{(y, b]} \psi e^{-\bar{f}} \Delta^- V^b + \sum_{[y, b)} \psi_+ e^{-\bar{f}_+} \Delta^+ V^b
\end{aligned}$$

for $a \leq y \leq b$. Thus the last two terms in (3.147) give zero to the equality, and hence the Doléans exponential $\mathcal{E}_b(f)$ solves the augmented backward linear Right λ -integral equation (3.144).

Finally, suppose that $G \in \text{dual}(\mathcal{W}_p) \cap Q_\lambda[a, b]$ solves the augmented backward linear Right λ -integral equation (3.144). We then show that $G = \mathcal{E}_b(f)$. To this aim let $V := e^{\bar{f}} G$, where $\bar{f} := f - f(b) + (1/2)[g]_\lambda^c(b) - (1/2)[g]_\lambda^c$. It is enough to prove that V is equal to the pure jump function $V_b = \gamma(f; \cdot, b)$. Let

$$\mathbf{f} := (\bar{f}, G) = (C - f(b) + \frac{1}{2}[g]_\lambda^c(b), 0) + (g, G) + (h - \frac{1}{2}[g]_\lambda^c, 0) \quad \text{and} \quad \phi(u, v) := e^u v \tag{3.148}$$

for $u, v \in \mathbb{R}$. Then $V = \phi \circ \mathbf{f}$. We wish to apply a chain rule to show that V is the unique solution to the linear Right Young integral equation with respect to a pure jump function of bounded variation. First, we claim that the 2-vector function $\mathbf{g} := (g, G)$ has the quadratic λ -variation. By condition (b) of Definition 50, (f, G) has the quadratic λ -variation. To apply Lemma 49 we have to check its assumption (3.136). By condition (a) of Definition 50 with $\psi \equiv 1$, and by Proposition 23, the Right λ -integral $(R) \int G d_\lambda f$ is defined on $[a, b]$ with respect to $D_{\lambda, p}^-(f)$, and (3.143) holds. By Definition 18 of the Right Cauchy λ -integral in conjunction with Proposition 2.27 concerning jumps of the indefinite Right Young integral, we have that

$$\Delta^- G(y) = -(G \Delta^- f)(y) \quad \text{and} \quad \Delta^+ G(x) = -(G \Delta^+ f)(x) \tag{3.149}$$

for $a \leq y < x \leq b$. In particular, it then follows that (3.136) holds. Therefore by Lemma 49, the 2-vector function $\mathbf{g} = (g, G)$ has the quadratic λ -covariation, and thereby has the quadratic λ -variation. Since the pair $(\mathbf{g}, \mathbf{h})$ is (λ, p) -dual and (3.148) holds, $\mathbf{f}$ is (λ, p) -decomposable by $(\mathbf{g}, \mathbf{h})$. By (3.149), it follows that $(\mathbf{g}, \mathbf{h}) \in D_{\lambda, p}^-(\mathbf{f})$. Therefore all assumptions of Theorem 34.(b) are satisfied by $\mathbf{f}$ and ϕ . By the first part of statement (b) of Theorem 34 with $\psi \equiv 1$, the Right Cauchy λ -integrals

$$(RC) \int \langle (1, -\nabla \phi \circ \mathbf{f}), d_\lambda(\phi \circ \mathbf{f}, \mathbf{g}) \rangle = \mu(\phi \circ \mathbf{f}) - (RC) \int \langle \nabla \phi \circ \mathbf{f}, d_\lambda \mathbf{g} \rangle$$

are defined on $[a, b]$, where $\nabla \phi(u, v) = (\frac{\partial \phi}{\partial u}, \frac{\partial \phi}{\partial v})(u, v) = (e^u v, e^u)$ for $u, v \in \mathbb{R}$, and μ is defined by (2.2). Further, by condition (a) of Definition 50 with $\psi = \exp\{\bar{f}\}$, we have the equality

$$(RC) \int \langle \nabla \phi \circ \mathbf{f}, d_\lambda \mathbf{g} \rangle \equiv (RC) \int \langle (e^{\bar{f}} G, e^{\bar{f}}), d_\lambda(g, G) \rangle = -(RY) \int e^{\bar{f}} G dh. \quad (3.150)$$

By condition (b) of Definition 50, by the second part of Lemma 49 and by (3.149), we have

$$(RY) \int_a^y G d[f]_\lambda = -[f, G]_\lambda(y) = -[g, G]_\lambda^c(y) + \sum_{(a, y]} G \{\Delta^- f\}^2 + \sum_{[a, y)} G_+ \{\Delta^+ f\}^2$$

for $a \leq y \leq b$. On the other hand, by the second part of Lemma 3, and since $[f]_\lambda^c = [g]_\lambda^c$, we have

$$(RY) \int_a^y G d[f]_\lambda = (RS) \int_a^y G d[g]_\lambda^c + \sum_{(a, y]} G \{\Delta^- f\}^2 + \sum_{[a, y)} G_+ \{\Delta^+ f\}^2$$

for $a \leq y \leq b$. Therefore $[g, G]_\lambda^c(y) = (RS) \int_a^y G d[g]_\lambda^c$ for $a \leq y \leq b$. This in conjunction with the substitution theorem for the Riemann-Stieltjes integral, yields the equality

$$(RS) \int_y^b e^{\bar{f}} d[g, G]_\lambda^c = -(RS) \int_y^b e^{\bar{f}} G d[g]_\lambda^c \quad (3.151)$$

for $a \leq y \leq b$. By the second part of statement (b) of Theorem 34 with $\psi \equiv 1$, we then have

$$\begin{aligned} V(b) - V(y) &= (RC) \int_y^b \langle \nabla \phi \circ \mathbf{f}, d_\lambda \mathbf{g} \rangle + (RY) \int_y^b e^{\bar{f}} G d(h - (1/2)[g]_\lambda^c) \\ &\quad - \frac{1}{2}(RS) \int_y^b e^{\bar{f}} G d[g]_\lambda^c - (RS) \int_y^b e^{\bar{f}} d[g, G]_\lambda^c \\ &\quad + \sum_{(y, b]} \left\{ \Delta^- V - e^{\bar{f}} \Delta^- G - e^{\bar{f}} G \Delta^- f \right\} \\ &\quad + \sum_{[y, b)} \left\{ \Delta^+ V - (e^{\bar{f}})_+ \Delta^+ G - (e^{\bar{f}} G)_+ \Delta^+ f \right\} \end{aligned}$$

$$\text{by Lemma 9} \quad = (RC) \int_y^b \langle \nabla \phi \circ \mathbf{f}, d_\lambda \mathbf{g} \rangle + (RY) \int_y^b e^{\bar{f}} G dh$$

$$\text{by (3.151)} \quad - \frac{1}{2}(RS) \int_y^b e^{\bar{f}} G d[g]_\lambda^c - \frac{1}{2}(RS) \int_y^b e^{\bar{f}} G d[g]_\lambda^c + (RS) \int_y^b e^{\bar{f}} G d[g]_\lambda^c$$

$$\begin{aligned}
& \text{by (3.149)} \quad + \sum_{(y,b]} \left\{ \Delta^- V + e^{\bar{f}} G \Delta^- f - e^{\bar{f}} G \Delta^- f \right\} \\
& \quad + \sum_{[y,b)} \left\{ \Delta^+ V + \left(e^{\bar{f}} \right)_+ G_+ \Delta^+ f - \left(e^{\bar{f}} \right)_+ G_+ \Delta^+ f \right\} \\
& \text{by (3.150)} \quad = \sum_{(y,b]} \Delta^- V + \sum_{[y,b)} \Delta^+ V
\end{aligned}$$

for $a \leq y < b$. Therefore V is pure jump function of bounded variation satisfying the relation

$$V(y) = 1 - \sum_{(y,b]} \Delta^- V - \sum_{[y,b)} \Delta^+ V \quad (3.152)$$

for $a \leq y < b$. Further by (3.149), we have

$$\Delta^- V = G e^{\bar{f}} - (G - \Delta^- G) e^{\bar{f}} e^{-\Delta^- f} = V \left\{ 1 - (1 + \Delta^- f) e^{-\Delta^- f} \right\} = -V \Delta^- A$$

and

$$\Delta^+ V = \left(G e^{\bar{f}} \right)_+ - (G_+ - \Delta^+ G) \left(e^{\bar{f}} \right)_+ e^{-\Delta^+ f} = V_+ \left\{ 1 - (1 + \Delta^+ f) e^{-\Delta^+ f} \right\} = -V_+ \Delta^+ A,$$

where A is the pure jump function of bounded variation defined by (3.142). By Theorem 2.22, we have

$$\begin{aligned}
(RY) \int_y^b V dA &= \lim_{m \rightarrow \infty} S_{RC}(V, A; \lambda_m \cap [y, b]) = \sum_{(y,b]} V \Delta^- A + \sum_{[y,b)} V_+ \Delta^+ A \\
&= - \sum_{(y,b]} \Delta^- V - \sum_{[y,b)} \Delta^+ V
\end{aligned}$$

for $a \leq y \leq b$. Thus by (3.152), the function V is a solution to the linear Right Young integral equation

$$V(y) = 1 + (RY) \int_y^b V dA, \quad a \leq y \leq b.$$

By Theorem 5.22 of Dudley and Norvaiša [23, Part II], the solution is unique in the class of functions of bounded variation. Therefore

$$V(y) = e^{A(b)-A(y)} \prod_{[y,b]} (1 + \Delta A) e^{-\Delta A} = \prod_{[y,b]} (1 + \Delta f) e^{-\Delta f},$$

for $a \leq y \leq b$. That is $V = V_b = \gamma(f; \cdot, b]$, and hence $G = \mathcal{E}_b(f)$. The proof of Theorem 51 is complete. $\square$

3.7 The evolution representation

Now we are prepared to prove the main result of the paper as formulated by Theorem 1.5 in the introduction. Recall that a family $U = \{U(t, s) : a \leq s \leq t \leq b\}$ of real numbers $U(t, s)$ is called an *evolution* on $[a, b]$ if

$$\begin{cases} U(r, t) U(t, s) = U(r, s) & \text{for } a \leq s \leq t \leq r \leq b \\ U(s, s) = 1 & \text{for } a \leq s \leq b. \end{cases}$$

Definition 52 An evolution U on $[a, b]$ is called *regulated* if there exists a multiplicative upper continuous bounded and nondegenerate function $\pi = \pi_U$ on the extended simplex $S[[a, b]]$ such that $U(t, s) = \pi(s, t)$ for all $(s, t) \in S[a, b]$.

Recall that for a function f on $[a, b]$, $f \gg 0$ means that there is a constant $C > 0$ such that $f(t) \geq C$ for each $t \in [a, b]$.

Proposition 53 An evolution U on $[a, b]$ is regulated if and only if $U_a \equiv U(\cdot, a)$ is a regulated function on $[a, b]$ and $|U_a| \gg 0$.

Proof. First, let an evolution U on $[a, b]$ be regulated. Since U_a is the restriction of the right distribution function of π to $[a, b]$, U_a is regulated on $[a, b]$ by statement (iv) of Theorem 2.4. If $\inf\{|U_a(u)|: u \in [a, b]\} = 0$, then by compactness of $[a, b]$ it follows that $\pi(a, b) = 0$, a contradiction proving that $|U_a| \gg 0$.

Now, let U_a be regulated on $[a, b]$ and let $|U_a| \gg 0$. For each $(u, v) \in S[[a, b]]$, let $\pi(u, v) := U_a(u)/U_a(v)$. It is easy to see that π is a multiplicative, bounded and nondegenerate function on $S[[a, b]]$. By statement (iv) of Theorem 2.4, it then follows that π is upper-continuous. Since $U = \pi$ on $S[a, b]$, the conclusion follows. The proof of the proposition is complete. $\square$

Similarly one can show that an evolution U on $[a, b]$ is regulated if and only if the function $U_b := U(b, \cdot)$ is regulated on $[a, b]$ and $|U_b| \gg 0$.

As it is mentioned in the introduction, we are interested in evolutions satisfying property (1.10). We formulate next its restricted variant. For a function π on $S[[a, b]]$ and for a partition $\kappa = \{x_i: i = 0, \dots, n\}$ of an extended interval $[[u, v]]$, $a \leq u \leq v \leq b$, let

$$S(\pi; \kappa) := \sum_{i=1}^n [\pi(x_{i-1}, x_i) - 1].$$

Recalling Notation 3.2 of the trace partition $\kappa \mathfrak{m} [[u, v]]$ we have the following:

Definition 54 Let U be a regulated evolution on $[a, b]$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. We say that U has the λ -generator on $[a, b]$ if there exists an additive and upper continuous function ρ on $S[[a, b]]$ such that for each $(u, v) \in S[[a, b]]$,

$$\rho(u, v) = \lim_{m \rightarrow \infty} S(\pi_U; \lambda_m \mathfrak{m} [[u, v]]). \quad (3.153)$$

A restriction to $[a, b]$ of the right distribution function of ρ , that is the function R_ρ defined by (2.4), is called the λ -generator of U .

Also as it is mentioned in the introduction, the λ -generator of an evolution generalizes the notion of the generator of a one-parameter semigroup. Alternatively the λ -generator on $[a, b]$ can be defined solely in terms of a regulated function on $[a, b]$ as follows:

Proposition 55 Let U be a regulated evolution on $[a, b]$, and let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$. Then U has the λ -generator on $[a, b]$ if and only if there exists a regulated function G on $[a, b]$ such that $G(a) = 0$ and the following hold for each $a \leq s < t \leq b$:

$$G(t) - G(s) = \lim_{m \rightarrow \infty} S(U; \lambda_m \mathfrak{m} [s, t]), \quad (3.154)$$

$$\Delta^- G(t) = \pi_U(t, t-) - 1 \quad \text{and} \quad \Delta^+ G(s) = \pi_U(s+, s) - 1. \quad (3.155)$$

If the two statements hold then G is the λ -generator of U .

Proof. First suppose that U has the λ -generator, and let G be the λ -generator of U . Then $G(a) = 0$, G is regulated by statement (iv) of Theorem 2.4, G satisfies (3.154) and (3.155), and so the “only if” part of the conclusion holds.

For the converse, let G be a regulated function on $[a, b]$ such that $G(a) = 0$, and (3.154), (3.155) hold. For each $(u, v) \in S[[a, b]]$, let $\rho(u, v) := G(v) - G(u)$. Then ρ is additive and upper continuous function on $S[[a, b]]$ by Theorem 2.4. We prove (3.153) only for the two cases $(u, v) = (a, t-)$ and $(u, v) = (a, t+)$ because the proofs for the other cases are similar. To this aim let $\lambda_m = \{x_i^m: i = 0, \dots, n(m)\}$, $m = 1, 2, \dots$, and let $t \in (a, b]$. For each $m \geq 1$, there is an index $i(t) \in \{1, \dots, n(m) - 1\}$ such that $x_{i(t)}^m < t \leq x_{i(t)+1}^m$. Then by (3.154) and (3.155), we have

$$\begin{aligned} \lim_{m \rightarrow \infty} S(\pi_U; \lambda_m \cap [a, t-]) &= \lim_{m \rightarrow \infty} \left\{ S(U; \lambda_m \cap [a, t]) - \pi_U(t, x_{i(t)}^m) + \pi_U(x_{i(t)}^m, t-) \right\} \\ &= G(t) - \pi_U(t, t-) + 1 = G(t-) = \rho(a, t-), \end{aligned}$$

proving (3.153) for $(u, v) = (a, t-) \in S[[a, b]]$. Now let $t \in [a, b)$. By the definition of the trace partition, we have for each $m \geq 1$,

$$S(\pi_U; \lambda \cap [a, t+]) = S(\pi_U; \lambda \cap [a, t]) + \Delta^+ G(t),$$

proving (3.153) for $(u, v) = (a, t+) \in S[[a, b]]$. The proof of Proposition 55 is complete. $\square$

Given $\lambda \in \Lambda[a, b]$ and a regulated evolution U , we prove the existence of the λ -generator if U is strictly positive and if the right discontinuity points of $U_a \equiv U(\cdot, a)$ are accessible by λ in the sense of (1.17), that is, if U_a is an element of the set $L_\lambda[a, b]$ defined by (1.23). Also we show that the λ -generator G of U in this case has discontinuities strictly larger than -1 and the right discontinuity points of G are accessible by λ , that is, G is an element of the set $E_\lambda[a, b]$ defined by (1.24). Moreover, the λ -generator G of U has a Left Cauchy λ -integral representation.

Proposition 56 *Let $f \in L_\lambda[a, b]$ for some $\lambda \in \Lambda[a, b]$. Then the Left Cauchy λ -integral $(LC) \int f^{-1} d_\lambda f$ is defined on $[a, b]$, and for $a \leq t \leq b$,*

$$\begin{aligned} \mathcal{L}_\lambda f(t) &:= (LC) \int_a^t f^{-1} d_\lambda f \\ &= \log \frac{f(t)}{f(a)} + \frac{1}{2} (RS) \int_a^t \frac{d[f]_\lambda^c}{f^2} - \sum_{(a, t]} \left(\log \frac{f}{f_-} - \frac{\Delta^- f}{f_-} \right) - \sum_{[a, t)} \left(\log \frac{f_+}{f} - \frac{\Delta^+ f}{f} \right), \end{aligned} \tag{3.156}$$

where the two sums converge absolutely. Moreover, $\mathcal{L}_\lambda f \in E_\lambda[a, b]$ with the bracket function

$$[\mathcal{L}_\lambda f]_\lambda(t) = (LY) \int_a^t \frac{d[f]_\lambda}{f^2} \quad a \leq t \leq b. \tag{3.157}$$

Proof. The Left Cauchy λ -integral $(LC) \int f^{-1} d_\lambda f$ is defined and (3.156) holds by the chain rule of statement (a) in Theorem 24 applied to $\phi \circ f$ with $\phi(u) = \log u$ for $u > 0$. By Theorem 26 with the same ϕ , $\mathcal{L}_\lambda f$ has the quadratic λ -variation with the bracket function (3.157). By Definition 17 of the Left Cauchy λ -integral and by Proposition 19, $1 + \Delta^- \mathcal{L}_\lambda f \wedge \Delta^+ \mathcal{L}_\lambda f \gg 0$ since $f \gg 0$, $N_{(a, b)}(\Delta^+ \mathcal{L}_\lambda f) \subset \cup \lambda$, and so $\mathcal{L}_\lambda f \in E_\lambda[a, b]$. The proof is complete. $\square$

Now we are ready to prove existence of the λ -generator for a class of regulated evolutions.

Theorem 57 *Let U be a regulated evolution on $[a, b]$, and let $\lambda \in \Lambda[a, b]$. If $U_a \in L_\lambda[a, b]$ then on $[a, b]$, $\mathcal{L}_\lambda U_a$ is the λ -generator of U .*

Proof. Let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[a, b]$ be such $U_a \in L_\lambda[a, b]$. The function $\mathcal{L}_\lambda U_a$ is defined by Proposition 56. By Proposition 19, for $a \leq s < t \leq b$, we have

$$1 + \Delta^-(\mathcal{L}_\lambda U_a)(t) = 1 + \frac{\Delta^- U_a(t)}{U_a(t-)} = \frac{U_a(t)}{U_a(t-)} = \pi_U(t, t-) \quad (3.158)$$

and

$$1 + \Delta^+(\mathcal{L}_\lambda U_a)(s) = 1 + \frac{\Delta^+ U_a(s)}{U_a(s)} = \frac{U_a(s+)}{U_a(s)} = \pi_U(s+, s). \quad (3.159)$$

Thus by Proposition 55, it is enough to prove that for $a \leq s < t \leq b$,

$$\lim_{m \rightarrow \infty} S(U; \lambda_m \cap [s, t]) = (LC) \int_s^t U_a^{-1} d_\lambda U_a. \quad (3.160)$$

Fix $a \leq s < t \leq b$, and suppose that $\lambda_m \cap [s, t] = \{x_i^m: i = 0, \dots, n(m)\}$ for each $m \geq 1$. For $i = 1, \dots, n(m)$ and $m \geq 1$, letting

$$d(x_i^m, x_{i-1}^m) := U(x_i^m, x_{i-1}^m) - 1 - [\mathcal{L}_\lambda U_a(x_i^m) - \mathcal{L}_\lambda U_a(x_{i-1}^m)], \quad (3.161)$$

by additivity of the Left Cauchy λ -integral, it amounts to show that

$$\lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} d(x_i^m, x_{i-1}^m) = 0. \quad (3.162)$$

Let $\epsilon \in (0, 1/2)$. Since U_a is regulated and $U_a \in L_\lambda[a, b]$, there exists a partition $\zeta = \{z_j: j = 0, \dots, k\}$ of $[s, t]$ such that

$$\sum_{j=1}^k \sum_{(z_{j-1}, z_j)} \left((\Delta^- U_a)^2 + (\Delta^+ U_a)^2 \right) < \epsilon \quad \text{and} \quad \max_{1 \leq j \leq k} \text{Osc}(U_a; (z_{j-1}, z_j)) < \epsilon C, \quad (3.163)$$

where C is the constant from the definition of $L_\lambda[a, b]$ such that $|U_a(t)| \geq C$ for $t \in [a, b]$. Let I be the set of indices $j \in \{1, \dots, k-1\}$ such that $z_j \in \cup_m \lambda_m$, and let m_0 be the minimal integer such that each intersection $\lambda_m \cap (z_{j-1}, z_j)$, $j = 1, \dots, k$, contains at least two different points, and $\{z_j: j \in I\} \subset \lambda_m$. For each $m \geq m_0$ and $j = 0, \dots, k$, let $i(j) = i_m(j)$ be the index in $\{0, \dots, n(m)\}$ such that $x_{i(j)}^m = z_j$ if $z_j \in \cup_m \lambda_m$, or $x_{i(j)-1}^m < z_j < x_{i(j)}^m$ if $j \in \{1, \dots, k-1\} \setminus I$. Then let

$$I_1(\zeta, m) := \{i(j): j = 1, \dots, k\} \cup \{1, i(j)+1: j \in I\} \quad \text{and} \quad I_2(\zeta, m) := \{1, \dots, n(m)\} \setminus I_1(\zeta, m).$$

By the definition of $L_\lambda[a, b]$, $\Delta^+ U_a(z_j) = 0$ if $j \notin I$. Thus by (3.158), for each $j \in \{1, \dots, k\}$, we have

$$\lim_{m \rightarrow \infty} d(x_{i(j)}^m, x_{i(j)-1}^m) = \pi_U(z_j, z_j-) - 1 - \Delta^-(\mathcal{L}_\lambda U_a)(z_j) = 0.$$

For each $j \in \{0\} \cup I$, by (3.159), we have

$$\lim_{m \rightarrow \infty} d(x_{i(j)+1}^m, x_{i(j)}^m) = \pi_U(z_j+, z_j) - 1 - \Delta^+(\mathcal{L}_\lambda U_a)(z_j) = 0.$$

Therefore

$$\lim_{m \rightarrow \infty} \sum_{i \in I_1(\zeta, m)} d(x_i^m, x_{i-1}^m) = 0.$$

Since U_a has the quadratic λ -variation and $\{t \in (a, b): (\Delta^+ U_a)(t) \neq 0\} \subset \cup_m \lambda_m$, by Lemmas 11 and 9, we have

$$\lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^2 = (LY) \int_s^t \frac{d[U_a]_\lambda}{U_a^2} \quad (3.164)$$

$$= (RS) \int_s^t \frac{d[U_a]_\lambda^c}{U_a^2} + \sum_{(s,t]} \left(\frac{\Delta^- U_a}{(U_a)_-} \right)^2 + \sum_{[s,t)} \left(\frac{\Delta^+ U_a}{U_a} \right)^2. \quad (3.165)$$

For each $i \in I_2(\zeta, m)$, by the second relation in (3.163),

$$|1 - U_a(x_i^m)/U_a(x_{i-1}^m)| \leq C^{-1} \max_{1 \leq j \leq k} \text{Osc}(U_a; (z_{j-1}, z_j)) < \epsilon. \quad (3.166)$$

Thus by Taylor's theorem with remainder (3.108), for $i \in I_2(\zeta, m)$

$$\log \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} \right) = \frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 - \frac{1}{2} \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^2 + 3\theta_i^m \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^3 \quad (3.167)$$

with $|\theta_i^m| \leq 1$. By (3.161) and (3.156), for $i \in I_2(\zeta, m)$,

$$\begin{aligned} d(x_i^m, x_{i-1}^m) &= \frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 - \log \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} \right) - \frac{1}{2} (RS) \int_{x_{i-1}^m}^{x_i^m} \frac{d[U_a]_\lambda^c}{U_a^2} \\ &\quad + \sum_{(x_{i-1}^m, x_i^m]} \left(\log \frac{U_a}{(U_a)_-} - \frac{\Delta^- U_a}{(U_a)_-} \right) + \sum_{[x_{i-1}^m, x_i^m)} \left(\log \frac{(U_a)_+}{U_a} - \frac{\Delta^+ U_a}{U_a} \right) \\ \text{by (3.167)} \quad &= \frac{1}{2} \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^2 - 3\theta_i^m \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^3 - \frac{1}{2} (LY) \int_{x_{i-1}^m}^{x_i^m} \frac{d[U_a]_\lambda}{U_a^2} \\ \text{and (3.165)} \quad &+ \sum_{(x_{i-1}^m, x_i^m]} \left(\log \frac{U_a}{(U_a)_-} - \frac{\Delta^- U_a}{(U_a)_-} \right) + \frac{1}{2} \sum_{(x_{i-1}^m, x_i^m]} \left(\frac{\Delta^- U_a}{(U_a)_-} \right)^2 \\ &+ \sum_{[x_{i-1}^m, x_i^m)} \left(\log \frac{(U_a)_+}{U_a} - \frac{\Delta^+ U_a}{U_a} \right) + \frac{1}{2} \sum_{[x_{i-1}^m, x_i^m)} \left(\frac{\Delta^+ U_a}{U_a} \right)^2. \end{aligned} \quad (3.168)$$

As for (3.161), we conclude that

$$\lim_{m \rightarrow \infty} \sum_{i \in I_1(\zeta, m)} \left\{ \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^2 - (LY) \int_{x_{i-1}^m}^{x_i^m} \frac{d[U_a]_\lambda}{U_a^2} \right\} = 0.$$

By (3.166), we have

$$\limsup_{m \rightarrow \infty} \left| \sum_{i \in I_2(\zeta, m)} 3\theta_i^m \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^3 \right| \leq 3\epsilon C^{-2} [U_a]_\lambda.$$

By the Taylor series expansion (3.89) and by the first relation in (3.163), we have

$$\limsup_{m \rightarrow \infty} \left| \sum_{i \in I_2(\zeta, m)} \sum_{(x_{i-1}^m, x_i^m]} \left(\log \frac{U_a}{(U_a)_-} - \frac{\Delta^- U_a}{(U_a)_-} \right) \right| \leq \frac{2}{C^2} \limsup_{m \rightarrow \infty} \sum_{i \in I_2(\zeta, m)} \sum_{(x_{i-1}^m, x_i^m]} (\Delta^- U_a)^2 \leq \frac{2\epsilon}{C^2}.$$

Similar bounds hold for the last three terms in (3.168). All things considered, it then follows that the sum

$$\begin{aligned}
\sum_{i=1}^{n(m)} d(x_{i-1}^m, x_i^m) &= \sum_{i \in I_1(\zeta, m)} d(x_{i-1}^m, x_i^m) + \frac{1}{2} \left\{ \sum_{i=1}^{n(m)} \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^2 - (LY) \int_a^b \frac{d[U_a]_\lambda}{U_a^2} \right\} \\
&\quad - \frac{1}{2} \sum_{i \in I_1(\zeta, m)} \left\{ \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^2 - (LY) \int_{x_{i-1}^m}^{x_i^m} \frac{d[U_a]_\lambda}{U_a^2} \right\} - \sum_{i \in I_2(\zeta, m)} 3\theta_i^m \left(\frac{U_a(x_i^m)}{U_a(x_{i-1}^m)} - 1 \right)^3 \\
&\quad + \sum_{i \in I_2(\zeta, m)} \left\{ \sum_{(x_{i-1}^m, x_i^m]} \left(\log \frac{U_a}{(U_a)_-} - \frac{\Delta^- U_a}{(U_a)_-} \right) + \frac{1}{2} \sum_{(x_{i-1}^m, x_i^m]} \left(\frac{\Delta^- U_a}{(U_a)_-} \right)^2 \right\} \\
&\quad + \sum_{i \in I_2(\zeta, m)} \left\{ \sum_{[x_{i-1}^m, x_i^m)} \left(\log \frac{(U_a)_+}{U_a} - \frac{\Delta^+ U_a}{U_a} \right) + \frac{1}{2} \sum_{[x_{i-1}^m, x_i^m)} \left(\frac{\Delta^+ U_a}{U_a} \right)^2 \right\}
\end{aligned}$$

is small as m is large, proving (3.162). The proof of Theorem 57 is complete. $\square$

By Proposition 56, given $\lambda \in \Lambda[a, b]$, $\mathcal{L}_\lambda$ is a mapping from $L_\lambda[a, b]$ to $E_\lambda[a, b]$. The following statement shows that this mapping is one-to-one and the inverse mapping is the indefinite product λ -integral $\mathcal{P}_\lambda$.

Theorem 58 *Let $\lambda \in \Lambda[a, b]$ and let $f \in L_\lambda[a, b]$. Then both $\mathcal{L}_\lambda f$ and $\mathfrak{Y}(1 + d_\lambda(\mathcal{L}_\lambda f))$ are defined on $[a, b]$, and for each $a \leq y \leq z \leq b$,*

$$\mathfrak{Y}_y^z(1 + d_\lambda(\mathcal{L}_\lambda f)) = f(z)/f(y). \quad (3.169)$$

Proof. Let $\lambda = \{\lambda_m : m \geq 1\} \in \Lambda[a, b]$ be such that $f \in L_\lambda[a, b]$. The Left Cauchy λ -integral $(LC) \int f^{-1} d_\lambda f$ is defined on $[a, b]$ by Proposition 56. To prove the conclusion concerning the product λ -integral we use Proposition 3.39 with $G := f/f(a)$. It is clear that G is a regulated function on $[a, b]$, $G(a) = 1$ and $G(x) \neq 0$ for $x \in (a, b]$. Also by Proposition 19, for $a \leq y < z \leq b$, we have

$$\begin{cases} 1 + \Delta^-(\mathcal{L}_\lambda f)(z) = 1 + \Delta^- f(z)/f(z-) = G(z)/G(z-) \\ 1 + \Delta^+(\mathcal{L}_\lambda f)(y) = 1 + \Delta^+ f(y)/f(y) = G(y+)/G(y) \end{cases} \quad (3.170)$$

Therefore by Proposition 3.39, it is enough to prove that for each $a \leq y < z \leq b$,

$$\lim_{m \rightarrow \infty} |\log \{P(\mathcal{L}_\lambda f; \lambda_m \mathfrak{M} [y, z])\} - \log \{f(z)/f(y)\}| = 0. \quad (3.171)$$

Fix $a \leq y < z \leq b$, and suppose that $\lambda_m \mathfrak{M} [y, z] = \{x_i^m : i = 0, \dots, n(m)\}$. By the second part of Proposition 56, $\mathcal{L}_\lambda f$ has the quadratic λ -variation and

$$\lim_{m \rightarrow \infty} s_2(\mathcal{L}_\lambda f; \lambda_m \mathfrak{M} [y, z]) = (LY) \int_y^z \frac{d[f]_\lambda}{f^2} \quad (3.172)$$

$$= (RS) \int_y^z \frac{d[f]_\lambda^c}{f^2} + \sum_{(y, z]} \left(\frac{\Delta^- f}{f_-} \right)^2 + \sum_{[y, z)} \left(\frac{\Delta^+ f}{f} \right)^2, \quad (3.173)$$

where the second equality holds by Lemma 9. To begin the proof of (3.171) let $\epsilon > 0$. Since the sums in (3.156) and (3.173) converge unconditionally, and $\mathcal{L}_\lambda f$ is regulated, there exists a partition $\zeta = \{z_j: j = 0, \dots, k\}$ of $[y, z]$ such that for each finite set $\mu \subset [y, z]$ disjoint from ζ ,

$$\left| \sum_{\mu} \left(\log \frac{f}{f_-} - \frac{\Delta^- f}{f_-} \right) + \sum_{\mu} \left(\log \frac{f_+}{f} - \frac{\Delta^+ f}{f} \right) \right| < \epsilon, \quad (3.174)$$

$$\sum_{\mu} \left(\frac{\Delta^- f}{f_-} \right)^2 + \sum_{\mu} \left(\frac{\Delta^+ f}{f} \right)^2 < \epsilon \quad \text{and} \quad \text{Osc}(\mathcal{L}_\lambda f; (z_{j-1}, z_j)) < \epsilon \wedge \frac{1}{2}. \quad (3.175)$$

Let I be the set of indices $j \in \{1, \dots, k-1\}$ such that $z_j \in \cup_m \lambda_m$, and let m_0 be the minimal integer such that $\{z_j: j \in I\} \subset \lambda_m$ and each intersection $\lambda_m \cap (z_{j-1}, z_j)$, $j = 1, \dots, k$, contains at least two different points. For each $m \geq m_0$ and $j = 0, \dots, k$, let $i(j) = i_m(j)$ be the index in $\{0, \dots, n(m)\}$ such that $x_{i(j)}^m = z_j$ if $z_j \in \cup_m \lambda_m$, or $x_{i(j)-1}^m < z_j < x_{i(j)}^m$ otherwise. Then let

$$I_1(\zeta, m) := \{i(j): j = 0, \dots, k\} \cup \{1, i(j) + 1: j \in I\} \quad \text{and} \quad I_2(\zeta, m) := \{0, \dots, n(m)\} \setminus I_1(\zeta, m).$$

For each $m \geq m_0$, let $\Delta_i^m(\mathcal{L}_\lambda f) := \mathcal{L}_\lambda f(x_i^m) - \mathcal{L}_\lambda f(x_{i-1}^m) = (LC) \int_{x_{i-1}^m}^{x_i^m} f^{-1} d_\lambda f$, $i = 1, \dots, n(m)$, and

$$\begin{aligned} & \log \{P(\mathcal{L}_\lambda f; \lambda_m \cap [y, z])\} - \log \{f(z)/f(y)\} \\ &= \left[\log \left\{ \prod_{i \in I_1(\zeta, m)} (1 + \Delta_i^m(\mathcal{L}_\lambda f)) \right\} - \log \left\{ \prod_{i \in I_1(\zeta, m)} \frac{f(x_i^m)}{f(x_{i-1}^m)} \right\} \right] \\ &+ \left[\log \left\{ \prod_{i \in I_2(\zeta, m)} (1 + \Delta_i^m(\mathcal{L}_\lambda f)) \right\} - \log \left\{ \prod_{i \in I_2(\zeta, m)} \frac{f(x_i^m)}{f(x_{i-1}^m)} \right\} \right] =: D_1(\zeta, m) + D_2(\zeta, m). \end{aligned}$$

First consider the difference $D_1(\zeta, m)$. Since $\Delta^+ f(z_j) = 0$ for $z_j \notin \cup_m \lambda_m$, we have

$$\begin{aligned} \lim_{m \rightarrow \infty} D_1(\zeta, m) &= \log \left\{ \prod_{j=1}^k (1 + \Delta^-(\mathcal{L}_\lambda f)(z_j))(1 + \Delta^+(\mathcal{L}_\lambda f)(z_{j-1})) \right\} \\ &\text{by (3.170)} \quad - \log \left\{ \prod_{j=1}^k (1 + \left(\frac{\Delta^- f}{f_-} \right)(z_j))(1 + \left(\frac{\Delta^+ f}{f} \right)(z_{j-1})) \right\} = 0. \end{aligned}$$

Now we examine the difference $D_2(\zeta, m)$. By Taylor's theorem with remainder (3.108),

$$\begin{aligned} D_2(\zeta, m) &= \sum_{i \in I_2(\zeta, m)} \left\{ \Delta_i^m(\mathcal{L}_\lambda f) - \frac{1}{2}(\Delta_i^m(\mathcal{L}_\lambda f))^2 + 3\theta_i(\Delta_i^m(\mathcal{L}_\lambda f))^3 - \log \left(\frac{f(x_i^m)}{f(x_{i-1}^m)} \right) \right\} \\ \text{by (3.156)} &= \sum_{i \in I_2(\zeta, m)} \left\{ \frac{1}{2}(RS) \int_{x_{i-1}^m}^{x_i^m} \frac{d[f]_\lambda^c}{f^2} - \sum_{(x_{i-1}^m, x_i^m]} \left(\log \frac{f}{f_-} - \frac{\Delta^- f}{f_-} \right) \right. \\ &\quad \left. - \sum_{[x_{i-1}^m, x_i^m)} \left(\log \frac{f_+}{f} - \frac{\Delta^+ f}{f} \right) - \frac{1}{2}(\Delta_i^m(\mathcal{L}_\lambda f))^2 + 3\theta_i(\Delta_i^m(\mathcal{L}_\lambda f))^3 \right\}. \end{aligned}$$

By (3.173), we have

$$\begin{aligned}
& \sum_{i \in I_2(\zeta, m)} \left\{ (\Delta_i^m(\mathcal{L}_\lambda f))^2 - (RS) \int_{x_{i-1}^m}^{x_i^m} \frac{d[f]_\lambda^c}{f^2} \right\} \\
&= \left\{ s_2(\mathcal{L}_\lambda f; \lambda_m \mathbb{M}[y, z]) - (LY) \int_y^z \frac{d[f]_\lambda^c}{f^2} \right\} + \sum_{i \in I_1(\zeta, m)} (RS) \int_{x_{i-1}^m}^{x_i^m} \frac{d[f]_\lambda^c}{f^2} \\
&+ \sum_{i \in I_2(\zeta, m)} \left\{ \sum_{(x_{i-1}^m, x_i^m)} \left(\frac{\Delta^- f}{f_-} \right)^2 + \sum_{[x_{i-1}^m, x_i^m)} \left(\frac{\Delta^+ f}{f} \right)^2 - (\Delta_i^m(\mathcal{L}_\lambda f))^2 \right\} \\
&+ \sum_{i \in I_2(\zeta, m)} \left\{ \sum_{(x_{i-1}^m, x_i^m)} \left(\frac{\Delta^- f}{f_-} \right)^2 + \sum_{[x_{i-1}^m, x_i^m)} \left(\frac{\Delta^+ f}{f} \right)^2 \right\} =: \sum_{j=1}^4 A_j(m).
\end{aligned}$$

By (3.172), $A_1(m) \rightarrow 0$ as $m \rightarrow \infty$. By continuity of $[f]_\lambda^c$, $A_2(m) \rightarrow 0$ as $m \rightarrow \infty$. Since $\Delta^+ f(z_j) = 0$ for $z_j \notin \cup_m \lambda_m$, and by (3.32), it follows that $A_3(m) \rightarrow 0$ as $m \rightarrow \infty$. This in conjunction with (3.174) and (3.175) yields that

$$\begin{aligned}
\limsup_{m \rightarrow \infty} |D_2(\zeta, m)| &\leq 3[\mathcal{L}_\lambda f]_\lambda(b) \limsup_{m \rightarrow \infty} \max_{i \in I_2(\zeta, m)} |\Delta_i^m(\mathcal{L}_\lambda f)| + \frac{1}{2} \limsup_{m \rightarrow \infty} |A_4(m)| \\
&+ \limsup_{m \rightarrow \infty} \left| \sum_{i \in I_2(\zeta, m)} \left\{ \sum_{(x_{i-1}^m, x_i^m)} \left(\log \frac{f}{f_-} - \frac{\Delta^- f}{f_-} \right) + \sum_{[x_{i-1}^m, x_i^m)} \left(\log \frac{f_+}{f} - \frac{\Delta^+ f}{f} \right) \right\} \right| \\
&\leq 3\epsilon[\mathcal{L}_\lambda f]_\lambda(b) + \epsilon/2 + \epsilon.
\end{aligned}$$

All things considered, (3.171) holds. The proof of Theorem 58 is complete. $\square$

The following shows that the mapping $\mathcal{L}_\lambda$ from $L_\lambda[a, b]$ to $E_\lambda[a, b]$ is *onto*. Recall that the indefinite product λ -integral $\mathcal{P}_\lambda$ is defined by $\mathcal{P}_\lambda f(t) = \mathfrak{Y}_a^t(1 + d_\lambda f)$ for $t \in [a, b]$.

Theorem 59 *Let $\lambda \in \Lambda[a, b]$ and $f \in E_\lambda[a, b]$. Then $\mathfrak{Y}(1 + d_\lambda f)$ and $(LC) f(\mathcal{P}_\lambda f)^{-1} d_\lambda(\mathcal{P}_\lambda f)$ are defined on $[a, b]$, and for all $a \leq y \leq z \leq b$,*

$$(LC) \int_y^z (\mathcal{P}_\lambda f)^{-1} d_\lambda(\mathcal{P}_\lambda f) = f(z) - f(y). \quad (3.176)$$

Proof. Let $\lambda \in \Lambda[a, b]$ be such that $f \in E_\lambda[a, b]$. The product λ -integral $\mathfrak{Y}(1 + d_\lambda f)$ exists and equals to $\beta_\lambda(f)$ by Theorem 42. The function $\beta_\lambda(f)$ on $S[a, b]$ defined by (3.101) is nondegenerate, bounded, multiplicative and upper continuous by Corollary 41. Therefore $\mathcal{P}_\lambda(f)$ exists and equals to the forward Doléans exponential $\mathcal{E}_\lambda(f) := \mathcal{E}_{\lambda, a}(f)$. By Theorem 44, $\mathcal{E}_\lambda(f)$ has the quadratic λ -variation, and so $\mathcal{E}_\lambda(f) \in L_\lambda[a, b]$ since $f \in E_\lambda[a, b]$. Therefore the Left Cauchy λ -integral in (3.176) exists by Proposition 56. To prove (3.176) notice that for $a \leq s < t \leq b$,

$$\Delta^- \mathcal{L}_\lambda(\mathcal{P}_\lambda(f))(t) = (\mathcal{P}_\lambda(f)(t-))^{-1} \Delta^- \mathcal{P}_\lambda(f)(t) = \Delta^- f(t)$$

and

$$\Delta^+ \mathcal{L}_\lambda(\mathcal{P}_\lambda(f))(s) = (\mathcal{P}_\lambda(f)(s))^{-1} \Delta^+ \mathcal{P}_\lambda(f)(s) = \Delta^+ f(s).$$

Hence by Proposition 19, it is enough to prove (3.176) for ordinary points $a \leq y < z \leq b$. Let $a \leq y < z \leq b$, $\lambda = \{\lambda_m: m \geq 1\}$, and for each $m \geq 1$, let $\lambda_m \cap [y, z] = \{x_i^m: i = 0, \dots, n(m)\}$. Thus letting $\Delta_i^m \chi := \chi(x_i^m) - \chi(x_{i-1}^m)$, it is enough to prove that

$$\lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} \left\{ \beta_\lambda(f; x_{i-1}^m, x_i^m) - 1 - \Delta_i^m f \right\} = 0. \quad (3.177)$$

Let $\epsilon \in (0, 1/2)$. There exists a partition $\zeta = \{z_j: j = 0, \dots, k\}$ of $[y, z]$ such that for each $j = 1, \dots, k$,

$$\text{Osc}(f; (z_{j-1}, z_j)) < \epsilon, \quad \text{Osc}([f]_\lambda^c; [z_{j-1}, z_j]) < \epsilon \quad \text{and} \quad \sum_{j=1}^k \sum_{(z_{j-1}, z_j)} (\Delta f)^2 < \epsilon. \quad (3.178)$$

Let I be the set of indices $j \in \{1, \dots, k-1\}$ such that $z_j \in \cup_m \lambda_m$, and let m_0 be the minimal integer such that $\{z_j: j \in I\} \subset \lambda_m$ and each intersection $\lambda_m \cap (z_{j-1}, z_j)$, $j = 1, \dots, k$, contains at least two different points. For each $m \geq m_0$ and $j = 0, \dots, k$, let $i(j) = i_m(j)$ be the index in $\{0, \dots, n(m)\}$ such that $x_{i(j)}^m = z_j$ if $z_j \in \cup_m \lambda_m$, or $x_{i(j)-1}^m < z_j < x_{i(j)}^m$ otherwise. Then let

$$I_1(\zeta, m) := \{i(j): j = 0, \dots, k\} \cup \{1, i(j) + 1: j \in I\} \quad \text{and} \quad I_2(\zeta, m) := \{0, \dots, n(m)\} \setminus I_1(\zeta, m).$$

Since $\Delta^+ f(z_j) = 0$ for $z_j \notin \cup_m \lambda_m$, by (3.102), we have

$$\lim_{m \rightarrow \infty} \sum_{i \in I_1(\zeta, m)} \left\{ \beta_\lambda(f; x_{i-1}^m, x_i^m) - 1 - \Delta_i^m f \right\} = 0.$$

For each $i \in \{1, \dots, n(m)\}$ and $m \geq m_0$, we have

$$\begin{aligned} \beta_\lambda(f; x_{i-1}^m, x_i^m) - 1 - \Delta_i^m f &= \exp \left\{ \Delta_i^m f - \frac{1}{2} \Delta_i^m [f]_\lambda^c \right\} \left[\gamma(f; x_{i-1}^m, x_i^m) - 1 \right] \\ &\quad + \exp \left\{ \Delta_i^m f - \frac{1}{2} \Delta_i^m [f]_\lambda^c \right\} - 1 - \Delta_i^m f. \end{aligned}$$

By (3.86) and (3.178), there exists a finite constant $C = C(f)$ such that for each $m \geq m_0$,

$$\left| \sum_{i \in I_2(\zeta, m)} \exp \left\{ \Delta_i^m f - \frac{1}{2} \Delta_i^m [f]_\lambda^c \right\} \left[\gamma(f; x_{i-1}^m, x_i^m) - 1 \right] \right| \leq C \sum_{i \in I_2(\zeta, m)} \sum_{[x_{i-1}^m, x_i^m]} (\Delta f)^2 \leq \epsilon C.$$

By Taylor's theorem for the exponential function, for each $i \in \{1, \dots, n(m)\}$ and $m \geq m_0$, there is a θ_i^m , uniformly bounded in i and m , and such that

$$d_i^m := \exp \left\{ \Delta_i^m f - \frac{1}{2} \Delta_i^m [f]_\lambda^c \right\} - 1 - \Delta_i^m f = \frac{1}{2} \left\{ (\Delta_i^m f)^2 - \Delta_i^m [f]_\lambda^c \right\} + r_i^m,$$

where

$$r_i^m := -\frac{1}{2} \Delta_i^m f \Delta_i^m [f]_\lambda^c + \frac{1}{8} (\Delta_i^m [f]_\lambda^c)^2 + \theta_i^m \max \{ |\Delta_i^m f|, \Delta_i^m [f]_\lambda^c \} \left\{ (\Delta_i^m f)^2 + \Delta_i^m [f]_\lambda^c \right\}.$$

For each $m \geq m_0$, we have

$$\begin{aligned}
& \sum_{i \in I_2(\zeta, m)} \left\{ (\Delta_i^m f)^2 - \Delta_i^m [f]_\lambda^c \right\} \\
&= \left\{ s_2(f; \lambda_m \cap [y, z]) - \alpha_\lambda(f; y, z) \right\} + \sum_{i \in I_1(\zeta, m)} \Delta_i^m [f]_\lambda^c \\
&+ \sum_{i \in I_1(\zeta, m)} \left\{ \sum_{(x_{i-1}^m, x_i^m]} (\Delta^- f)^2 + \sum_{[x_{i-1}^m, x_i^m)} (\Delta^+ f)^2 - (\Delta_i^m f)^2 \right\} \\
&+ \sum_{i \in I_2(\zeta, m)} \left\{ \sum_{(x_{i-1}^m, x_i^m]} (\Delta^- f)^2 + \sum_{[x_{i-1}^m, x_i^m)} (\Delta^+ f)^2 \right\} =: \sum_{j=1}^4 A_j(m).
\end{aligned}$$

Since f has the quadratic λ -variation, $A_1(m) \rightarrow 0$ as $m \rightarrow \infty$. By continuity of $[f]_\lambda^c$, $A_2(m) \rightarrow 0$ as $m \rightarrow \infty$. Since $\Delta^+ f(z_j) = 0$ for $z_j \notin \cup_m \lambda_m$, it follows that $A_3(m) \rightarrow 0$ as $m \rightarrow \infty$. Therefore by (3.178), there exists a finite constant $C = C(f)$ such that

$$\limsup_{m \rightarrow \infty} \left| \sum_{i \in I_2(\zeta, m)} d_i^m \right| \leq \limsup_{m \rightarrow \infty} \left\{ A_4(m) + \sum_{i \in I_2(\zeta, m)} |r_i^m| \right\} \leq \epsilon + \epsilon C.$$

Since $\epsilon \in (0, 1/2)$ is arbitrary, (3.177) holds. The proof of Theorem 59 is complete. $\square$

Summing up we have:

Corollary 60 *For $\lambda \in \Lambda[a, b]$, the mapping $\mathcal{L}_\lambda$ from $L_\lambda[a, b]$ to $E_\lambda[a, b]$ is one-to-one and onto, with the inverse $\mathcal{P}_\lambda$.*

3.8 Non-existence of λ -integrals

In this section for a sequence λ of dyadic partitions of an interval $[0, 1]$, we show that the Left Cauchy and Right Cauchy λ -integrals do not exist for almost all pairs of sample functions of two jointly Gaussian but dependent Brownian motions.

Integration with respect to itself of a Brownian motion. Let $B = \{B(t); t \geq 0\}$ be a standard Brownian motion. It is well-known that the integral

$$\int_0^T B(t) dB(t), \quad 0 < T < \infty, \tag{3.179}$$

does not exist in the Riemann-Stieltjes sense. To see this notice that the difference of the two Riemann-Stieltjes sums evaluated at the same partition $\kappa = \{0 = t_0 < t_1 < \dots < t_n = T\}$ may be written as follows:

$$\sum_{i=1}^n B(t_i)[B(t_i) - B(t_{i-1})] - \sum_{i=1}^n B(t_{i-1})[B(t_i) - B(t_{i-1})] = s_2(B; \kappa).$$

It follows from Théorème 9 of Lévy [63, p. 516] that with probability 1,

$$\limsup_{\epsilon \downarrow 0} \{s_2(B; \kappa): \kappa \in \Xi[0, T], |\kappa| \leq \epsilon\} = +\infty,$$

where $|\kappa|$ denotes the mesh of $\kappa \in \Xi[0, T]$. Thus the integral (3.179) does not exist in the Riemann-Stieltjes sense.

Let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[0, T]$ and let $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ for $m \geq 1$. Since $B(0) = 0$, for each $m \geq 1$, we have

$$S_{LC}(B, B; \lambda_m) = \sum_{i=1}^{n(m)} B(t_{i-1}^m)[B(t_i^m) - B(t_{i-1}^m)] = \frac{1}{2}[B(T)^2 - s_2(B; \lambda_m)].$$

Moreover, by Théorème 5 of Lévy [63, p. 510], the limit $\lim_{m \rightarrow \infty} s_2(B; \lambda_m) = T$ exists with probability 1. Thus with probability 1, there exists the limit

$$(LC) \int_0^T B d_\lambda B = \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} B(t_{i-1}^m)[B(t_i^m) - B(t_{i-1}^m)] = \frac{1}{2}[B(T)^2 - T].$$

We show next that such a limit may be infinite if a pair of Brownian motions (B, B) is replaced by a pair of two different Brownian motions.

A special construction. Let L be a linear isometry between Hilbert spaces $L^2[0, 1]$ and $L^2(\Omega, \text{Pr})$. Then $B(t) := L(1_{[0, t]})$, $0 \leq t \leq 1$, is the Brownian motion stochastic process. The trigonometric system

$$1, \sqrt{2} \cos(2\pi t), \sqrt{2} \sin(2\pi t), \dots, c_k(t) := \sqrt{2} \cos(2\pi kt), s_k(t) := \sqrt{2} \sin(2\pi kt), \dots \quad (3.180)$$

is the basis of $L^2[0, 1]$. Thus L maps the trigonometric system onto a sequence of independent identically distributed (iid) standard normal random variables $\xi_0, \xi_1, \eta_1, \dots, \xi_k, \eta_k, \dots$. Let

$$1_{[0, t]} = a_0(t) + \sum_{k=1}^{\infty} \left\{ a_k(t) c_k(t) + b_k(t) s_k(t) \right\}$$

be the Fourier series representation of the indicator function $1_{[0, t]}$ in $L^2[0, 1]$, so that $a_0(t) = \langle 1_{[0, t]}, 1 \rangle = t$,

$$a_k(t) = \langle 1_{[0, t]}, c_k \rangle = \frac{1}{2\pi k} s_k(t) \quad \text{and} \quad b_k(t) = \langle 1_{[0, t]}, s_k \rangle = \frac{1}{2\pi k} (\sqrt{2} - c_k(t)).$$

Then the Fourier-Wiener series representation of the Brownian motion B is given by

$$\begin{aligned} B(t) &= \xi_0 t + \frac{1}{2\pi} \sum_{k=1}^{\infty} \frac{1}{k} \left\{ \xi_k s_k(t) + \eta_k (\sqrt{2} - c_k(t)) \right\} \\ &= \theta_0 t + \frac{\sqrt{2}}{\pi} \sum_{k=1}^{\infty} \theta_k \frac{\sin(\pi kt)}{k}, \quad 0 \leq t \leq 1 \end{aligned} \quad (3.181)$$

where $\theta_0 \equiv \xi_0$ and $\theta_k := [\xi_k s_k(t) + \eta_k (\sqrt{2} - c_k(t))]/2s_k(t/2)$, $k = 1, 2, \dots$ are iid standard normal random variables. Indeed, we then have

$$EB(t)B(s) = ts + \frac{2}{\pi^2} \sum_{k=1}^{\infty} \frac{\sin(\pi kt) \sin(\pi ks)}{k^2} = t \wedge s.$$

The right side of (3.181) without the linear term $\xi_0 t$ has the distribution of a Brownian bridge. The Fourier-Wiener series representation of a Brownian bridge was used in Paley, Wiener and Zygmund [91] and in Paley and Wiener [90, Chapter IX] to prove several of its sample function properties. For example, by Theorem XLIII of Paley and Wiener [90], a subsequence of partial sums of (3.181) converges uniformly to a limit almost surely which yields its almost sure continuity. Almost sure uniform convergence of the sequence of partial sums to its limit was proved by Hunt [50] for a more general class of random series. Later on the almost sure uniform convergence of (3.181) was proved by several authors using different methods including general methods of highly developed theory of probability distributions on Banach spaces. A new approach to proving this fact have been recently suggested by Kwapień and Woyczyński [61, Section 2.5]. The series

$$C(t) := \frac{1}{2\pi} \sum_{k=1}^{\infty} \frac{1}{k} \left\{ \eta_k s_k(t) - \xi_k(\sqrt{2} - c_k(t)) \right\}, \quad 0 \leq t \leq 1.$$

is called the series conjugate to (3.181) because C is the imaginary part and B is the real part of the complex series

$$Z(t) = \zeta_0 t + \frac{1}{\pi} \sum_{k=1}^{\infty} \frac{\zeta_k}{ik} \left(1 - e^{-2\pi i k t} \right), \quad 0 \leq t \leq 1,$$

where $\zeta_k = (\xi_k + i\eta_k)/\sqrt{2}$, $k = 0, 1, \dots$. As for B , one can check that $EC(t)C(s) = t \wedge s - ts$, that is C has the distribution of a Brownian bridge. For $0 \leq s, t \leq 1$, we have

$$EB(t)C(s) = \frac{1}{2\pi^2} \sum_{k=1}^{\infty} \frac{1}{k^2} \left\{ \sin(2\pi k(t-s)) - [\sin(2\pi kt) - \sin(2\pi ks)] \right\}.$$

Thus $B(t)$ and $C(s)$ are independent if either $t = s \in (0, 1)$ or $t, s \in \{0, 1\}$.

A behaviour of the Riemann-Stieltjes sums of $S_{RS}(B, C; \kappa)$ is defined by a behaviour of the Riemann-Stieltjes sums $S_{RS}(Y, X; \kappa)$, where

$$X(t) := B(t) - \xi_0 t - \frac{1}{\sqrt{2}\pi} \sum_{k=1}^{\infty} \frac{\eta_k}{k} = \frac{1}{2\pi} \sum_{k=1}^{\infty} \frac{1}{k} \left\{ \xi_k s_k(t) - \eta_k c_k(t) \right\}, \quad 0 \leq t \leq 1.$$

and

$$Y(t) := C(t) + \frac{1}{\sqrt{2}\pi} \sum_{k=1}^{\infty} \frac{\xi_k}{k} = \frac{1}{2\pi} \sum_{k=1}^{\infty} \frac{1}{k} \left\{ \eta_k s_k(t) + \xi_k c_k(t) \right\}, \quad 0 \leq t \leq 1.$$

It is easy to check that for any tagged partition κ of $[0, 1]$

$$S_{RS}(C, B; \kappa) = \frac{-\xi_0}{\sqrt{2}\pi} \sum_{k=1}^{\infty} \frac{\xi_k}{k} + S_{RS}(Y, X; \kappa) \quad \text{and} \quad S_{RS}(Y, X; \kappa) = S^1(\kappa) + S^2(\kappa), \quad (3.182)$$

where

$$S^1(\kappa) := \frac{1}{4\pi^2} \sum_{k,l=1}^{\infty} \left\{ \frac{\xi_k \xi_l}{kl} S_{RS}(c_k, s_l; \kappa) - \frac{\eta_k \eta_l}{kl} S_{RS}(s_k, c_l; \kappa) \right\}, \quad (3.183)$$

$$S^2(\kappa) := \frac{1}{4\pi^2} \sum_{k,l=1}^{\infty} \left\{ \frac{\eta_k \xi_l}{kl} S_{RS}(s_k, s_l; \kappa) - \frac{\xi_k \eta_l}{kl} S_{RS}(c_k, c_l; \kappa) \right\}. \quad (3.184)$$

For $0 \leq s, t \leq 1$, we have

$$EX(t)Y(s) = \frac{1}{2\pi^2} \sum_{k=1}^{\infty} \frac{\sin(2\pi k(t-s))}{k^2}.$$

Thus the dependence structure between X and Y is the same as between B and C . Also, for $0 \leq s, t \leq 1$, we have

$$EX(t)X(s) = EY(t)Y(s) = \frac{1}{2\pi^2} \sum_{k=1}^{\infty} \frac{\cos(2\pi k(t-s))}{k^2}.$$

The random series X and Y are examples of the class of Gaussian Fourier series

$$\sum_{k=0}^{\infty} a_k \{\xi_k \cos(2\pi kt) + \eta_k \sin(2\pi kt)\}, \quad 0 \leq t \leq 1,$$

for a sequence $\{a_k: k \geq 1\}$ of non-negative real numbers (see Section 14 in Kahane [53]). By known results for Gaussian Fourier series, the series X and Y are Fourier series of continuous functions almost surely. Clearly, their sample functions have similar properties to sample functions of a Brownian motion. In particular, for a $\lambda \in \Lambda[0, 1]$, almost all sample functions of X and Y have the quadratic λ -variation and their bracket functions $[X]_{\lambda}(t) = [Y]_{\lambda}(t) = t$ for $0 \leq t \leq 1$.

Theorem 61 *Let λ be the sequence of partitions $\lambda_m = \{i2^{-m}: i = 0, \dots, 2^m\}$, $m \geq 1$, of $[0, 1]$. Then $S_{LC}(Y, X; \lambda_m)$ and $S_{RC}(Y, X; \lambda_m)$ do not converge in probability to finite limits as $m \rightarrow \infty$.*

Proof. More generally, we show that the statement of the theorem holds for any subsequence of the sequence of partitions $\kappa_n = \{i/n: i = 0, \dots, n\}$, $n \geq 1$. For $n \geq 1$, let

$$G_n := \frac{n}{4\pi^2} \sum_{(k,l) \in C_-(n) \cup C_+(n)} \left\{ \frac{\xi_k \xi_l}{kl} \sin\left(\frac{2\pi l}{n}\right) + \frac{\eta_k \eta_l}{kl} \sin\left(\frac{2\pi k}{n}\right) \right\}$$

and

$$F_n := \frac{n}{4\pi^2} \sum_{(k,l) \in C(n)} \left\{ \frac{\xi_k \eta_l + \eta_k \xi_l}{kl} (1 - \cos\left(\frac{2\pi l}{n}\right)) \right\} = \frac{n}{4\pi^2} \sum_{(k,l) \in C(n)} \left\{ \frac{\xi_k \eta_l + \eta_k \xi_l}{kl} (1 - \cos\left(\frac{2\pi k}{n}\right)) \right\},$$

where $C(n) := \{(k, l) \in \mathbb{N} \times \mathbb{N}: k + l \in n\mathbb{N}\}$ and

$$\begin{cases} C_-(n) := \{(k, l) \in \mathbb{N} \times \mathbb{N}: k - l \in n\mathbb{Z} \text{ and } k + l \notin n\mathbb{N}\} \\ C_+(n) := \{(k, l) \in \mathbb{N} \times \mathbb{N}: k - l \notin n\mathbb{Z} \text{ and } k + l \in n\mathbb{N}\}. \end{cases} \quad (3.185)$$

Lemma 62 *For an integer $n \geq 1$,*

$$S_{LC}(Y, X; \kappa_n) = G_n + F_n \quad \text{and} \quad S_{RC}(Y, X; \kappa_n) = G_n - F_n.$$

Proof. Let n be a positive integer. Let $\kappa_n^l = \{((i-1)/n, i/n], (i-1)/n): i = 1, \dots, n\}$ and $\kappa_n^r = \{((i-1)/n, i/n], i/n): i = 1, \dots, n\}$ be two tagged partitions of $[0, 1]$. Then by the second relation in (3.182), we have

$$S_{LC}(Y, X; \kappa_n) = S^1(\kappa_n^l) + S^2(\kappa_n^l) \quad \text{and} \quad S_{RC}(Y, X; \kappa_n) = S^1(\kappa_n^r) + S^2(\kappa_n^r), \quad (3.186)$$

where functions S^1 and S^2 are defined by (3.183) and (3.184), respectively. For functions ϕ, ψ defined on $[0, 1]$, let $\Sigma^n \phi := \sum_{i=1}^n \phi(i/n)$, and $(\Delta \psi)(i/n) := \psi(i/n) - \psi((i-1)/n)$ for $i = 1, \dots, n$. Notice that if ϕ is periodic: $\phi(0) = \phi(1)$, then $\Sigma^n \phi = \sum_{i=1}^n \phi((i-1)/n)$. For each positive integer l , let $s_{l,n} := s_l(1/n) = \sqrt{2} \sin(2\pi l/n)$ and $c_{l,n} := \sqrt{2} - c_l(1/n) = \sqrt{2}(1 - \cos(2\pi l/n))$. We have by trigonometric identities for each l and $i = 1, \dots, n$

$$\Delta s_l\left(\frac{i}{n}\right) = \begin{cases} [c_l\left(\frac{i}{n}\right)s_{l,n} + s_l\left(\frac{i}{n}\right)c_{l,n}]/\sqrt{2} \\ [c_l\left(\frac{i-1}{n}\right)s_{l,n} - s_l\left(\frac{i-1}{n}\right)c_{l,n}]/\sqrt{2} \end{cases} \quad (3.187)$$

and

$$\Delta c_l\left(\frac{i}{n}\right) = \begin{cases} [c_l\left(\frac{i}{n}\right)c_{l,n} - s_l\left(\frac{i}{n}\right)s_{l,n}]/\sqrt{2} \\ -[c_l\left(\frac{i-1}{n}\right)c_{l,n} + s_l\left(\frac{i-1}{n}\right)s_{l,n}]/\sqrt{2}. \end{cases} \quad (3.188)$$

Recall the formulas

$$D_n(x) = \frac{1}{2} + \sum_{i=1}^n \cos ix = \frac{\sin(n+1/2)x}{2 \sin x/2} \quad \text{and} \quad \tilde{D}_n(x) = \sum_{i=1}^n \sin ix = \frac{\cos x/2 - \cos(n+1/2)x}{2 \sin x/2}$$

for Dirichlet's kernel D_n and Dirichlet's conjugate kernel $\tilde{D}_n$ respectively, valid for all $x \in (0, 2\pi)$. Then we have

$$\sum_{i=1}^n \cos(2\pi ji/n) = \begin{cases} 0 & \text{if } j \notin n\mathbb{Z} \\ n & \text{if } j \in n\mathbb{Z} \end{cases} \quad \text{and} \quad \sum_{i=1}^n \sin(2\pi ji/n) = 0 \quad \text{for any integer } j. \quad (3.189)$$

By the preceding relation it follows that for each $k, l \in \mathbb{N}$,

$$\Sigma^n c_k s_l = \sum_{i=1}^n [\sin(2\pi(l-k)i/n) + \sin(2\pi(l+k)i/n)] = 0 \quad (3.190)$$

and

$$\Sigma^n s_k c_l = \sum_{i=1}^n [\sin(2\pi(k-l)i/n) + \sin(2\pi(k+l)i/n)] = 0. \quad (3.191)$$

Recalling definition (3.185) of $C_- = C_-(n)$ and $C_+ = C_+(n)$, by (3.187) and (3.190), we have for each k and l ,

$$S_{LC}(c_k, s_l; \kappa_n) = S_{RC}(c_k, s_l; \kappa_n) = \frac{s_{l,n}}{2} \Sigma^n (c_{k-l} + c_{k+l}) = \begin{cases} n \frac{s_{l,n}}{\sqrt{2}} & \text{if } (k, l) \in C_- \cup C_+ \\ 0 & \text{otherwise.} \end{cases}$$

The last equation holds by (3.189) and because $s_{l,n} = 0$ if $k-l \in n\mathbb{Z}$ and $k+l \in n\mathbb{Z}$. Likewise by (3.188), (3.191) and (3.189), we have

$$S_{LC}(s_k, c_l; \kappa_n) = S_{RC}(s_k, c_l; \kappa_n) = -\frac{s_{l,n}}{2} \Sigma^n (c_{k-l} - c_{k+l}) = \begin{cases} -n \frac{s_{k,n}}{\sqrt{2}} & \text{if } (k, l) \in C_- \cup C_+ \\ 0 & \text{otherwise} \end{cases}$$

because $s_{l,n} = s_{k,n}$ if $(k, l) \in C_-$, and $s_{l,n} = -s_{k,n}$ if $(k, l) \in C_+$. Thus we arrive at formulas for the first terms on the right sides of (3.186)

$$S^1(\kappa_n^l) = S^1(\kappa_n^r) = G_n. \quad (3.192)$$

Next by (3.187) and (3.190), for each k and l , we have

$$S_{LC}(c_k, c_l; \kappa_n) = -\frac{c_{l,n}}{2} \Sigma^n(c_{k-l} + c_{k+l}) = -S_{RC}(c_k, c_l; \kappa_n).$$

Likewise, by (3.188) and (3.191), we have

$$S_{LC}(s_k, s_l; \kappa_n) = -\frac{c_{l,n}}{2} \Sigma^n(c_{k-l} - c_{k+l}) = -S_{RS}(s_k, s_l; \kappa_n).$$

Therefore by the first relation in (3.189), it follows that

$$\begin{aligned} S_2(\kappa_n^l) &= \frac{1}{4\pi^2} \sum_{k,l=1}^{\infty} \frac{\xi_k \eta_l - \xi_l \eta_k}{kl} \frac{c_{l,n}}{2} \Sigma^n c_{k-l} + \sum_{k,l=1}^{\infty} \frac{\xi_k \eta_l + \xi_l \eta_k}{kl} \frac{c_{l,n}}{2} \Sigma^n c_{k+l} \\ &= \frac{n}{4\pi^2} \sum_{(k,l) \in C} \frac{\xi_k \eta_l + \xi_l \eta_k}{kl} \frac{c_{l,n}}{\sqrt{2}}. \end{aligned}$$

The first series gives zero because $(\xi_k \eta_l - \xi_l \eta_k) c_{l,n} = -(\xi_l \eta_k - \xi_k \eta_l) c_{k,n}$ for each pair of positive integers (k, l) such that $k - l \in n\mathbb{Z}$. Thus $S_2(\kappa_n^l) = F_n$. Likewise it follows that $S_2(\kappa_n^r) = -F_n$. This in conjunction with (3.192) and (3.186) yields the claim of Lemma 62. $\square$

Suppose that a subsequence of random variables $\{Z_{n_k} : k \geq 1\}$ converges in probability to a random variable Z as $k \rightarrow \infty$. Since for each k and any $0 < M < \infty$,

$$\Pr(\{|Z_{n_k}| > M\}) \leq \Pr(\{|Z| > M/2\}) + \Pr(\{|Z_{n_k} - Z| > M/2\}),$$

it then follows that $\limsup_k \Pr(\{|Z_{n_k}| < M\}) > 0$ for all large enough M . Therefore to prove the theorem it is enough to prove that for each $M < \infty$,

$$\lim_{n \rightarrow \infty} \Pr(\{|Z_n| < M\}) = 0 \quad (3.193)$$

if Z_n is either $S_{LC}(Y, X; \kappa_n)$ or $S_{RC}(Y, X; \kappa_n)$. For such Z_n , we show that

$$EZ_n \geq C_1 \ln n \quad \text{and} \quad \sup_n \text{Var}(Z_n) \leq C_2 \quad (3.194)$$

for some finite constants C_1 and C_2 . Then assuming this is true, for any $0 < M, R < \infty$,

$$\begin{aligned} \Pr(\{|Z_n| < M\}) &\leq \Pr(\{|EZ_n| < M + |Z_n - EZ_n|, |Z_n - EZ_n| < R\}) \\ &\quad + \Pr(\{|Z_n - EZ_n| \geq R\}) \\ &\leq \Pr(\{|EZ_n| < M + R\}) + R^{-2} \text{Var}(Z_n). \end{aligned}$$

Taking R large enough one can see that (3.194) yields (3.193).

We prove (3.194) only for $Z_n = S_{LC}(Y, X; \kappa_n)$, $n \geq 1$, because a proof for the other case is the same. By Lemma 62, since $E(\xi_k \eta_l) = E(\eta_k \xi_l) = 0$ for all k, l and $E(\xi_k \xi_l) = E(\eta_k \eta_l) = 0$ for $k \neq l$, the expectation of Z_n ,

$$EZ_n = EG_n + EF_n = \frac{n}{2\pi^2} \sum_{(k,k) \in C_-(n)} \frac{1}{k^2} \sin\left(\frac{2\pi k}{n}\right) = \frac{n}{2\pi^2} \sum_{k=1}^{\infty} \frac{1}{k^2} \sin\left(\frac{2\pi k}{n}\right),$$

because $\sin(2\pi k/n) = 0$ for $2k \in n\mathbb{N}$. For $0 \leq x \leq \pi/2$, $\sin x \geq \frac{2}{\pi}x$. Thus for $k \leq n/4$ we have $\sin(2\pi k/n) \geq 4k/n$. For $n/4 < k \leq n/2$, $\sin(2\pi k/n) \geq 0$, and $\sum_{k \geq n/2} k^{-2} \leq 2/(n-2)$. Thus for all $n \geq 4$

$$EZ_n \geq \frac{2}{\pi^2} \left(\sum_{1 \leq k \leq n/4} \frac{1}{k} \right) - \frac{n}{2\pi^2} \frac{2}{n-2} \geq \frac{2}{\pi^2} (\ln n - 1),$$

and hence the first inequality in (3.194) holds. For the variance of Z_n , we have $\sqrt{\text{Var}(Z_n)} \leq 4^{-1}\pi^{-2}(2\sqrt{V_1} + 4\sqrt{V_2} + 2\sqrt{V_3} + 2\sqrt{V_4})$ for suitable $V_1, \dots, V_4$ to be bounded next for each $n \geq 1$. Since $\sin^2 x \leq \min(1, x^2)$ for all x , we have

$$V_1 = \text{Var} \left\{ n \sum_{k=1}^{\infty} \frac{\xi_k^2}{k^2} \sin \left(\frac{2\pi k}{n} \right) \right\} = n^2 \sum_{k=1}^{\infty} \frac{E(\xi_k^2 - 1)^2}{k^4} \sin^2 \left(\frac{2\pi k}{n} \right) \leq 16\pi^2.$$

Since $\sin(2\pi l/n) = 0$ if $k+l \in n\mathbb{N}$ and $k-l \in n\mathbb{Z}$, we have

$$V_2 = E \left\{ n \sum_{(k,l) \in C_-(n), k>l} \frac{\xi_k \xi_l}{kl} \sin \left(\frac{2\pi l}{n} \right) \right\}^2 = n^2 \sum_{m,l=1}^{\infty} \frac{\sin^2(2\pi l/n)}{(l+mn)^2 l^2} \leq n^2 \sum_{m,l=1}^{\infty} \frac{1}{m^2 n^2 l^2} \leq 4.$$

Denoting the integer part of an r by $[r]$, likewise we get

$$V_3 = E \left\{ n \sum_{(k,l) \in C_+(n)} \frac{\xi_k \xi_l}{kl} \sin \left(\frac{2\pi l}{n} \right) \right\}^2 = n^2 \sum_{k=1}^{\infty} \sum_{m=[k/n]+1}^{\infty} \frac{\sin^2(2\pi k/n)}{k^2 (nm-k)^2} \leq 12\pi^2 + 2.$$

Since $1 - \cos x = 2 \sin^2(x/2) \leq \min(2, x^2/2)$ for all x , we have

$$V_4 = E \left\{ n \sum_{(k,l) \in C_+(n)} \frac{\xi_k \eta_l}{kl} (1 - \cos \left(\frac{2\pi l}{n} \right)) \right\}^2 = n^2 \sum_{k=1}^{\infty} \sum_{m=[k/n]+1}^{\infty} \frac{(1 - \cos(2\pi k/n))^2}{k^2 (nm-k)^2} \leq 12\pi^4 + 2.$$

Thus $\text{Var}(Z_n) < \pi^2$, and hence (3.194) holds proving the Theorem 61. $\square$

Chapter 4

Extension of the class of semimartingales

Mathematics is endangered by a loss of unity and interaction

Richard Courant (1888-1972)

A stochastic process X which is decomposable into a sum of a local martingale M and an adapted càdlàg stochastic process A with locally bounded variation is called a semimartingale. In this chapter we consider a stochastic process X with the same decomposition property except that A is assumed to have locally bounded p -variation for some $1 \leq p < 2$, so that A may have almost all sample functions with *unbounded* variation. We call the extended semimartingale X , a $(2 - \epsilon)$ -semimartingale (see Definition 8 below). The main results are that an extended stochastic integral with respect to X and a continuous part of an extended quadratic variation of X are *invariant* with respect to possibly different decompositions $X - X_0 = M + A$. The new constructions in conjunction with the results of Chapter 2 concerning the refinement Riemann-Stieltjes integral are then used to describe the unique solution to the linear extended stochastic integral equation with respect to a $(2 - \epsilon)$ -semimartingale.

4.1 Stochastic processes and p -variation

In this paper, a *stochastic process* $X = \{X(t) : t \geq 0\}$ is a family of random variables $X(t) = X(t, \cdot)$ defined on a complete probability space $(\Omega, \mathcal{F}, \text{Pr})$. Here a random variable is a real valued measurable function, rather than an equivalence class of almost surely equal measurable functions. For each $\omega \in \Omega$, the function $X(\cdot) = X(\cdot, \omega)$ is called a *sample function* of X . We say that two stochastic processes X and Y are *indistinguishable*, and write $X = Y$, if almost surely, $X(t) = Y(t)$ for each t . Thus we identify two stochastic processes whose almost all sample functions are the same. This point of view implies that for two random variables ξ and η their equivalence means $\xi = \eta$ almost surely. We say that two stochastic processes X and Y are *modifications* of each other if for each t , $X(t) = Y(t)$ almost surely.

Regulated and càdlàg stochastic processes. A stochastic process $X = \{X(t): t \geq 0\}$ has regulated sample functions if for almost all $\omega \in \Omega$, there exist the limits

$$X(t-, \omega) := \lim_{u \uparrow t} X(u, \omega) \quad \text{and} \quad X(s+, \omega) := \lim_{u \downarrow s} X(u, \omega)$$

for $0 \leq s < t < \infty$. In this case we say that X is a *regulated stochastic process*. By Theorem 11.5 of Doob [20, p. 361], if a separable stochastic process X is a martingale then X is regulated. In Probability Theory sample functions of stochastic processes are often assumed to be regulated and right-continuous, that is, càdlàg. Let X be a regulated stochastic process and let Ω_X be the set of all $\omega \in \Omega$ such that $X(\cdot, \omega)$ is regulated. Due to completeness of the underlying probability space, $\Omega_X \in \mathcal{F}$ and $\Pr(\Omega_X) = 1$. For each $t \in [0, \infty)$, let

$$X_+(t) := X_+(t, \omega) := \begin{cases} X(t+, \omega) & \text{if } \omega \in \Omega_X \\ X(t, \omega) & \text{otherwise.} \end{cases}$$

Similarly define $X_-(t)$ for each $t \in (0, \infty)$ and let $X_-(0) := X(0)$. For each $t \in [0, \infty)$, $X_+(t)$ is a random variable because it is the limit for $\omega \in \Omega_X$ of the random variables $X(r_n)$ for r_n rational, $r_n \downarrow t$, and equals $X(t)$ otherwise. Likewise, $X(t-)$ is a random variable. Therefore $X_+ = \{X_+(t): t \geq 0\}$ and $X_- = \{X_-(t): t \geq 0\}$ are stochastic processes on the same probability space as X .

Recall that $\Delta^+ X(t) := X_+(t) - X(t)$ and $\Delta^- X(t) := X(t) - X_-(t)$. For each $t \in [0, \infty)$, let

$$\Omega_d(t) := \{\omega \in \Omega: \text{either } \Delta^- X(t) \neq 0 \text{ or } \Delta^+ X(t) \neq 0\}.$$

For a regulated stochastic process X , a point $t \in [0, \infty)$ is called a point of *fixed discontinuity* if $\Pr(\Omega_d(t)) > 0$. If t is not a point of fixed discontinuity then

$$\lim_{s \rightarrow t} X(s) = X(t) \quad \text{almost surely.} \tag{4.1}$$

Indeed, for $t \in (0, \infty)$ and $\omega \in \Omega_X \setminus \Omega_d(t)$, we have

$$\lim_{s \uparrow t} X(s, \omega) = X(t-, \omega) = X(t, \omega) = X(t+, \omega) = \lim_{s \downarrow t} X(s, \omega).$$

Since $\Pr(\Omega_X \setminus \Omega_d(t)) = 1$, (4.1) holds when $t \in (0, \infty)$. The same argument yields (4.1) when $t = 0$. If the stochastic process X has no points of fixed discontinuity then the three processes X_- , X_+ and X are modifications of each other, that is

$$\Pr(\{X(t) = X(t-)\}) = \Pr(\{X(t) = X(t+)\}) = 1$$

for each $t \in [0, \infty)$.

A point $t \in [0, \infty)$ is a point of *stochastic continuity* of X if $X(s) \rightarrow X(t)$ in probability as $s \rightarrow t$.

Theorem 1 *Let X be a regulated stochastic process. The set of points of fixed discontinuity of X is at most countable and coincides with the set of points of stochastic discontinuity of X .*

Proof. Since the limit of a sequence convergent in probability is unique almost surely, one can show that X is not stochastically continuous at some $t \in [0, \infty)$ if t is a point of fixed discontinuity. Also, X is stochastically continuous at t whenever t is not a point of fixed

discontinuity. This yields the second part of the claim. The first part follows from Theorem 11.1, Ch. VII, of Doob [20]. $\square$

By the preceding theorem, the set of fixed discontinuities of any stochastic process with regulated sample functions is at most countable. However, almost every sample function of such a process may have a non-fixed discontinuity, e.g. $X(t, \omega) = 1_{\{t \geq \omega\}}$, where ω has a uniform distribution in $[0, 1]$.

The p -variation and its index. Suppose that almost all sample functions of a stochastic process X have bounded p -variation on $[0, t]$ for some $0 < p, t < \infty$. Since the set $\Xi[0, t]$ of all partitions of $[0, t]$ is uncountable, the function $\omega \mapsto v_p(X(\cdot, \omega); [0, t])$ need not be measurable. For example, let $\Pr$ be Lebesgue measure on $\Omega := [0, 1]$. For a non-Lebesgue measurable set $A \subset [0, 1]$, let $X = \{X(t): 0 \leq t \leq 1\}$ be a stochastic process defined by $X(t, \omega) = 0$ on $[0, 1]$ except at the point $t = \omega \in A$ when $X(\omega, \omega) = 1$. Then $v_p(X; [0, 1]) = 0$ if $\omega \notin A$, and $= 2$ if $\omega \in A$. Notice that all sample functions of X are regulated.

One possibility to overcome this difficulty is to understand the supremum

$$\sup \{s_p(X(\cdot, \omega); \kappa): \kappa \in \Xi[0, T]\}, \quad \omega \in \Omega,$$

as the lattice supremum in the space of equivalence classes of random variables $L_0 = L_0(\Omega, \mathcal{F}, \Pr)$. Recall that L_0 is a Dedekind complete Riesz space of countable type (see §IV.1 and §VI.2 in Vulikh [107]). Therefore every infinite set E in L_0 which is bounded above has a supremum (in L_0), and there exists a finite or countable subset $E' \subset E$ such that $\sup E' = \sup E$.

In Stochastic Analysis, traditionally a different route is used to deal with non-measurability problems. Next we show that for a càdlàg stochastic process, the p -variation over an interval is indistinguishable from the p -variation over a countable and everywhere dense set, which is always measurable. Let $\lambda = \{\lambda_m: m \geq 1\}$ be a sequence of nested partitions $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ of $[0, \infty)$ such that $\cup_m \lambda_m$ is dense in $[0, \infty)$, and let $\Lambda[0, \infty)$ be the set of all such sequences. For example, the sequence $\{i2^{-m}: i = 0, \dots, m2^m\}$, $m \geq 1$, belongs to $\Lambda[0, \infty)$. Let f be a function on $[0, \infty)$, and let $0 < p < \infty$. For each $m \geq 1$ and $t \geq 0$, let

$$v_p(f; \lambda_m)(t) := \max \left\{ \sum_{j=1}^k |f(s_j \wedge t) - f(s_{j-1} \wedge t)|^p: \{0, t_{n(m)}^m\} \subset \{s_j: j = 0, \dots, k\} \subset \lambda_m \right\},$$

which is the p -variation over the finite set $\{t_i^m \wedge t: i = 0, \dots, n(m)\}$, equal to the trace partition $\lambda_m \cap [0, t]$ when $t \leq t_{n(m)}^m$ (see Notation 3.2). Since λ_m , $m \geq 1$, are nested partitions, the sequence $v_p(f; \lambda_m)(t)$, $m \geq 1$, is nondecreasing for each $t \geq 0$. For $t \geq 0$, let

$$v_p(f)(t) := v_p(f; \lambda)(t) := \sup_{m \geq 1} v_p(f; \lambda_m)(t) = \lim_{m \rightarrow \infty} v_p(f; \lambda_m)(t). \quad (4.2)$$

For a stochastic process X and each $t \geq 0$, $v_p(X)(t, \omega) := v_p(X(\cdot, \omega))(t)$ is possibly unbounded but measurable function of $\omega \in \Omega$.

Theorem 2 *Let $X = \{X(t): t \geq 0\}$ be either a separable stochastic process continuous in probability, or a càdlàg stochastic process, let $\lambda \in \Lambda[0, \infty)$, and let $0 < p < \infty$. If for each integer $k \geq 1$,*

$$v_p(X; \lambda)(k) < \infty \quad \text{almost surely,} \quad (4.3)$$

then $\{v_p(X; [0, t]): t \geq 0\}$ is a stochastic process indistinguishable from $v_p(X)$. Moreover, if in addition $p \geq 1$ then almost surely for each $t > 0$,

$$v_p(X)(t)^{1/p} - v_p(X)(t-)^{1/p} \leq |\Delta^- X(t)| \leq \{v_p(X)(t) - v_p(X)(t-)\}^{1/p},$$

and for each $t \geq 0$,

$$v_p(X)(t+)^{1/p} - v_p(X)(t)^{1/p} \leq |\Delta^+ X(t)| \leq \{v_p(X)(t+) - v_p(X)(t)\}^{1/p}.$$

Proof. Let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[0, \infty)$. First, suppose that X is a separable stochastic process continuous in probability. Then by Theorem 2.2 of Doob [20, Section II.2], $S := \cup_m \lambda_m$ is a separating set. Thus there exists a Pr-null set Ω_0 such that if A is a closed subset of $\mathbb{R}$ and I is an open subset of $[0, \infty)$, then

$$\{\omega: X(t, \omega) \in A, t \in S \cap I\} \setminus \{\omega: X(t, \omega) \in A, t \in I\} \subset \Omega_0.$$

The following consequence of separability is what we need: for each $\omega \notin \Omega_0$ and any open interval $I \subset [0, \infty)$,

$$\overline{X(I, \omega)} = \overline{X(I \cap S, \omega)}. \quad (4.4)$$

Here $\overline{X(I, \omega)}$ denotes the closure in $\mathbb{R}$ of the set of values assumed by the sample function $X(\cdot, \omega)$ as $t \in I$, and the other set has similar meaning. In particular, for every $t \geq 0$ there exists a sequence $\{s_k: k \geq 1\} \subset S$ such that $s_k \rightarrow t$ and $X(s_k, \omega) \rightarrow X(t, \omega)$ for each $\omega \notin \Omega_0$ (the sequence $\{s_k\}$ may depend on ω). Second, if X is a càdlàg stochastic process then (4.4) holds for each càdlàg sample function $X(\cdot, \omega)$.

Let Ω_0 be a Pr-null set such that for each $\omega \notin \Omega_0$, (4.4) holds and $v_p(X; \lambda)(k, \omega) < \infty$ for each integer $k \geq 1$. Let $0 \leq r < t < \infty$. If $r \in S$ then

$$v_p(X; \lambda)(r, \omega) \leq v_p(X, \lambda)(t, \omega) < \infty \quad (4.5)$$

holds for each ω by definition (4.2). If $r \notin S$ then for each $\omega \notin \Omega_0$ and for each $m \geq 1$,

$$v_p(X; \lambda_m)(r, \omega) = \lim_{s_k \rightarrow r, s_k \in S} v_p(X; \lambda_m)(s_k, \omega) \leq v_p(X; \lambda)(t, \omega).$$

Therefore (4.5) holds for each $\omega \notin \Omega_0$ and any $0 \leq r < t < \infty$. Let $t > 0$ and let κ be a partition of $[0, t]$. By (4.4), for each $\omega \notin \Omega_0$, $s_p(X(\cdot, \omega); \kappa)$ can be approximated arbitrarily closely by sums $s_p(X(\cdot, \omega); \kappa_m)$ with $\{0, t\} \subset \kappa_m \subset \lambda_m \cap [0, t]$ as $m \rightarrow \infty$. Therefore for each $\omega \notin \Omega_0$,

$$v_p(X(\cdot, \omega); [0, t]) = v_p(X; \lambda)(t, \omega) < \infty,$$

proving the first part of the theorem. The second part of the theorem follows from the inequalities

$$v_p(f; [a, c]) + v_p(f; [c, b]) \leq v_p(f; [a, b]) \leq \{v_p(f; [a, c])^{1/p} + v_p(f; [c, b])^{1/p}\}^p$$

valid for $a < c < b$ (Lemma 4.6 in [24]), and from the relations,

$$\lim_{z \uparrow x} v_p(f; [z, x]) = |\Delta^- f(x)|^p \quad \text{and} \quad \lim_{z \downarrow y} v_p(f; [y, z]) = |\Delta^+ f(y)|^p$$

valid for $a \leq y < x \leq b$ (Lemma 2.19 in [23, Part II]). $\square$

For a càdlàg stochastic process X if (4.3) holds for some $\lambda \in \Lambda[0, \infty)$ then it also holds for any other such λ , and $v_p(X)$ is a càdlàg stochastic process. It is clear from definition (4.2) that if X is adapted then $v_p(X)$ is also adapted stochastic process whose almost all sample functions are nondecreasing.

Definition 3 Let $0 < p < \infty$. We say that a stochastic process X is *locally of bounded p -variation* if X is adapted, càdlàg and (4.3) holds for some $\lambda \in \Lambda[0, \infty)$ and each integer $k \geq 1$. Also, $v_p(X)$ is called the *p -variation process* for X .

For a function $f: [0, T] \mapsto \mathbb{R}$, the p -variation index of f is defined by

$$v(f) := v(f; [0, T]) := \begin{cases} \inf\{p > 0: v_p(f; [0, T]) < \infty\} & \text{if the set is nonempty,} \\ +\infty & \text{otherwise.} \end{cases} \quad (4.6)$$

For a stochastic process X on a complete probability space $(\Omega, \mathcal{F}, \Pr)$, the p -variation index $v(X; [0, T])$ is defined provided $v(X(\cdot, \omega); [0, T])$ is a constant for almost all $\omega \in \Omega$. We say that the *local p -variation index* $v(X) = p$ if with probability 1, the p -variation index $v(X; [0, T]) = p$ for each $T > 0$.

Local martingales. We continue with recalling a standard framework of Stochastic Analysis. Let $(\Omega, \mathcal{F}, \Pr)$ be a complete probability space, and let $\mathbb{F} = \{\mathcal{F}_t: 0 \leq t < \infty\}$ be a filtration of sub- σ -algebras of $\mathcal{F}$ satisfying the usual hypotheses, that is, $\mathbb{F}$ is right-continuous and $\mathcal{F}_0$ contains all the $\Pr$ -null sets of $\mathcal{F}$. A stochastic process $X = \{X(t): t \geq 0\}$ on $(\Omega, \mathcal{F}, \Pr)$ is said to be *adapted* if $X(t)$ is $\mathcal{F}_t$ measurable for each t . An adapted, càdlàg stochastic process X is called a *local martingale* with respect to the filtration $\mathbb{F}$ if there exists a sequence of increasing $\mathbb{F}$ -stopping times, τ_n , with $\lim_{n \rightarrow \infty} \tau_n = +\infty$ a.s. such that $X^{\tau_n} 1_{\{\tau_n > 0\}} = \{X(t \wedge \tau_n) 1_{\{\tau_n > 0\}}: t \geq 0\}$ is a uniformly integrable martingale for each n . This definition of a local martingale is used in the book by Protter [95, p. 33], which is used for references in this chapter.

The following result is due to Lépingle [62]. A different proof of this result was given by Pisier and Xu [92].

Theorem 4 *A local martingale is locally of bounded p -variation for each $p > 2$. Moreover, for $p > 2$ and $1 \leq r < \infty$ there is a finite constant $K_{p,r}$ such that for any martingale $M = \{M(t): 0 \leq t \leq T\}$, $T > 0$, in $L^r(\Omega, \mathcal{F}, \Pr)$,*

$$E[v_p(M; [0, T])]^{r/p} \leq K_{p,r} E\left[\sup_{0 \leq t \leq T} |M(t)|\right]^r. \quad (4.7)$$

Proof. Let $p > 2$, $1 \leq r < \infty$ and let $M = \{M(t): 0 \leq t \leq T\}$ be a martingale in L^r . Let $\lambda = \{\lambda_m: m \geq 1\}$ be a nested sequence of partitions $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ of $[0, T]$ such that $\cup_m \lambda_m$ is dense in $[0, T]$. By Theorem 2.4 of Pisier and Xu [92], there is finite constant $K_{p,r}$ such that for each $m \geq 1$,

$$E[s_p(M; \lambda_m)(T)]^{r/p} \leq K_{p,r} E\left[\max_{0 \leq i \leq n(m)} |M(t_i^m)|\right]^r \leq K_{p,r} E\left[\sup_{0 \leq t \leq T} |M(t)|\right]^r.$$

Thus (4.7) follows by Fatou lemma and Theorem 2. Suppose now that M is a local martingale. By the Fundamental Theorem of Local Martingales (Theorem III.13 in [95]), $M = N + A$, where N is locally bounded martingale and A is locally of bounded 1-variation. Thus an application of the first part of the proof yields the conclusion of the theorem. $\square$

It is well known that sample functions of martingales which are continuous and nonconstant must have unbounded 1-variation (see Lemma 3.2.1 of Fisk [33]). In fact they must have unbounded p -variation for every $p < 2$.

Theorem 5 *If for some $p < 2$ a sample continuous local martingale is locally of bounded p -variation then almost surely its sample functions are constants.*

The proof of this statement is based on a stopping time technique as follows. We will say that a stochastic process X is sample uniformly continuous if for each $0 < T < \infty$ there exists a Pr-null set Ω_0 such that given $\epsilon > 0$ one can choose $\delta > 0$ such that $|X(t, \omega) - X(s, \omega)| < \epsilon$ whenever $|t - s| < \delta$, $t, s \in [0, T]$ and $\omega \notin \Omega_0$. For a stopping time τ , let $X^\tau := \{X(t \wedge \tau): t \geq 0\}$.

Lemma 6 *Let X be a sample continuous stochastic process adapted to a filtration $\mathbb{F}$. For each $0 < T < \infty$ there exists a sequence $\{\tau_k: k \geq 1\}$ of increasing $\mathbb{F}$ stopping times such that $\Pr(\cup_{k=1}^\infty \{\tau_k = T\}) = 1$, and each X^{τ_k} is sample uniformly continuous and bounded by k .*

Proof. Let $0 < T < \infty$. Since almost every sample function of X is uniformly continuous on $[0, T]$, there exists a sequence $\{\delta_{kn}: k, n \geq 1\}$ of positive numbers such that, for each k , $\delta_{k1} > \delta_{k2} > \dots$, and for each k, n ,

$$\Pr(\{\sup\{|X(t) - X(s)|: |t - s| \leq \delta_{kn}\} \geq 1/k\}) \leq 2^{-n-k}. \quad (4.8)$$

Let τ'_{kn} be the least $t \in [0, T]$ such that

$$\sup\{|X(s'') - X(s')|: |s'' - s'| \leq \delta_{kn}, s'', s' \in [0, t]\} \geq 1/k.$$

If there is no such t , let $\tau'_{kn} := T$. Since X is sample continuous each τ'_{kn} is an $\mathbb{F}$ stopping time. Let $\tau'_k := \inf_n \tau'_{kn}$. Then for each k and any real r , $\{\tau'_k < r\} = \cup_n \{\tau'_{kn} < r\} \in \mathbb{F}_r$. Since $\mathbb{F}$ is right continuous, each τ'_k is an $\mathbb{F}$ stopping time. Then each stopped process $X^{\tau'_k}$ is sample uniformly continuous. Also, by (4.8), $\Pr(\{X^{\tau'_k} \neq X\}) \leq 2^{-k}$. It then follows from the Borell-Cantelli lemma that $\Pr(\cup_k \{\tau'_k = T\}) = 1$. Let τ''_k be the least $t \in [0, T]$ such that $|X(t)| > k$. If no such t exists, let $\tau''_k := T$. Then each τ''_k is an $\mathbb{F}$ stopping time and $X^{\tau''_k}$ is bounded by k . Clearly the sequence of $\mathbb{F}$ stopping times $\tau_k := \tau'_k \wedge \tau''_k$, $k \geq 1$, satisfies the conclusion of the lemma. $\square$

Now we are ready to prove Theorem 5.

Proof of Theorem 5. Let a sample continuous local martingale X be locally of bounded p -variation for some $1 \leq p < 2$. It is enough to prove that for any $0 < T < \infty$, almost surely, $X(t) = X(0)$ for each $0 \leq t \leq T$. Since X is sample continuous, the p -variation process $v_p(X)$ is also sample continuous by Theorem 2. By Lemma 6 applied to X and $v_p(X)$, and since X is a local martingale, there exists a sequence $\{\tau_k: k \geq 1\}$ of $\mathbb{F}$ stopping times such that $\Pr(\cup_k \{\tau_k = T\}) = 1$, each X^{τ_k} is sample uniformly continuous martingale bounded by k and each $v_p(X)^{\tau_k}$ is bounded by k . Notice that each $v_p(X)^{\tau_k}$ is the p -variation process for X^{τ_k} . Let $\lambda = \{\lambda_m: m \geq 1\}$ be a sequence of nested partitions $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ of $[0, T]$ such that $\max_i(t_i^m - t_{i-1}^m) \rightarrow 0$ as $m \rightarrow \infty$. Then $\cup_m \lambda_m$ is dense in $[0, T]$. For each $k, m \geq 1$, let

$$\epsilon_{k,m} := \text{esssup}_\omega \max_i |X^{\tau_k}(t_i^m) - X^{\tau_k}(t_{i-1}^m)|.$$

Let $t \in \cup_m \lambda_m$. Then there exists an integer m_t such that $t \in \lambda_m$ for each $m \geq m_t$. Since X^{τ_k} has orthogonal increments, for each $m \geq m_t$, we have

$$E[X^{\tau_k}(t) - X^{\tau_k}(0)]^2 = Es_2(X^{\tau_k}; \lambda_m) \leq E(\epsilon_{k,m}^{2-p} v_p(X)^{\tau_k}(t)) \leq kE(\epsilon_{k,m}^{2-p}).$$

Since for each $k \geq 1$, $\epsilon_{k,m} \leq 2k$ and $\epsilon_{k,m} \rightarrow 0$ almost surely as $m \rightarrow \infty$, by the dominated convergence theorem, the left side of the preceding inequality is zero for each $k \geq 1$. Therefore,

almost surely, for each $k \geq 1$, $X^{\tau_k}(t) = X^{\tau_k}(0)$ for each $t \in \cup_m \lambda_m$. Since X is sample continuous and since for almost every $\omega \in \Omega$, there is an integer $k \geq 1$ such that $X^{\tau_k}(t, \omega) = X(t, \omega)$ for $0 \leq t \leq T$, it follows that almost surely, $X(t) = X(0)$ for each $0 \leq t \leq T$. The proof of Theorem 5 is complete. $\square$

4.2 Stochastic integral and the Left Young integral

We show here that values of the Left Young integral for pairs of sample functions of stochastic processes and values of the corresponding stochastic integral agree almost surely under conditions ensuring the existence of both. Let Y, X be two regulated stochastic processes such that for almost all $\omega \in \Omega$, the Left Young integral

$$(LY) \int_0^t Y(\cdot, \omega) dX(\cdot, \omega)$$

exists for all $t > 0$. For the rest of this chapter we consider an integration with respect to a stochastic process X which in addition is right-continuous. In that case, by Definition 2.21 of the Left Young integral, for $0 \leq s < t < \infty$,

$$(LY) \int_s^t Y dX = (RRS) \int_s^t Y_-^{(s)} dX = (RRS) \int_s^t Y_-^{(0)} dX.$$

The second equality holds also due to the right-continuity of X . We skip the superscript (0) in what follows for notation simplicity, and always assume that $Y(0-) = Y(0)$. So that the Left Young integration of Y with respect to a càdlàg stochastic process X reduces to the refinement Riemann-Stieltjes integration of the left modification Y_- with respect to X . Then we define the *integral stochastic process* by

$$\int Y_- dX := \left\{ (RRS) \int_0^t Y_- dX : t \geq 0 \right\}. \quad (4.9)$$

Also for $1 \leq p < \infty$, we say that a function f on $[0, \infty)$ belongs to $\text{dual}(\mathcal{W}_p)^{\text{loc}}$ if for each $0 < T < \infty$, f belongs to $\text{dual}(\mathcal{W}_p)[0, T]$ defined by (1.16).

Proposition 7 *Let $1 \leq p < \infty$, let M be a local martingale of locally bounded p -variation, and let H be an adapted stochastic process with almost all sample functions in $\text{dual}(\mathcal{W}_p)^{\text{loc}}$. Then the stochastic integral $H_- \cdot M$ and the integral stochastic process $\int H_- dM$ are defined and indistinguishable.*

Proof. The integral stochastic process $\int H_- dM$ exists by Theorem 2.22. The stochastic integral $H_- \cdot M$ exists by Theorem III.17 in [95, p. 106]. We have to prove that for almost all $\omega \in \Omega$, the equality

$$(RRS) \int_0^t H(s-, \omega) dM(s, \omega) = (H_- \cdot M)(t, \omega) \quad (4.10)$$

holds for all $t \geq 0$. There exists $\Omega_0 \subset \Omega$ such that $\Pr(\Omega_0) = 1$ and for all $\omega \in \Omega_0$ the left and right sides of (4.10) are regulated right-continuous functions of t . Therefore it is enough to prove that (4.10) holds for an arbitrary t and for almost all ω . Fix $t > 0$. To prove (4.10) we use term by term integration theorems for the RRS integral and the dominated convergence in probability for the stochastic integral.

For each integer $m \geq 1$, let $\lambda_m := \{u_j^m := tj2^{-m} : j = 0, \dots, 2^m\}$ be a partition of $[0, t]$. Let $H_m(s) := H(u_{j-1}^m)$ if $s \in [u_{j-1}^m, u_j^m)$ for some $j = 1, \dots, 2^m$ and $H_m(t) := H(t)$. We claim that for each $m \geq 1$, for each right-continuous regulated sample function of M , and for each bounded sample function of H , we have

$$(RRS) \int_0^t H_m(s-) dM(s) = \sum_{j=1}^{2^m} H(u_{j-1}^m) [M(u_j^m) - M(u_{j-1}^m)] = S_{LC}(H, M; \lambda_m), \quad (4.11)$$

where $H_m(0-) = H_m(0) := 0$. Indeed, let $m \geq 1$, and let $\kappa = \{([t_{i-1}, t_i], s_i) : i = 1, \dots, n\}$ be a tagged partition of $[0, t]$ and a refinement of λ_m . Therefore for each $j = 0, \dots, 2^m$, there exists an index $i(j) \in \{0, \dots, n\}$ such that $t_{i(j)} = u_j^m$. Then the Riemann-Stieltjes sum based on κ is given by

$$\begin{aligned} S_{RS}((H_m)_-, M; \kappa) &= \sum_{i=1}^n H_m(s_i-) [M(t_i) - M(t_{i-1})] \\ &= \sum_{j=1}^{2^m} \left\{ H_m(s_{i(j-1)+1}-) [M(t_{i(j-1)+1}) - M(u_{j-1}^m)] + H(u_{j-1}^m) [M(u_j^m) - M(t_{i(j-1)+1})] \right\}. \end{aligned}$$

The right side of the preceding display approaches the right side of (4.11) as $t_{i(j-1)+1} \downarrow u_{j-1}^m$ for each $j = 1, \dots, 2^m$ because sample functions of M are right-continuous and sample functions of H are bounded. Therefore given $\epsilon > 0$, one can choose a partition λ of $[0, t]$ containing all points of λ_m and points $v_j \in (u_{j-1}^m, u_j^m)$, $j = 1, \dots, 2^m$ sufficiently close to u_{j-1}^m , so that the Riemann-Stieltjes sums based on refinements of λ are within ϵ to the right side of (4.11). Thus the RRS integral in (4.11) exists and has the stated value.

Next we prove that

$$\lim_{m \rightarrow \infty} (RRS) \int_0^t H_m(s-) dM(s) = (RRS) \int_0^t H(s-) dM(s). \quad (4.12)$$

We have that almost every sample function of M is right-continuous and when restricted to $[0, t]$ is in $\mathcal{W}_p[0, t]$, while almost every sample function of H when restricted to $[0, t]$ is in $\text{dual}(\mathcal{W}_p)[0, t]$. First suppose that $p > 1$. In this case we use Theorem 5.6 on term by term integration of L. C. Young [115, p. 602]. Let $q < \infty$ be such that the sample function of H is in $\mathcal{W}_q[0, t]$ and $p^{-1} + q^{-1} > 1$. It is clear that the q -variation $v_q(H_m; [0, t]) \leq v_q(H; [0, t])$ for each $m \geq 1$. The other assumptions of L. C. Young's theorem will follow once we show that $(H_m)_-$ converge to H_- uniformly on the left at each point of $(0, t]$, that is, for $s \in (0, t]$ and given $\epsilon > 0$ there is an integer M and a $\delta > 0$ such that for all $m \geq M$ and all $x \in [s - \delta, s)$, $|H_m(x-) - H(x-)| < \epsilon$. Let $s \in (0, t]$ and let $\epsilon > 0$. Choose a $\delta > 0$ such that $|H(x) - H(s-)| < \epsilon/2$ for each $x \in [s - \delta, s)$. Let M be the minimal positive integer such that $t2^{-M} \leq \delta/2$. Then for all $x \in [s - \delta/2, s)$ and all $m \geq M$, $|H_m(x-) - H(x-)| \leq \epsilon/2 + |H(s-) - H(u_{i-1}^m)| < \epsilon$ for some $u_{i-1}^m \in [s - \delta, s)$. Thus all assumptions of L. C. Young's theorem are satisfied, and hence (4.12) holds in the case $p > 1$. Now suppose that $p = 1$. In this case we use the Osgood convergence theorem proved for the Young-Stieltjes integrals by Hildebrandt [49, Theorem 19.3.14]. Since $\|(H_m)_-\|_{\sup} \leq \|H\|_{\sup} < \infty$ and $(H_m)_-$ converge pointwise to H_- , (4.12) holds by the Osgood theorem in the case $p = 1$. Therefore (4.12) holds. On the other hand, by Theorem VIII.15 of Dellacherie and Meyer [16, p. 324], the limit

$$\lim_{m \rightarrow \infty} S_{LC}(H, M; \lambda_m) = (Y_- \cdot M)(t)$$

exists in probability. This in conjunction with (4.11) and (4.12) yields that (4.10) holds for almost all $\omega \in \Omega$. The proof of Proposition 7 is complete. $\square$

4.3 $(2 - \epsilon)$ -semimartingales and the stochastic $\ominus$ -integral

In this section we define the class of $(2 - \epsilon)$ -semimartingales and the stochastic $\ominus$ -integral with respect to a $(2 - \epsilon)$ -semimartingale.

Definition 8 Let $1 \leq p < 2$, and let X be an adapted càdlàg stochastic process. We say that X is a p -semimartingale if there exist stochastic processes M and A such that

$$X - X(0) = M + A \quad \text{almost surely,} \quad (4.13)$$

$M(0) = A(0) = 0$, M is a local martingale and A is a stochastic process of locally bounded p -variation. The set of all such pairs (M, A) is denoted by $\mathcal{D}_p(X)$, and $\mathcal{D}(X) := \cup\{\mathcal{D}_p(X) : 1 \leq p < 2\}$. A stochastic process X is called a $(2 - \epsilon)$ -semimartingale if X is a p -semimartingale for some $1 \leq p < 2$, that is, if the set $\mathcal{D}(X)$ is nonempty.

The 1-semimartingale is the same as semimartingale. As in the case of semimartingales, decomposition (4.13) may not be unique. For example, the Lévy decomposition can be viewed in two different ways: either as a 1-semimartingale (see Theorem I.40 in [95, p. 31]), or as a p -semimartingale with the decomposition into a Brownian motion part and a pure jump part. However we prove that in most interesting cases a "natural" integral with respect to a p -semimartingale and a continuous part of the quadratic variation of a p -semimartingale are unique in spite of a possible non-uniqueness of decomposition (4.13).

In the definition of a $(2 - \epsilon)$ -semimartingale we require the second component A in decomposition (4.13) to have locally finite p -variation for some $1 \leq p < 2$. Instead, one may define a class of stochastic processes with decomposition (4.13), where A has almost all sample functions locally in $\mathcal{W}_2^*$. Stochastic processes with such a property are reminiscent to Dirichlet stochastic processes defined by Fölmer [35], and further modified in different directions by other authors.

Let X be a $(2 - \epsilon)$ -semimartingale, and let H be an adapted regulated stochastic process. Let $(M, A) \in \mathcal{D}(X)$. The stochastic integral $H_- \cdot M = \{(H_- \cdot M)(t) : t \geq 0\}$ of the left-continuous modification H_- with respect to a local martingale M is well-known. We can and do assume that $H_- \cdot M$ is a càdlàg stochastic process. It is natural to define a stochastic integral of H with respect to a $(2 - \epsilon)$ -semimartingale X by

$$(H \ominus X)(t) := (H_- \cdot M)(t) + (RRS) \int_0^t H_- dA, \quad t \geq 0, \quad (4.14)$$

provided the integral stochastic process $\int H_- dA$ is defined (see (4.9)). If A is a stochastic process of locally bounded variation then the refinement Riemann-Stieltjes integral in (4.14) is defined and has the same value as the Lebesgue-Stieltjes integral. If A is a stochastic process of locally bounded p -variation with $1 < p < 2$ then the refinement Riemann-Stieltjes integral in (4.14) is defined provided H is a stochastic process of locally bounded q -variation with $p^{-1} + q^{-1} > 1$ by the L. C. Young Theorem on Stieltjes integrability (see (1.6)). In particular, the refinement Riemann-Stieltjes integral in (4.14) is defined provided H is a local martingale by Theorem 4. However, since decomposition (4.13) may not be unique a value of sum (4.14) also need not be the same for different decompositions of a $(2 - \epsilon)$ -semimartingale X . We define the stochastic

$\ominus$ -integral with respect to a $(2 - \epsilon)$ -semimartingale provided a sum of the two integrals in (4.14) does not depend on decomposition (4.13).

Definition 9 Let X be a $(2 - \epsilon)$ -semimartingale, and let H be an adapted regulated stochastic process such that the integral stochastic process $\int H_- dA$ is defined for each $(M, A) \in \mathcal{D}(X)$. If the stochastic processes

$$\left\{ (H_- \cdot M)(t) + (RRS) \int_0^t H_- dA : t \geq 0 \right\} \quad (4.15)$$

are indistinguishable for different pairs $(M, A) \in \mathcal{D}(X)$ then define the *stochastic $\ominus$ -integral* $H \ominus X = \{(H \ominus X)(t) : t \geq 0\}$ by (4.14).

The next statement follows from the bilinearity of the stochastic integral and of the refinement Riemann-Stieltjes integral.

Proposition 10 Let X, Y be $(2 - \epsilon)$ -semimartingales and let H, G be adapted regulated stochastic processes. Then

$$(H + G) \ominus (X + Y) = H \ominus X + H \ominus Y + G \ominus X + G \ominus Y$$

provided all five stochastic $\ominus$ -integrals are defined.

For a stochastic process $Y = \{Y(t) : t \geq 0\}$, the *jump process* $\Delta^- Y = \{\Delta^- Y(t) : t \geq 0\}$ is defined by $\Delta^- Y(t) = Y(t) - Y(t-)$ if $t > 0$, and $\Delta^- Y(0) := 0$.

Proposition 11 Let X be a $(2 - \epsilon)$ -semimartingale, and let H be an adapted regulated stochastic process such that the stochastic $\ominus$ -integral $H \ominus M$ is defined. Then almost all sample functions of $H \ominus X$ are càdlàg and

$$\{\Delta^- (H \ominus X)(t) : t \geq 0\} = \{H_-(t) \Delta^- X(t) : t \geq 0\}.$$

Proof. By the same property for the stochastic integral with respect to a local martingale (Theorem II.13 in [95, p. 53]), and by Proposition 2.27, we have

$$\Delta^- (H \ominus X) = \Delta^- (H_- \cdot M) + \Delta^- (\int H_- dA) = H_- \Delta^- M + H_- \Delta^- A = H_- \Delta^- X$$

almost surely, and the proof is complete. $\square$

Next is a sufficient condition for the stochastic $\ominus$ -integrability with respect to a $(2 - \epsilon)$ -semimartingale.

Theorem 12 Let X be a $(2 - \epsilon)$ -semimartingale, and let H be an adapted stochastic process with the local p -variation index $v(H) = 2$. Then the stochastic $\ominus$ -integral $H \ominus X$ is defined, is a $(2 - \epsilon)$ -semimartingale, and for each $t > 0$ and $\lambda = \{\lambda_m : m \geq 1\} \in \Lambda[0, t]$,

$$\lim_{m \rightarrow \infty} S_{LC}(H, X; \lambda_m) = (H \ominus X)(t) \quad \text{in probability.} \quad (4.16)$$

Proof. Let $(M, A) \in \mathcal{D}(X)$. Then M is a local martingale, A is an adapted càdlàg stochastic process of locally bounded p -variation for some $1 \leq p < 2$, $M(0) = A(0) = 0$, and the decomposition (4.13) holds. By Theorem III.17 in [95, p. 106], the stochastic integral $H_- \cdot M$ is defined and is a local martingale. Since $1 \leq p < 2$ and the local p -variation index $v(H) = 2$, there exists $2 < q < \infty$ such that $p^{-1} + q^{-1} > 1$ and H has almost all sample functions in $\mathcal{W}_q^{\text{loc}}$. Thus by Theorem 2.22, there exists the integral stochastic process $\int H_- dA$. Hence stochastic process (4.15) is defined for (M, A) . We have to show that this process is unique up to indistinguishability for different elements of $\mathcal{D}(X)$. If the set $\mathcal{D}(X)$ has a single element then there is nothing to prove. Suppose that there are two different elements in $\mathcal{D}(X)$, that is, $(M_1, A_1) \in \mathcal{D}_{p_1}(X)$ and $(M_2, A_2) \in \mathcal{D}_{p_2}(X)$ for some $p_1, p_2 \in [1, 2)$. Taking the maximum between p_1 and p_2 if they are different, we can and do assume that $p_1 = p_2 = p$. By the decomposability property of X , we have that stochastic process $M_1 - M_2 = -(A_1 - A_2)$ has almost all sample functions in $\mathcal{W}_p^{\text{loc}}$. On the other hand, there exists $2 < q < \infty$ such that $p^{-1} + q^{-1} > 1$ and H has almost all sample functions in $\mathcal{W}_q^{\text{loc}}$, so that the refinement Riemann-Stieltjes integral

$$(RRS) \int_0^t H_-(\cdot, \omega) d(M_1 - M_2)(\cdot, \omega)$$

exists for almost all $\omega \in \Omega$ and for all $t > 0$. Thus by Proposition 7, for almost all $\omega \in \Omega$ and for all $t > 0$,

$$\begin{aligned} (H_- \cdot M_1)(t, \omega) - (H_- \cdot M_2)(t, \omega) &= (RRS) \int_0^t H_-(\cdot, \omega) d(M_1 - M_2)(\cdot, \omega) \\ &= (RRS) \int_0^t H_-(\cdot, \omega) dA_2(\cdot, \omega) - (RRS) \int_0^t H_-(\cdot, \omega) dA_1(\cdot, \omega). \end{aligned}$$

Therefore the stochastic processes (4.15) corresponding to (M_1, A_1) and (M_2, A_2) are indistinguishable, and so the stochastic $\ominus$ -integral $H \ominus X$ is defined.

By Theorem 2.22 and Proposition 2.27, the integral stochastic process $\int H_- dA$ is an adapted càdlàg stochastic process. By Corollary 2.13, $\int H_- dA$ has the same p -variation property as A . Since $H_- \cdot M$ is a local martingale, it follows that the stochastic $\ominus$ -integral $H \ominus X = H_- \cdot M + \int H_- dA$ is a $(2 - \epsilon)$ -semimartingale with value 0 at 0. The approximation in probability of the stochastic $\ominus$ -integral by Left Cauchy sums follows from Theorem 2.22 and from Theorem II.21 in [95, p. 57]. The proof of Theorem 12 is complete. $\square$

The preceding theorem shows that the property of being a $(2 - \epsilon)$ -semimartingale is preserved by the stochastic $\ominus$ -integration. Also, it provides a class of stochastic $\ominus$ -integrable processes which is small as compared with the class of all predictable stochastic processes integrable in the sense of the stochastic integral with respect to a semimartingale. However, the class of possible integrators allowed by Theorem 12 is larger.

In Stochastic Analysis the property proved next is called associativity of the stochastic integral.

Theorem 13 *Let X be a $(2 - \epsilon)$ -semimartingale, and let G, H be adapted stochastic processes with the local p -variation indices $v(G) = v(H) = 2$. Then the stochastic $\ominus$ -integral $H \ominus X$ is a $(2 - \epsilon)$ -semimartingale and*

$$G \ominus (H \ominus X) = (GH) \ominus X. \quad (4.17)$$

Proof. The stochastic $\ominus$ -integral $H \ominus X$ is defined and is a $(2 - \epsilon)$ -semimartingale by Theorem 12. Let $(M, A) \in \mathcal{D}(X)$. Then $(H_- \cdot M, \int H_- dA) \in \mathcal{D}(H \ominus X)$ by Theorem III.17 in [95, p. 106] and by Corollary 2.13. By the associativity of the stochastic integral with respect to a local martingale (Theorem II.19 in [95, p. 55]) and by Proposition 2.28, for any $0 < t < \infty$, the equalities

$$\begin{aligned} (G \ominus (H \ominus X))(t) &= (G_- \cdot (H_- \cdot M))(t) + (MPS) \int_0^t G_- d\left(\int H_- dA\right) \\ &= ((GH)_- \cdot M)(t) + (MPS) \int_0^t (GH)_- dA = ((GH) \ominus X)(t) \end{aligned}$$

hold almost surely. Since almost all sample functions of the stochastic integrals are càdlàg, the relation (4.17) follows. $\square$

4.4 Quadratic covariation of $(2 - \epsilon)$ -semimartingales

Let X, Y be two $(2 - \epsilon)$ -semimartingales. Since the local p -variation index of a $(2 - \epsilon)$ -semimartingale is equal to 2 by Theorem 4, the stochastic $\ominus$ -integrals $X \ominus Y$ and $Y \ominus X$ are defined by Theorem 12. Therefore the stochastic process $[X, Y] = \{[X, Y](t) : t \geq 0\}$ defined by

$$[X, Y](t) := X(t)Y(t) - X(0)Y(0) - (X \ominus Y)(t) - (Y \ominus X)(t), \quad t \geq 0, \quad (4.18)$$

exists and is called the *quadratic covariation* of the pair X, Y . In the case X and Y are semimartingales, definition (4.18) differ from the definition of the quadratic covariation in Section II.6 of [95, p. 58] by the random variable $X(0)Y(0)$. Let $\bar{X} := X - X(0)$ and $\bar{Y} := Y - Y(0)$. By (4.16), we have $X \ominus \bar{Y} = X \ominus Y$ and $Y \ominus \bar{X} = Y \ominus X$. Therefore it follows that

$$[X, Y] = \bar{X}\bar{Y} - \bar{X} \ominus \bar{Y} - \bar{Y} \ominus \bar{X}. \quad (4.19)$$

The above notion of the quadratic covariation for $(2 - \epsilon)$ -semimartingales is consistent with the quadratic λ -covariation for functions (Definition 3.12) in the following sense.

Proposition 14 *Let X, Y be $(2 - \epsilon)$ -semimartingales such that for some $\lambda \in \Lambda[0, T]$, $0 < T < \infty$, the 2-vector function $(X(\cdot, \omega), Y(\cdot, \omega))$ has the quadratic λ -covariation for almost all $\omega \in \Omega$. Then for almost all $\omega \in \Omega$,*

$$[X, Y](t, \omega) = [X(\cdot, \omega), Y(\cdot, \omega)]_\lambda(t), \quad 0 \leq t \leq T. \quad (4.20)$$

Proof. Let $\lambda = \{\lambda_m : m \geq 1\} \in \Lambda[0, T]$ be the sequence of partitions $\lambda_m = \{t_i^m : i = 0, \dots, n(m)\}$ of $[0, T]$ satisfying the assumptions of the proposition. By (4.16), we have that for each $0 \leq t \leq T$,

$$\begin{aligned} [X, Y](t) &= \lim_{m \rightarrow \infty} \sum_{i=1}^{n(m)} [X(t_i^m \wedge t) - X(t_{i-1}^m \wedge t)][Y(t_i^m \wedge t) - Y(t_{i-1}^m \wedge t)] \\ &= \lim_{m \rightarrow \infty} C(X, Y; \lambda_m \cap [0, t]) \quad \text{in probability.} \end{aligned}$$

It then follows that there exists a countable dense subset D of $[0, T]$, and a subset $\Omega_0 \subset \Omega$ such that $\Pr(\Omega_0) = 1$ and for each $\omega \in \Omega_0$, the functions $[X, Y](\cdot, \omega)$ and $[X(\cdot, \omega), Y(\cdot, \omega)]_\lambda$ on $[0, T]$ are càdlàg and

$$[X, Y](t, \omega) = [X(\cdot, \omega), Y(\cdot, \omega)]_\lambda(t), \quad t \in D.$$

Therefore (4.20) holds. $\square$

Let X, Y be $(2 - \epsilon)$ -semimartingales, and let $(M, A) \in \mathcal{D}(X)$, $(N, B) \in \mathcal{D}(Y)$. By (4.19) and by Proposition 10, we have the equality

$$[X, Y] = [M + A, N + B] = [M, N] + [M, B] + [A, N] + [A, B].$$

Since $M(0) = N(0) = 0$, by definition (4.18), the equality

$$[M, N] = MN - M_- \cdot N - N_- \cdot M$$

holds and shows that $[M, N]$ agree with the quadratic covariation for semimartingales used in Stochastic Analysis, and hence inherits standard properties of the quadratic covariation. For example, almost all sample functions of $[M, N]$ are càdlàg, have locally bounded variation, and $\Delta^- [M, N] = \Delta^- M \Delta^- N$. The same is true for the quadratic covariations $[M, B]$, $[A, N]$ and $[A, B]$. This follows from the preceding proposition in conjunction with Proposition 3.14. Also the last three quadratic covariation stochastic processes have pure jump sample functions. Since almost all sample functions of $[X, Y]$ are càdlàg, have locally bounded variation and $\Delta^- [X, Y] = \Delta^- X \Delta^- Y$, almost all sample functions of the stochastic process

$$[X, Y]^c := \left\{ [X, Y](t) - \sum_{(0, t]} \Delta^- X \Delta^- Y : t \geq 0 \right\}$$

are continuous. The following shows that the continuous part of the quadratic covariation of $(2 - \epsilon)$ -semimartingales agree with the analogous Stochastic Analysis construction and does not depend on their decompositions into local martingale and bounded p -variation parts.

Proposition 15 *Let X, Y be $(2 - \epsilon)$ -semimartingales. Then for each $(M, A) \in \mathcal{D}(X)$ and $(N, B) \in \mathcal{D}(Y)$, $[M, N]^c$ is indistinguishable from $[X, Y]^c$.*

Proof. Let $(M, A) \in \mathcal{D}(X)$ and $(N, B) \in \mathcal{D}(Y)$. Since $M(0) = N(0) = A(0) = B(0) = 0$, by (4.19) and Proposition 10, we have for each $t > 0$,

$$\begin{aligned} [X, Y](t) &= (M + A)(t)(N + B)(t) - ((M + A) \ominus (N + B))(t) - ((N + B) \ominus (M + A))(t) \\ &= [M, N](t) + \left\{ M(t)B(t) - (B_- \cdot M)(t) - (RRS) \int_0^t M_- dB \right\} \\ &\quad + \left\{ A(t)N(t) - (A_- \cdot N)(t) - (RRS) \int_0^t N_- dA \right\} \\ &\quad + \left\{ A(t)B(t) - (RRS) \int_0^t A_- dB - (RRS) \int_0^t B_- dA \right\}. \end{aligned}$$

Let $A = A(\cdot, \omega)$ and $B = B(\cdot, \omega)$ be càdlàg sample functions which have locally bounded p -variation for some $1 \leq p < 2$. For any $t > 0$ and any $\lambda \in \Lambda[0, t]$, by Propositions 3.14, 3.21 and Theorem 2.22, there exists the quadratic λ -covariation $[A, B]_\lambda$ on $[0, t]$ and the equality

$$A(t)B(t) - (RRS) \int_0^t A_- dB - (RRS) \int_0^t B_- dA = [A, B]_\lambda(t) = \sum_{(0, t]} \Delta^- A \Delta^- B$$

holds. In addition to sample functions A and B , let $M = M(\cdot, \omega)$ and $N = N(\cdot, \omega)$ be càdlàg sample functions which have the local p -variation index equal to 2. For any $t > 0$ and any

$\lambda \in \Lambda[0, t]$, by the preceding argument, there exist the quadratic λ -covariations $[M, B]_\lambda$, $[A, N]_\lambda$ on $[0, t]$, and the equalities

$$M(t)B(t) - (RRS) \int_0^t B_- dM - (RRS) \int_0^t M_- dB = [M, B]_\lambda(t) = \sum_{(0, t]} \Delta^- M \Delta^- B,$$

$$A(t)N(t) - (RRS) \int_0^t A_- dN - (RRS) \int_0^t N_- dA = [A, N]_\lambda(t) = \sum_{(0, t]} \Delta^- A \Delta^- N$$

hold. By Proposition 7, the stochastic integrals $B_- \cdot M$ and $A_- \cdot N$ are indistinguishable from the integral stochastic processes $\int B_- dM$ and $\int A_- dN$, respectively. Then using the linearity of the unconditional sums, it follows that the quadratic covariation $[X, Y]$ is indistinguishable from the stochastic process

$$\left\{ [M, N](t) - \sum_{(0, t]} \Delta^- M \Delta^- N + \sum_{(0, t]} \Delta^- X \Delta^- Y : t \geq 0 \right\}.$$

The proof of the proposition is complete. $\square$

Proposition 16 *Let X, Z be $(2 - \epsilon)$ -semimartingales, let H be an adapted càdlàg stochastic process with the local p -variation index $v(H) = 2$, and let $Y := C + H \odot Z$ for some constant C . Then*

$$[X, Y] = (RRS) \int H_- d[X, Z]. \quad (4.21)$$

Proof. By the assumption there are local martingales M, N and adapted càdlàg stochastic processes A, B having locally bounded p -variation for some $1 \leq p < 2$ such that $M(0) = N(0) = A(0) = B(0) = 0$, $X - X(0) = M + A$ and $Z - Z(0) = N + B$ almost surely. Since $H_- \cdot M$ is a local martingale and $\int H_- dA$ is an adapted càdlàg stochastic process having locally bounded p -variation, Y is a $(2 - \epsilon)$ -semimartingale with $Y(0) = C$. Therefore in (4.21), the two quadratic covariations and the integral stochastic process are defined. Since almost all sample functions of the stochastic processes in (4.21) are càdlàg, it is enough to prove that for any $0 < t < \infty$

$$[X, Y](t) = (RRS) \int_0^t H_- d[X, Z] \quad \text{almost surely.}$$

To this aim fix an arbitrary $0 < t < \infty$. By Proposition 14 applied to the two pairs of $(2 - \epsilon)$ -semimartingales M, B and A, Z in conjunction with Proposition 3.14, the equalities

$$[M, B](t) = \sum_{(0, t]} \Delta^- M \Delta^- B, \quad [A, Z](t) = \sum_{(0, t]} \Delta^- A \Delta^- Z,$$

hold almost surely. By the preceding argument in conjunction with Propositions 2.27 and 11, the two pairs of $(2 - \epsilon)$ -semimartingales $A, H \odot Z$ and $M, \Phi := \int H_- dB$, satisfy the relations

$$[M, \Phi](t) = \sum_{(0, t]} \Delta^- M \Delta^- \Phi = \sum_{(0, t]} H_- \Delta^- M \Delta^- B = (RRS) \int_0^t H_- d[M, B],$$

$$[A, H \odot Z](t) = \sum_{(0, t]} \Delta^- A \Delta^- (H \odot Z) = \sum_{(0, t]} H_- \Delta^- A \Delta^- Z = (RRS) \int_0^t H_- d[A, Z]$$

almost surely. Since M, N are local martingales such that $M(0) = N(0) = 0$, by Theorem II.29 in [95, p. 68], the equality

$$[M, H_- \cdot N](t) = (LS) \int_{[0,t]} H_- d[M, N] = (RRS) \int_0^t H_- d[M, N]$$

holds almost surely, where LS refers to the Lebesgue-Stieltjes integration. Since the quadratic covariation and the RRS integral are bilinear, by relation (4.19), we have that equalities

$$\begin{aligned} [X, Y](t) &= [M + A, H \odot Z](t) = [M, H_- \cdot N](t) + [M, \Phi](t) + [A, H \odot Z](t) \\ &= (RRS) \int_0^t H_- d[M, N] + (RRS) \int_0^t H_- d[M, B] + (RRS) \int_0^t H_- d[A, Z] \\ &= (RRS) \int_0^t H_- d[X, Z] \end{aligned}$$

hold almost surely. The proof of Proposition 16 is complete. $\square$

4.5 The Itô formula and the linear stochastic $\ominus$ -integral equation

First we extend the Itô formula to $(2 - \epsilon)$ -semimartingales using $\ominus$ -integral, and second we show that the stochastic Doléans exponential is the unique $(2 - \epsilon)$ -semimartingale satisfying the linear $\ominus$ -integral equation.

As before we say that a function $\phi: \mathbb{R}^d \mapsto \mathbb{R}$ is a C^2 class function if its all second order partial derivatives exist and are continuous. Then we write $\phi'_k := \frac{\partial \phi}{\partial x_k}$ and $\phi''_{kl} := \frac{\partial^2 \phi}{\partial x_k \partial x_l}$ for $k, l = 1, \dots, d$.

Theorem 17 *Let $X = (X_1, \dots, X_d)$ be a d -tuple of $(2 - \epsilon)$ -semimartingales, and let $\phi: \mathbb{R}^d \mapsto \mathbb{R}$ be a C^2 class function. Then (1) for $k = 1, \dots, d$, the stochastic $\ominus$ -integral $(\phi'_k \circ X) \ominus X_k$ is defined, (2) for $k, l = 1, \dots, d$, the Riemann-Stieltjes integral stochastic process*

$$(RS) \int (\phi''_{kl} \circ X)_- d[X_k, X_l]^c := \left\{ (RS) \int_0^t (\phi''_{kl} \circ X)_- d[X_k, X_l]^c : t \geq 0 \right\}$$

is defined, (3) the unconditional sum stochastic process

$$\sum \left\{ \Delta^-(\phi \circ X) - \sum_{k=1}^d (\phi'_k \circ X)_- \Delta^- X_k \right\} := \left\{ \sum_{(0,t]} \left\{ \Delta^-(\phi \circ X) - \sum_{k=1}^d (\phi'_k \circ X)_- \Delta^- X_k \right\} : t \geq 0 \right\}$$

is adapted càdlàg stochastic process with locally bounded variation, and (4) the composition $\phi \circ X$ is a $(2 - \epsilon)$ -semimartingale and

$$\begin{aligned} \phi \circ X &= (\phi \circ X)(0) + \sum_{k=1}^d (\phi'_k \circ X) \ominus X_k + \frac{1}{2} \sum_{k,l=1}^d (RS) \int (\phi''_{kl} \circ X)_- d[X_k, X_l]^c \\ &\quad + \sum \left\{ \Delta^-(\phi \circ X) - \sum_{k=1}^d (\phi'_k \circ X)_- \Delta^- X_k \right\}. \end{aligned} \tag{4.22}$$

Proof. Composing with a Lipschitz smooth function does not change the p -variation property. Thus the stochastic $\ominus$ -integrals $(\phi'_k \circ X) \ominus X_k$, $k = 1, \dots, d$, exist by Theorem 12. The Riemann-Stieltjes stochastic processes are defined and have almost all sample functions continuous because each $[X_k, X_t]^c$ has almost all sample functions continuous with locally bounded variation and each integrand is a regulated stochastic process. Applying the mean value theorem and Hölder's inequality it follows that the unconditional sum stochastic process is defined because almost surely

$$\sum_{(0,t]} \{\Delta^- X\}^2 \leq [X](t) < \infty$$

for each $t > 0$. By Theorem 12 again, the stochastic $\ominus$ -integral is a $(2 - \epsilon)$ -semimartingale, so that the composition $\phi \circ X$ is also a $(2 - \epsilon)$ -semimartingale provided formula (4.22) holds. For notation simplicity the proof of (4.22) is given in the case $d = 1$.

The proof will be complete once we show that the equality

$$\begin{aligned} (\phi \circ X)(t) &= (\phi \circ X)(0) + ((\phi' \circ X)_- \ominus X)(t) + \frac{1}{2}(RS) \int_0^t (\phi'' \circ X)_- d[X]^c \\ &\quad + \sum_{(0,t]} \left\{ \Delta^-(\phi \circ X) - (\phi' \circ X)_- \Delta^- X \right\} \end{aligned} \quad (4.23)$$

holds almost surely for an arbitrary $0 < t < \infty$. Indeed, it then holds almost surely for countably many t 's, and since all the stochastic processes are càdlàg the equality will hold almost surely for all $0 < t < \infty$. Thus let $0 < t < \infty$ be fixed, and let $(M, A) \in \mathcal{D}(X)$, that is, $X = X(0) + M + A$, M is a local martingale, A is an adapted stochastic process with locally bounded p -variation, and $M(0) = A(0) = 0$. For a positive integer m , let $\tau_m := \inf\{s \in [0, t]: |M(s)| > m \text{ or } |A(s)| > m\}$. Then each τ_m is a stopping time such that $\Pr(\cup_m \{\tau_m = t\}) = 1$. Therefore (4.23) holds provided it holds when M and A are replaced by M^{τ_m} and A^{τ_m} respectively. For notation simplicity again, let $|M(s)| \vee |A(s)| < K$ for some finite K and all $s \in [0, t]$ almost surely. Since ϕ'' is uniformly continuous over $[-K, K]$, we then are able to control increments $\phi''(X(s)) - \phi''(X(r))$ almost surely with s and r having values in the domain of X outside of an arbitrary small set.

Let $\epsilon \in (0, 1)$. Then there exists $\delta > 0$ such that $|\phi''(u) - \phi''(v)| < \epsilon$ whenever $|u - v| < \delta$ and $u, v \in [-K, K]$. For almost all $\omega \in \Omega$ there exists a finite set $\mu = \{s_j: j = 1, \dots, k\} \subset (0, t]$ (depending on ω) such that sample functions $M = M(\cdot, \omega)$ and $A = A(\cdot, \omega)$ satisfy the conditions:

$$\left| \sum_{\mu} \{\Delta^-(\phi \circ X) - (\phi' \circ X)_- \Delta^- X\} - \sum_{(0,t]} \{\Delta^-(\phi \circ X) - (\phi' \circ X)_- \Delta^- X\} \right| < \epsilon, \quad (4.24)$$

$$\left| \sum_{\mu} \{(\phi'' \circ X)_- (\Delta^- M)^2\} - \sum_{(0,t]} \{(\phi'' \circ X)_- (\Delta^- M)^2\} \right| < \epsilon, \quad (4.25)$$

$$\max_{1 \leq j \leq k} \text{Osc}(X; [s_{j-1}, s_j]) < \delta \quad \text{and} \quad \sum_{j=1}^k v_2(A; [s_{j-1}, s_j]) < \epsilon, \quad (4.26)$$

where $s_0 := 0$. The first condition in (4.26) holds because almost all sample functions of X are càdlàg, and the second one follows from Lemma 2.9. Let $\{\lambda_m: m \geq 1\}$ be a nested sequence of partitions $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$ of $[0, t]$ such that $\cup_m \lambda_m$ is dense in $[0, t]$, and let $m_0 = m_0(\omega)$ be the minimal integer such that each $[s_{j-1}, s_j] \cap \lambda_m \neq \emptyset$. For each $m \geq m_0$,

let $I_1(m) := \{i = 1, \dots, n(m) : \mu \cap (t_{i-1}^m, t_i^m] \neq \emptyset\}$ and $I_2(m) := \{1, \dots, n(m)\} \setminus I_1(m)$. By a telescoping sum and Taylor's theorem (3.46), for each $m \geq m_0$ we have

$$\begin{aligned} (\phi \circ X)(t) - (\phi \circ X)(0) &= \sum_{i=1}^{n(m)} \phi'(X(t_{i-1}^m)) \Delta_i^m X + \frac{1}{2} \sum_{i \in I_2(m)} \phi''(X(t_{i-1}^m)) (\Delta_i^m M)^2 \\ &+ \sum_{i \in I_1(m)} \{\Delta_i^m (\phi \circ X) - \phi'(X(t_{i-1}^m)) \Delta_i^m X\} + \frac{1}{2} \sum_{i \in I_2(m)} \phi''(y_i^m) [2\Delta_i^m M \Delta_i^m A + (\Delta_i^m A)^2] \\ &+ \frac{1}{2} \sum_{i \in I_2(m)} [\phi''(y_i^m) - \phi''(X(t_{i-1}^m))] (\Delta_i^m M)^2 =: \sum_{l=1}^5 S_l(m), \end{aligned} \quad (4.27)$$

where $\Delta_i^m f := f(t_i^m) - f(t_{i-1}^m)$, $y_i^m := X(t_{i-1}^m) + \theta_i^m \Delta_i^m X \subset [-K, K]$ and $\theta_i^m \in (0, 1)$ for $i = 1, \dots, n(m)$. By Theorem 12 and Theorem 2.22, we have

$$\begin{aligned} S_1(m) &= \sum_{i=1}^{n(m)} \phi'(X(t_{i-1}^m)) \Delta_i^m M + \sum_{i=1}^{n(m)} \phi'(X(t_{i-1}^m)) \Delta_i^m A \\ &\rightarrow ((\phi' \circ X)_- \cdot M)(t) + (RRS) \int_0^t (\phi' \circ X)_- dA = ((\phi' \circ X) \odot X)(t) \end{aligned}$$

in probability as $m \rightarrow \infty$. By Theorem II.30 in [95, p. 69] and by definition of the set $I_1(m)$, we have

$$\begin{aligned} S_2(m) &= \frac{1}{2} \sum_{i=1}^m \phi''(X(t_{i-1}^m)) (\Delta_i^m M)^2 - \frac{1}{2} \sum_{i \in I_1(m)} \phi''(X(t_{i-1}^m)) (\Delta_i^m M)^2 \\ &\rightarrow \frac{1}{2} (RRS) \int_0^t (\phi'' \circ X)_- d[M] - \frac{1}{2} \sum_{\mu} \{(\phi'' \circ X)_- (\Delta^- M)^2\} \end{aligned} \quad (4.28)$$

in probability as $m \rightarrow \infty$. Since almost all sample functions of X are càdlàg, by definition of $I_1(m)$ and because $\cup_m \lambda_m$ is dense in $[0, t]$, it follows that

$$S_3(m) \rightarrow \sum_{\mu} \{\Delta^- (\phi \circ X) - (\phi' \circ X)_- \Delta^- X\} \quad (4.29)$$

almost surely as $m \rightarrow \infty$. Notice that for each $i \in I_2(m)$, $[t_{i-1}^m, t_i^m] \subset [s_{j-1}, s_j]$ for some $j = 1, \dots, k$. Then by Hölder's inequality, and using the second condition in (4.26), we get the bound

$$\begin{aligned} |S_4(m)| &\leq \frac{1}{2} \|\phi''\|_{\infty} \left\{ 2s_2(M; \lambda_m)^{1/2} \left(\sum_{i \in I_2(m)} (\Delta_i^m A)^2 \right)^{1/2} + \sum_{i \in I_2(m)} (\Delta_i^m A)^2 \right\} \\ &\leq \frac{\sqrt{\epsilon}}{2} \|\phi''\|_{\infty} \{2s_2(M; \lambda_m)^{1/2} + 1\}. \end{aligned}$$

Finally using the first condition in (4.26), we get the bound

$$|S_5(m)| \leq \frac{1}{2} \max_{i \in I_2(m)} |\phi''(y_i^m) - \phi''(X(t_{i-1}^m))| s_2(M; \lambda_m) \leq \frac{\epsilon}{2} s_2(M; \lambda_m).$$

Since the sums $S_1(m)$, $S_2(m)$ and $S_3(m)$ converge in probability as $m \rightarrow \infty$, due to identity (4.27), it follows that so does the sum $S_4(m) + S_5(m)$. Let $R(\epsilon)$ be the limit of $S_4(m) + S_5(m)$. Notice that bounds of $S_4(m)$ and $S_5(m)$ also converge in probability as $m \rightarrow \infty$. Thus letting $\epsilon \downarrow 0$ it follows that $R(\epsilon) \rightarrow 0$ in probability, while the sums over μ in (4.28) and (4.29) converge to the corresponding sums over $(0, t]$ by (4.24) and (4.25), respectively. By Lemma 3.9, we have that

$$(RRS) \int_0^t (\phi'' \circ X)_- d[M] = (RS) \int_0^t (\phi'' \circ X)_- d[M]^c + \sum_{(0,t]} \{(\phi'' \circ X)_- (\Delta^- M)^2\}.$$

Therefore equality (4.23) holds. The proof of Theorem 17 is complete. $\square$

By the preceding theorem and by Theorem 12 we have:

Corollary 18 *The class of all $(2 - \epsilon)$ -semimartingales is closed by taking a composition with a C^2 class function.*

Now we turn to the linear stochastic $\ominus$ -integral equation. Let X be a $(2 - \epsilon)$ -semimartingale. Since almost surely $\sum_{(0,t]} \{\Delta^- X\}^2 \leq [X](t) < \infty$ for each $t > 0$, by Proposition 3.37 and Lemma 3.43, the product

$$V(X; t) := \prod_{(0,t]} (1 + \Delta^- X) e^{-\Delta^- X}, \quad t > 0,$$

converges absolutely for almost all sample functions of X . The *stochastic Doléans exponential* $\mathcal{E}(X) = \{\mathcal{E}(X; t): t \geq 0\}$ is then defined by

$$\mathcal{E}(X; t) := \exp \{X(t) - X(0) - \frac{1}{2}[X]^c(t)\} V(X; t), \quad t \geq 0,$$

where $V(X; 0) := 1$. Suppose that for some $\lambda \in \Lambda[0, T]$, $0 < T < \infty$, almost all sample functions of X have the quadratic λ -variation on $[0, T]$. Then by Proposition 14, for almost all $\omega \in \Omega$,

$$[X]^c(t, \omega) = [X(\cdot, \omega)]_\lambda^c(t) \quad \text{and} \quad \mathcal{E}(X; t)(\omega) = \mathcal{E}_{\lambda, 0}(X(\cdot, \omega); t)$$

for $0 \leq t \leq T$. That is the stochastic Doléans exponential for X agree with the Doléans exponential for sample functions of X considered in Section 3.5.

For a semimartingale X , Doléans-Dade [18] proved that $\mathcal{E}(X)$ is the unique semimartingale which satisfies the linear stochastic integral equation

$$Y = 1 + Y_- \cdot X.$$

Next this result is generalized to $(2 - \epsilon)$ -semimartingales X .

Theorem 19 *Let X be a $(2 - \epsilon)$ -semimartingale. The stochastic Doléans exponential $\mathcal{E}(X)$ is the unique $(2 - \epsilon)$ -semimartingale which satisfies the linear stochastic $\ominus$ -integral equation:*

$$Y = 1 + Y \ominus X. \tag{4.30}$$

Proof. To begin with we show that $\mathcal{E}(X)$ is a $(2 - \epsilon)$ -semimartingale and satisfies equation (4.30). Let $\bar{X} = \{\bar{X}(t): t \geq 0\}$ and $V = \{V(t): t \geq 0\}$ be stochastic processes defined by

$$\bar{X}(t) := X(t) - X(0) - \frac{1}{2}[X]^c(t) \tag{4.31}$$

and $V(t) := V(X; t)$ for $t \geq 0$. By Lemma 3.43, $V = \{V(t): t \geq 0\}$ is a càdlàg stochastic process with almost all pure jump sample functions and locally of bounded variation. Since $[X]^c$ has continuous sample functions locally of bounded variation, $\bar{X}$ and V are $(2 - \epsilon)$ -semimartingales. Let $\phi(u, v) := e^u v$ for $u, v \in \mathbb{R}$. Then $\mathcal{E}(X) = \phi(\bar{X}, V)$, $(\bar{X}, V)$ is a 2-tuple $(2 - \epsilon)$ -semimartingale and $\phi: \mathbb{R}^2 \mapsto \mathbb{R}$ is a C^2 class function. By Theorem 17, $\mathcal{E}(X)$ is a $(2 - \epsilon)$ -semimartingale and

$$\begin{aligned}\mathcal{E}(X) &= 1 + \mathcal{E}(X) \ominus \bar{X} + e^{\bar{X}} \ominus V + \frac{1}{2}(RS) \int \mathcal{E}(X)_- d[X]^c \\ &\quad + \sum \left\{ \Delta^- \mathcal{E}(X) - \mathcal{E}(X)_- \Delta^- X - e^{\bar{X}-} \Delta^- V \right\} \\ &= 1 + \mathcal{E}(X) \ominus X + e^{\bar{X}} \ominus V - \sum \left\{ e^{\bar{X}-} \Delta^- V \right\}.\end{aligned}$$

The last equality holds by Proposition 10 and because $\Delta^- \mathcal{E}(X) = \mathcal{E}(X)_- \Delta^- X$ by (3.114). Since almost all sample functions of V are pure jump functions of locally bounded variation, the equality

$$e^{\bar{X}} \ominus V = \left\{ (RRS) \int_0^t e^{\bar{X}-} dV: t \geq 0 \right\} = \left\{ \sum_{(0,t]} e^{\bar{X}-} \Delta^- V: t \geq 0 \right\} = \sum \left\{ e^{\bar{X}-} \Delta^- V \right\}$$

holds almost surely. Therefore $\mathcal{E}(X)$ is indistinguishable from $1 + \mathcal{E}(X) \ominus X$, that is, the stochastic Doléans exponential $\mathcal{E}(X)$ satisfies equation (4.30).

Now we prove that $\mathcal{E}(X)$ is the unique solution in the class of all $(2 - \epsilon)$ -semimartingales. Let $Y = \{Y(t): t \geq 0\}$ be a $(2 - \epsilon)$ -semimartingale and a solution of (4.30). Let $V = \{V(t): t \geq 0\}$ be a stochastic process defined by $V(t) := Y(t)e^{-\bar{X}(t)}$, $t \geq 0$, where $\bar{X}(t)$ is defined by (4.31), and let $\psi(u, v) := e^{-u}v$ for $u, v \in \mathbb{R}$. Then $V = \psi(\bar{X}, Y)$, $(\bar{X}, Y)$ is a 2-tuple $(2 - \epsilon)$ -semimartingale, and $\psi: \mathbb{R}^2 \mapsto \mathbb{R}$ is a C^2 class function. It is enough to prove that V is indistinguishable from the product stochastic process

$$\{V(X; t): t \geq 0\} = \left\{ \prod_{(0,t]} (1 + \Delta^- X) e^{-\Delta^- X}: t \geq 0 \right\}. \quad (4.32)$$

To this aim we again apply Theorem 17, which in this case yields the first equality

$$\begin{aligned}V &= 1 - V \ominus \bar{X} + e^{-\bar{X}} \ominus Y + \frac{1}{2}(RS) \int V_- d[X]^c - (RS) \int e^{-\bar{X}-} d[X, Y]^c \\ &\quad + \sum \left\{ \Delta^- V + V_- \Delta^- X - e^{-\bar{X}-} \Delta^- Y \right\} \\ &= 1 - V \ominus X + \frac{1}{2}(RS) \int V_- d[X]^c + V \ominus X + \frac{1}{2}(RS) \int V_- d[X]^c - (RS) \int V_- d[X]^c \\ &\quad + \sum \left\{ \Delta^- V + V_- \Delta^- X - V_- \Delta^- X \right\} = 1 + \sum \Delta^- V.\end{aligned} \quad (4.33)$$

The second equality follows by Proposition 10, Theorem 13 and Proposition 16. By (4.33), almost all sample functions of V are right-continuous and pure jump functions of locally bounded variation. Since Y is a solution to the equation (4.30), we have the equality

$$\Delta^- V = e^{-\bar{X}- - \Delta^- X} (Y_- + \Delta^- Y) - e^{-\bar{X}-} Y_- = V_- \left\{ (1 + \Delta^- X) e^{-\Delta^- X} - 1 \right\}$$

valid almost surely. For $t > 0$, let

$$W(t) := \sum_{(0,t]} \left\{ (1 + \Delta^- X) e^{-\Delta^- X} - 1 \right\}$$

and $W(0) := 0$. Due to relation (3.87), $W = \{W(t) : t \geq 0\}$ is a stochastic process with almost all càdlàg and pure jump sample functions locally of bounded variation. This in conjunction with (4.33), implies that V is a solution to the linear stochastic integral equation

$$V = 1 + (RRS) \int V_- dW = 1 + (LY) \int V dW.$$

By Theorem 3.47, the solution is unique in the class of all stochastic processes with regulated and right-continuous sample functions. Thus V is indistinguishable from the product stochastic process (4.32). The proof of Theorem 19 is complete. $\square$

4.6 Concluding remarks

The initial motivation for introducing the class of $(2 - \epsilon)$ -semimartingales was the wish to make a relation between calculi based on the p -variation and martingale properties more transparent. Due to time and space constraints we touch only most elementary and basic aspects of the $(2 - \epsilon)$ -semimartingale calculus.

In Stochastic Analysis, it is a tradition (with some exceptions) to assume that stochastic processes are càdlàg. However it would be interesting to develop a calculus for stochastic processes whose sample functions are regulated functions. Such stochastic processes have different names in different books. Gihman and Skorohod [41, Chapter III, §4] use the name stochastic process without discontinuities of the second kind, while Dellacherie and Meyer [16, IV.20] use the name: functions which are free of oscillatory discontinuities. In these texts one can find conditions of existence of regulated stochastic processes. Concerning this assumption Gihman and Skorohod [41, p. 174] notice that when dealing with regulated functions one does not distinguish between two functions having at each point the same left and right limits (sic!). Therefore it is natural to choose a certain convention concerning the value of such a function at the discontinuity point. Gihman and Skorohod [41] consider the space of regulated functions which are continuous either from the left or from the right at each point. However from the point of view of applications it seems useful to have a calculus which makes no such restrictions in advance.

Chapter 5

Stock price modelling in continuous time

In practice, both market practitioners and regulators are aware that equilibrium is an illusion. It is rare to find a field in which theory and practice are so far apart, leaving ample room for alchemy and other forms of magic.

G. Soros. The Crisis of Global Capitalism [Open Society Endangered]. Public Affairs, New York, 1998, p. 41.

In continuous time finance the uncertainty related to a future stock price behaviour is modelled by introducing a probability space with a (semi)martingale $X = \{X_t \equiv X(t): 0 \leq t \leq T\}$. A stock price dynamics $P = \{P_t \equiv P(t): 0 \leq t \leq T\}$ is then described by means of a solution of a stochastic differential equation driven by X . Usually, it is justifiable by the fact that the driving process (the return) satisfies a one of several variants of the Random Walk Hypotheses. In this chapter we continue developing a continuous time stock price model undertaken in [84] without using Stochastic Analysis constructions and suggest a different a motivation.

5.1 Evolutionary asset pricing model

Model introduction. Prevailing stock market models are based on Probability Theory. In financial applications, a probability space $(\Omega, \mathcal{F}, \Pr)$ is considered as a set of possible "states of the world", where the probability $\Pr$ attach suitable weights on admissible events in the σ -algebra $\mathcal{F}$. In continuous time stock price models, a set of possible "states of the world" can be expressed by a set of possible price trajectories in continuous time. Under the Random Walk Hypotheses, typical distributions of price trajectories have support in a set of functions having unbounded variation, and so we need a calculus applicable to such functions. Stochastic Analysis provides several forms of calculi, such as the semimartingale theory or the Malliavin calculus, applicable to families of functions with unbounded variation, which are sample functions of suitable stochastic processes. The fact that possible "states of the world", that is, possible stock price trajectories, are indistinguishable by intrinsic constructions of Stochastic Analysis, such as the Itô integral, is an imperfection of this calculus. Therefore it is natural to look for alternative approaches to continuous time stock price models which allow a consideration of a single price

trajectory, a single trading strategy and so on (see Willinger and Taqqu [110] for a thorough discussion).

An interpretation of theoretical constructions, such as a set of possible "states of the world", is an important aspect of a model building. In fact, an interpretation renders a formal theory into a model. Here by a formal theory we mean an area of Mathematics, and not an Economic Theory, and therefore a resulting model do not provide economically meaningful explanations or predictions. Rather our model may serve as a motivation for further developments both in an area of Mathematics and in an area of Financial Theory. As it is usual in social sciences, a suggested interpretation includes a subjective character of stock price behaviour. Due to a human (scientific) activity, stock markets differ fundamentally from usual subjects of natural sciences. While the latter do not depend on existing theories, stock markets are very sensitive to various scientific interpretations. Therefore it is common to incorporate a human factor into market models. For example, in stochastic stock price models, a flow of σ -algebras of events is interpreted as a flow of information available for market traders. As compared to stock price models motivated by the Random Walk Hypothesis, our interpretation concerns a transformation relating a price and its return, or equivalently, a form of a return. Such an interpretation has no meaning in models based on Stochastic Analysis due to the aforementioned indistinguishability of price trajectories.

For simplicity we consider a market which consists of a single stock and an investor-specialist to be called a participant of the market. The basic assumption of our model is that participant's thinking effects and is effected by a stock price behaviour. In other words, we want to model a connection between a stock and market participants' expectations of the stock. Due to this connection a stock price and participants' expectations becomes equally (un)predictable. To say informally, the model consists of an ordered pair (M, A) of two sets of functions defined on $[0, T]$, and a one-to-one mapping R from M onto A , called a *reflexivity mapping for (M, A)* . Elements of the set M are called *price scenarios* $P = \{P(t): 0 \leq t \leq T\}$ during the time period $[0, T]$, and the stock is identified with the set M . Participants' expectations are represented by the set A of functions $X = \{X(t): 0 \leq t \leq T\}$, each of which is called a *return* of a price scenario $P \in M$ assigned by the reflexivity mapping R . Different elements of the pair (M, A) may be interpreted as different states of the world.

In continuous time finance it is customary to define a price evolution by a unique (strong) solution $P = \{P_t \equiv P(t): 0 \leq t \leq T\}$ of a stochastic differential equation

$$dP_t = \mu(t, P_t) dt + \sigma(t, P_t) dX_t, \quad 0 \leq t \leq T, \quad (5.1)$$

with the initial condition $P_0 = x$, driven by a semimartingale $X = \{X_t \equiv X(t): 0 \leq t \leq T\}$. Following the suggested construction, we take the sets M and A to be supports of the distributions of P and X , respectively. The reflexivity mapping between M and A then could be defined provided one can solve (5.1) path by path. Typically such a mapping may not be possible to express using intrinsic constructions of Stochastic Analysis. Unique solutions of (5.1) give rise to the solution mapping, sometimes called the Itô mapping, defined on a class of driving processes X . But except for special cases, the Itô mapping need not be given by a mapping defined on sample functions of the driving process X because the Itô integral is not defined for sample functions. This aspect is important in econometric analysis of stock price models, and sometimes makes simple exponential models preferable over the models given by solutions of (5.1). The pathwise aspect of the stochastic calculus is also related to the problem of robustness as discussed e.g. in Section 5.C of Bouleau and Lépingle [11]. By the above introductory remarks,

the reflexivity mapping, if exists, need not be continuous in a usual sense. However roughly speaking our reflexivity mapping may be considered as a substitute of the stochastic differential equation (5.1) in continuous time models based on Stochastic Analysis.

In (5.1) taking X to be a standard Brownian motion B , $\sigma(t, u) \equiv u$, $\mu \equiv 0$ and the initial condition $x = 1$, we get a simple case of the Black and Scholes model with the price evolution $P_B(t) := \exp\{B(t) - t/2\}$ for $0 \leq t \leq T$. Let A be a set of continuous sample functions of B . Since the mapping $b \mapsto \{\exp\{b(t) - t/2\}: 0 \leq t \leq T\}$, $b \in A$, is invertible on its range, say M , the set (M, A) gives an example of a pair of sets with a reflexivity mapping. If A is a subset of the set of all right-continuous functions on $[0, T]$ of bounded p -variation for some $0 < p < 2$ then by the results of Dudley and Norvaiša [23, Part II], some natural reflexivity mappings can be extended to analytic mappings.

Reflexivity mapping. More specifically, the reflexivity mapping will be modelled by means of a duality mapping between multiplicative and additive functions on the simplex of an extended interval (see Section 2.1). Recall that a function $\pi = \{\pi(s, t): (s, t) \in S[[0, T]]\}$ is multiplicative if $\pi(s, r) = \pi(s, t)\pi(t, r)$ for each $s, t, r \in [[0, T]]$ such that $0 \leq s \leq t \leq r \leq T$, and $\pi(t, t) = 1$ for $t \in [[0, T]]$. Also, a function $\mu = \{\mu(s, t): (s, t) \in S[[0, T]]\}$ is additive if $\mu(s, r) = \mu(s, t) + \mu(t, r)$ for each $s, t, r \in [[0, T]]$ such that $0 \leq s \leq t \leq r \leq T$, and $\mu(t, t) = 0$ for $t \in [[0, T]]$. For a given regulated function f on $[0, T]$, $\mu(f)$ is additive and upper continuous function on $S[[0, T]]$ defined by $\mu(f; s, t) := f(t) - f(s)$ for $(s, t) \in S[[0, T]]$, and $\pi(f)$ is multiplicative and upper continuous function on $S[[0, T]]$ defined by $\pi(f; s, t) := f(t)/f(s)$ for $(s, t) \in S[[0, T]]$.

Definition 1 Let $\mathcal{M} = \mathcal{M}[0, T]$ be a set of multiplicative functions, and let $\mathcal{A} = \mathcal{A}[0, T]$ be a set of additive functions. If there is a one-to-one mapping $\mathcal{R}$ from $\mathcal{M}$ onto $\mathcal{A}$, then we say that $(\mathcal{R}, \mathcal{M}, \mathcal{A})$ is a *duality triple* over $[0, T]$, and $\mathcal{R}$ is called a *reflexivity mapping*.

It will be seen from what follows that a reflexivity mapping for a given $\mathcal{M}$ and $\mathcal{A}$ may *not* be unique. We start with a simple example. Let $\mathcal{M}[0, T]$ be the set of all multiplicative functions on $S[[0, T]]$ which are positive, and let $\mathcal{A}[0, T]$ be the set of all real-valued additive functions on $S[[0, T]]$. Then taking the logarithm $\mathcal{R}_{\log}\pi(s, t) := \log \pi(s, t)$, $(s, t) \in S[[0, T]]$, induces the reflexivity mapping $\mathcal{R}_{\log}$ from $\mathcal{M}[0, T]$ onto $\mathcal{A}[0, T]$. In the next subsection we describe a different reflexivity mapping acting between subsets of $\mathcal{M}[0, T]$ and $\mathcal{A}[0, T]$.

In the cases considered below a natural domain of a reflexivity mapping is quite a large set. Therefore for modelling purposes, it is reasonable restrict a consideration to pairs of subsets defined as follows:

Definition 2 Let $(\mathcal{R}, \mathcal{M}, \mathcal{A})$ be a duality triple on $[0, T]$, and let M, A be two sets of regulated functions on $[0, T]$ such that $f(0) = 1$ for $f \in M$, $g(0) = 0$ for $g \in A$,

$$\pi(M) := \{\pi(f): f \in M\} \subset \mathcal{M} \quad \text{and} \quad \mu(A) := \{\mu(g): g \in A\} \subset \mathcal{A}.$$

If $\mathcal{R}(\pi(M)) = \mu(A)$ then we say that the pair (M, A) is an $\mathcal{R}$ -*system* and the mapping $R: M \rightarrow A$ defined by $R(f) := R_{\mathcal{R}\pi(f)}$, $f \in M$, is called the *reflexivity mapping* for (M, A) , where $R_\mu(t) := \mu(0, t)$, $t \in [0, T]$ is the right distribution function of μ restricted to $[0, T]$.

Let $(\mathcal{R}, \mathcal{M}, \mathcal{A})$ be a duality triple on $[0, T]$, and let (M, A) be an $\mathcal{R}$ -system. By Corollaries 2.5 and 2.6, the reflexivity mapping R for (M, A) is one-to-one with the inverse $R^{-1}: A \mapsto M$,




Figure 5.1: A reflexivity mapping for (M, A) and its inverse.

and diagrams of both mappings are given by Figure 5.1. For $(P, X) \in (M, A)$ and for each $0 \leq s \leq t \leq T$, we have

$$\frac{P(t)}{P(s)} = \pi(P; s, t) = (\mathcal{R}^{-1}\mu(X))(s, t) \quad \text{and} \quad X(t) - X(s) = \mu(X; s, t) = (\mathcal{R}\pi(P))(s, t). \quad (5.2)$$

As it was said earlier a stock is identified with a set M of possible price scenarios, and market participants' expectations are represented by a set of returns A . This model may be considered as a form of the "Concept of Reflexivity" introduced and advocated by Soros [103, p. 42]. The first relation in (5.2) means that investors make decisions in terms of the percentage changes in prices rather than in terms of absolute prices. In the context of finance this may be called the *property of relative price changes*. This idea was first formulated and discussed in the literature by Osborne [89] and Samuelson [101] (see also Bernstein [5, pp. 103-105]). It differs in an important way from Bachelier's hypothesis concerning differences of a stock price values. The second relation in (5.2) means that returns X allow for aggregation over different time periods, the property preferable when one is trying to test various assumptions made on the return X . Let us call it as the *aggregational property of returns*. The two "properties" are so important for the present interpretation that in a formal theory of a market they should be called as axioms.

Evolutionary model. Here we describe a reflexivity mapping induced by the λ -generator of a regulated evolution introduced in Section 3.7.

Definition 3 Let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[0, T]$. A set $\mathcal{M} = \mathcal{M}[0, T]$ of multiplicative interval functions on $S[[0, T]]$ will be called a *set of λ -evolutions*, and a mapping $\mathcal{R}_\lambda$ on $\mathcal{M}$ will be called a *λ -reflexivity mapping* if (a) and (b) hold, where

(a) for each $\pi \in \mathcal{M}$ there is a function $\mathcal{R}_\lambda(\pi) := \mu$ on $S[[0, T]]$ such that for each $(s, t) \in S[[0, T]]$,

$$\mu(s, t) = \lim_{m \rightarrow \infty} S(\pi; \lambda_m \cap [s, t]), \quad (5.3)$$

where $S(\pi; \kappa) := \sum_{i=1}^n [\pi(t_{i-1}, t_i) - 1]$ for a partition $\kappa = \{t_i: i = 0, \dots, n\}$;

(b) for each $\pi_1, \pi_2 \in \mathcal{M}$, if $\mathcal{R}_\lambda(\pi_1) = \mathcal{R}_\lambda(\pi_2)$ then $\pi_1 = \pi_2$.

It is clear that a function μ arising in condition (a) of the preceding definition, if it exists, is additive and unique, and so $\mathcal{R}_\lambda$ is a mapping on $\mathcal{M}$ with values in a class of additive interval functions on $[0, T]$. By (b), $\mathcal{R}_\lambda$ is a one-to-one mapping onto the range $\mathcal{A} := \mathcal{R}_\lambda(\mathcal{M})$. Therefore,

$(\mathcal{R}_\lambda, \mathcal{M}, \mathcal{A})$ is a duality triple over $[0, T]$ provided a set $\mathcal{M}$ of λ -evolutions and a λ -reflexivity mapping $\mathcal{R}_\lambda$ on $\mathcal{M}$ exist.

To construct a set of λ -evolutions we use the set $L_\lambda[0, T]$ defined by (1.23). This and related set $E_\lambda[0, T]$ defined by (1.24), were used in Section 3.7 to solve the evolution representation problem for functions having the quadratic λ -variation. For $\lambda \in \Lambda[0, T]$, let

$$\mathcal{M}_\lambda = \mathcal{M}_\lambda[0, T] := \pi(L_\lambda[0, T]) := \{\pi(f) : f \in L_\lambda[0, T]\}, \quad (5.4)$$

$$\mathcal{A}_\lambda = \mathcal{A}_\lambda[0, T] := \mu(E_\lambda[0, T]) := \{\mu(g) : g \in E_\lambda[0, T]\}. \quad (5.5)$$

For each $f \in L_\lambda[0, T]$, let

$$\mathcal{R}_\lambda \pi(f) := (LC) \int f^{-1} d_\lambda f, \quad (5.6)$$

where the right side is defined by Proposition 3.56. By Proposition 3.53, each $\pi(f)$, $f \in L_\lambda[0, T]$, is a regulated evolution on $[0, T]$, and condition (a) of Definition 3 holds provided $\pi(f)$ has the λ -generator of Definition 3.54. The latter holds by Theorem 3.57, and so condition (a) of Definition 3 holds for the set $\mathcal{M}_\lambda$. While condition (b) of Definition 3 holds for $\mathcal{M}_\lambda$ by Theorem 3.58. Moreover by Propositions 3.56 and 3.19, $\mathcal{R}_\lambda(\mathcal{M}_\lambda) \subset \mathcal{A}_\lambda$, and by Theorem 3.59, $\mathcal{R}_\lambda(\mathcal{M}_\lambda) \supset \mathcal{A}_\lambda$. We thus proved the following:

Corollary 4 *For $\lambda \in \Lambda[0, T]$, $\mathcal{M}_\lambda$ is a λ -evolutionary set, $\mathcal{R}_\lambda$ is a λ -reflexivity mapping from $\mathcal{M}_\lambda$ onto $\mathcal{A}_\lambda$, and $(\mathcal{R}_\lambda, \mathcal{M}_\lambda, \mathcal{A}_\lambda)$ is a duality triple.*

In our financial interpretation, $\mathcal{M}_\lambda$ is the set of price scenarios, $\mathcal{A}_\lambda$ is the set of returns of price scenarios assigned by the reflexivity mapping $\mathcal{R}_\lambda$. Therefore given $\lambda \in \Lambda[0, T]$, the duality triple $(\mathcal{R}_\lambda, \mathcal{M}_\lambda, \mathcal{A}_\lambda)$ is called the λ -evolutionary asset pricing model. If $\mu \in \mathcal{A}_\lambda$ then $\mu = \mathcal{R}_\lambda \pi(f)$ for some $f \in L_\lambda[0, T]$, and hence by Theorem 3.58, the inverse to $\mathcal{R}_\lambda$ is given by

$$\mathcal{R}_\lambda^{-1} \mu = \mathfrak{V}(1 + d_\lambda(\mathcal{L}_\lambda f)) = \pi(f), \quad (5.7)$$

where $\mathcal{L}_\lambda f(t) = \mu(0, t) = (LC) \int_0^t f^{-1} d_\lambda f$ for $t \in [0, T]$. Next is a special case of Definition 2.

Definition 5 Let $(\mathcal{R}_\lambda, \mathcal{M}_\lambda, \mathcal{A}_\lambda)$ be the λ -evolutionary asset pricing model for some $\lambda \in \Lambda[0, T]$, and let P, X be two sets of regulated functions on $[0, T]$ such that $f(0) = 1$ for $f \in P$, $g(0) = 0$ for $g \in X$, $P \subset L_\lambda[0, T]$ and $X \subset E_\lambda[0, T]$. The pair (P, X) is called a λ -evolutionary system if it is an $\mathcal{R}_\lambda$ -system, that is, if $\mathcal{R}_\lambda(\pi(P)) = \mu(X)$. Moreover, the reflexivity mapping for (P, X) , say R_λ , is called the λ -reflexivity mapping for (P, X) . The set P is called a *stock of the λ -evolutionary model*, or simply a *stock*, and each element of P is called a *stock price scenario*.

Remark 6 For notation convenience P may denote a set (= a stock) and an element of a set (= a stock price scenario). If there is a possibility of a confusion we also may use a notation for a stock price scenario different from a stock. The same rule applies to a return X . Typical examples of a λ -evolutionary system are pairs of stochastic process, and so our rule conforms to the tradition in Stochastic Analysis to denote a stochastic process and a sample function by the same letter.

Let (P, X) be a λ -evolutionary system for some $\lambda \in \Lambda[0, T]$. By (5.6), the λ -reflexivity mapping for (P, X) is $\mathcal{L}_\lambda$, that is, the mapping from P onto X induced by the right distribution function of the Left Cauchy λ -integral. By (5.7), the inverse of $R_\lambda = \mathcal{L}_\lambda$ is the right distribution function $\mathcal{P}_\lambda$ of the product λ -integral (see Figure 5.1). By Theorem 3.42, the inverse $R_\lambda^{-1} = \mathcal{P}_\lambda$ is also equal to the mapping induced by the forward Doléans exponential $\mathcal{E}_\lambda(g) := \mathcal{E}_{\lambda,0}(g)$ defined by (3.109) with $[a, b] = [0, T]$, for $g \in E_\lambda[0, T]$.

Proposition 7 For $\lambda \in \Lambda[0, T]$ and $X \subset E_\lambda[0, T]$, $(\mathcal{E}_\lambda(X), X)$ is a λ -evolutionary system. Also for each $g \in X$, $f := \mathcal{E}_\lambda(g) \in \mathcal{E}_\lambda(X)$ and $0 \leq t \leq T$,

$$f(t) = 1 + (LC) \int_0^t f d_\lambda g, \quad (5.8)$$

$$[f]_\lambda(t) = (LY) \int_0^t f^2 d[g]_\lambda \quad \text{and} \quad [f, g]_\lambda(t) = (LY) \int_0^t f d[g]_\lambda. \quad (5.9)$$

Proof. Let $\lambda \in \Lambda[0, T]$ and $X \subset E_\lambda[0, T]$. For $g \in X$, by Theorem 3.44, $\mathcal{E}_\lambda(g)$ has the quadratic λ -variation. Also since $1 + \Delta^- g \wedge \Delta^+ g \gg 0$, by Lemma 3.37 and Corollary 2.6, $\mathcal{E}_\lambda(g) \gg 0$. Since $g \in X$ is arbitrary, $\mathcal{E}_\lambda(X) \subset L_\lambda[0, T]$. Again let $g \in X$. By Theorem 2.4 and Corollary 3.41, $\pi(\mathcal{E}_\lambda(g)) = \beta_\lambda(g)$, and so by Theorem 3.42, $\mathfrak{V}(1 + d_\lambda g) = \pi(\mathcal{E}_\lambda(g))$. Therefore by Theorem 3.59,

$$\mathcal{R}_\lambda \pi(\mathcal{E}_\lambda(g)) = (LC) \int (\mathcal{P}_\lambda g)^{-1} d_\lambda (\mathcal{P}_\lambda g) = \mu(g).$$

Since $g \in X$ is arbitrary, $\mathcal{R}_\lambda \pi(\mathcal{E}_\lambda(X)) = \mu(X)$, that is, $(\mathcal{E}_\lambda(X), X)$ is a λ -evolutionary system. Finally (5.8) is a consequence of Theorem 3.48, and (5.9) is a consequence of Theorem 3.44. $\square$

The following is a λ -evolutionary version of the Black and Scholes model.

Example 8 Let B be a standard Brownian motion on a complete probability space $(\Omega, \mathcal{F}, \text{Pr})$. Given $\lambda \in \Lambda[0, T]$, almost all sample functions $b := B(\cdot, \omega)$, $\omega \in \Omega$, of B are continuous and have the quadratic λ -variation. Thus denoting by B_λ the set of all such functions b , and denoting by P_{B_λ} the set of all functions $\exp\{b(t) - t/2\}$, $0 \leq t \leq T$, we have that $(P_{B_\lambda}, B_\lambda)$ is a λ -evolutionary system. To see this notice that $P_{B_\lambda} = \{\mathcal{E}_\lambda(b) : b \in B_\lambda\}$ and apply the preceding proposition. Letting $N(\lambda) := \{\omega \in \Omega : B(\cdot, \omega) \notin B_\lambda\}$, we have $\cup\{N(\lambda) : \lambda \in \Lambda[0, T]\} = \Omega$, Pr -almost surely.

5.2 Almost sure approximation by a discrete time model

Here we prove that the λ -evolutionary version of the Black and Scholes model of Example 8 is a limit of discrete time binomial models. It is boring to say once again that the point here is an almost sure convergence because in probability variant of such approximation is well-known.

Returns in discrete time. We start by showing that the two continuous time price dynamics described by the reflexivity mappings $\mathcal{R}_\lambda$ and $\mathcal{R}_{\log}$, accord to two discrete time models based on different notions of returns. Consider a discrete time model of a stock price $\{P(t) : t = 0, 1, \dots, T\}$ with the *simple net return* $R_{\text{net}}(t)$ defined by

$$R_{\text{net}}(t) := \frac{P(t)}{P(t-1)} - 1, \quad t = 1, \dots, T.$$

Also we assume that $P(0) = 1$ and $P(t) > 0$ for $t = 1, \dots, T$. As it is usual for discrete time models, their returns are defined at the right endpoints t of unitary time lags $[t-1, t]$ since such returns always represent a change on the stock price between dates $t-1$ and t . To imbed a discrete time model into a continuous time model one has to redefine the discrete time returns as follows: let $X(0) := 0$ and $X(t) - X(t-1) := R_{\text{net}}(t)$ for $t = 1, \dots, T$. Adding up the increments of X , we get the relations

$$X(t) = \sum_{s=1}^t R_{\text{net}}(s) = \sum_{s=1}^t \frac{P(s) - P(s-1)}{P(s-1)}, \quad t = 1, \dots, T. \quad (5.10)$$

Let $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[0, T]$ with $\lambda_m = \{t_i^m: i = 0, \dots, n(m)\}$. Suppose that a price scenario P has its positive values defined on the interval $[0, T]$. Then for each $m \geq 1$, P restricted to λ_m gives the discrete time model with the simple net return X_m such that

$$X_m(t_i^m) - X_m(t_{i-1}^m) = \frac{P(t_i^m)}{P(t_{i-1}^m)} - 1, \quad i = 1, \dots, n(m).$$

Relation (5.3) with $\pi(t_i^m, t_{i-1}^m) = P(t_i^m)/P(t_{i-1}^m)$ for each i , means that $\mu(t, s)$ can be approximated arbitrarily closely by $X_m(t) - X_m(s)$ as $m \rightarrow \infty$. Thus the function μ satisfying (5.3), if exists, extends the simple net return X to a continuous time setting.

The following relations in the discrete setting illustrate the duality relations (5.2) used to define a reflexivity mapping $\mathcal{R}$. Relations (5.10) and

$$P(t) = \prod_{s=1}^t [1 + X(s) - X(s-1)], \quad t = 1, \dots, T, \quad (5.11)$$

provide a one-to-one correspondence between a positive price P and a simple net return X having jumps bigger than -1 . Instead of (5.11) one can use the equivalent relation

$$P(t) = 1 + \sum_{s=1}^t P(s-1)[X(s) - X(s-1)].$$

For each $t = 1, \dots, T$, let $\mathcal{L}(P)(t)$ be defined by the right side of (5.10), and let $\mathcal{E}(X)(t)$ be defined by the right side of (5.11). Then for $t = 1, \dots, T$, we have

$$\mathcal{L}(\mathcal{E}(X))(t) = \sum_{s=1}^t \left[\frac{\mathcal{E}(X)(s)}{\mathcal{E}(X)(s-1)} - 1 \right] = X(t)$$

and

$$\mathcal{E}(\mathcal{L}(P))(t) = \prod_{s=1}^t \left(1 + \mathcal{L}(P)(s) - \mathcal{L}(P)(s-1) \right) = P(t).$$

This duality is used in discrete time stock price models by specifying simple net returns rather than prices (see [93, Section 3.2]).

In econometric analysis instead of the simple net return, one uses the *log-return* R_{log} , defined by

$$R_{log}(t) := \log \left(\frac{P(t)}{P(t-1)} \right), \quad t = 1, \dots, T.$$

The log return is preferable to the simple net return in data analysis because of the aggregational (additivity) property:

$$R_{log}(t) + R_{log}(t-1) + \dots + R_{log}(t-s+1) = \log \left(\frac{P(t)}{P(t-s)} \right)$$

valid for any $s, t \in \{0, 1, \dots, T\}$, $s < t$. Clearly such a property does not hold for the simple net return R_{net} . In contrast, we see that if (5.3) holds then the continuous time analog of the simple net return possess the aggregational property. In sum, a continuous time stock price model based on the reflexivity mapping induced by the logarithm is a limit of discrete time models based on the log-returns, while discrete time models based on the simple net returns give rise to a continuous time models based on the λ -reflexivity mapping to be discussed in more detail in what follows.

Approximation by the binomial model. The binomial model is a simple but yet very important discrete time model of price evolution of a single stock. We show that the binomial price evolution approximates the λ -evolutionary system $(P_{B_\lambda}, B_\lambda)$ defined in Example 8. To recall a simplest variant of the binomial model, let $\epsilon_1, \epsilon_2, \dots$ be a sequence of independent identically distributed random variables on $(\Omega, \mathcal{F}, \Pr)$ such that $\Pr(\{\epsilon_1 = 1\}) = 1 - \Pr(\{\epsilon_1 = 0\}) = 1/2$. For each $t = 1, 2, \dots$ and for any $a \in (0, 1)$, let

$$P(t) := P(0)(1+a)^{N_t}(1-a)^{t-N_t},$$

where $N_t := \epsilon_1 + \dots + \epsilon_t$. According to this time evolution, called the *binomial model*, the price either goes up by the factor $1+a$, or it goes down by the factor $1-a$. The probability of an up move or down move is the same, and the moves over time are independent of each other. The binomial model is a discrete time analogue of the Black and Scholes continuous time model. To justify this one usually refers to a weak convergence of corresponding returns. Next we show that this analogy is much stronger in the sense that the approximation of price scenarios holds with probability 1.

To this aim we use the random walk approximation of a standard Brownian motion originated in Knight [56] and further developed in Section 6.2 of Révész [96] and in Szabados [104]. Let $\{r_1, r_2, \dots\}$ be a sequence of independent identically distributed random variables on $(\Omega, \mathcal{F}, \Pr)$ with the distribution $\Pr(\{r_1 = 1\}) = 1 - \Pr(\{r_1 = -1\}) = 1/2$. Letting $S(0) := 0$ and $S(k) := r_1 + \dots + r_k$ for each $k \geq 1$, we have the *random walk* process $S = \{S(k): k = 0, 1, \dots\}$ on the lattice of all non-negative integers. On a probability space carrying a countable array of independent random walks one can construct a sequence of "twisted" and "shrunk" random walks giving rise to a standard Brownian motion. More specifically the following holds:

Theorem 9 *On a rich enough probability space $(\Omega, \mathcal{F}, \Pr)$ one can construct stochastic processes B and $\{W_m: m \geq 1\}$ such that*

(a) *for a random walk process S and for each $m \geq 1$,*

$$\{W_m(k2^{-2m}): k = 0, 1, \dots\} \stackrel{\text{f.d.d.}}{=} \{2^{-m}S(k): k = 0, 1, \dots\};$$

(b) *for each $m \geq 1$, there is a random partition $\{T_{m+1}(k): k \geq 1\}$ of $[0, \infty)$ such that for each $k \geq 1$, $W_{m+1}(T_{m+1}(k)) = W_m(k2^{-2m})$;*

(c) *B is a standard Brownian motion, and for each $T > 0$, $\|B - \widehat{W}_m\|_{[0, T], \sup} \rightarrow 0$ almost surely as $m \rightarrow \infty$, where $\widehat{W}_m$ is a process indexed by $t \in [0, \infty)$ obtained from W_m by linear interpolation;*

(d) *with probability 1, for any $T' > T > 0$ and for all but finitely many m there exists a random partition $\{\tau_m(k): 0 \leq k \leq T2^{2m}\}$ of $[0, T')$ such that $B(\tau_m(k)) = W_m(k2^{-2m})$ for $0 \leq k \leq T2^{2m}$, where $\max_k |\tau_m(k) - \tau_m(k-1)| \rightarrow 0$ and $\max_k |\tau_m(k) - k2^{-2m}| \rightarrow 0$ both as $m \rightarrow \infty$.*

The construction of B and $\{W_m: m \geq 1\}$ satisfying (a), (b) and (c) is given in [96, Section 6.2] and in [104]. Statement (d) follows from Lemma 7(b) in [104].

Let $m \geq 1$, and let the (discrete time) simple net return X_m be defined by

$$X_m(t) := W_m(t2^{-2m}) \quad \text{for } t = 0, 1, \dots, T2^{2m}.$$

Then by (a) of the preceding theorem, $X_m(0) = 0$ almost surely and $\{X_m(t) - X_m(t-1) : t = 1, \dots, T2^{2m}\}$ are independent identically distributed random variables on $(\Omega, \mathcal{F}, \Pr)$ such that $\Pr(\{X_m(1) = 2^{-m}\}) = 1 - \Pr(\{X_m(1) = -2^{-m}\}) = 1/2$. For each $k = 1, \dots, T2^{2m}$, let $\epsilon_k := 1$ if $X_m(k) - X_m(k-1) = 2^{-m}$ and let $\epsilon_k := 0$ if $X_m(k) - X_m(k-1) = -2^{-m}$. Then the discrete time evolutionary price P_m is given by $P_m(0) := 1$ and for each $t = 1, \dots, T2^{2m}$,

$$P_m(t) = \prod_{k=1}^t [1 + W_m(k2^{-2m}) - W_m((k-1)2^{-2m})] = (1 + 2^{-m})^{N_t} (1 - 2^{-m})^{t-N_t},$$

where $N_t := \epsilon_1 + \dots + \epsilon_t$. Therefore for the probability space $(\Omega, \mathcal{F}, \Pr)$ of Theorem 9, $\{(P_m(\cdot, \omega), X_m(\cdot, \omega)) : \omega \in \Omega\}$ are discrete time binomial models for each integer $m \geq 1$. We show next that on the same probability space we can construct the λ -evolutionary system $(P_{B_\lambda}, B_\lambda)$ of Example 8 for $\lambda = \{\lambda_m : m \geq 1\}$ with $\lambda_m = \{k2^{-2m} : k = 0, \dots, T2^{2m}\}$.

Theorem 10 *Let B and W_m , $m \geq 1$, be stochastic processes on the probability space $(\Omega, \mathcal{F}, \Pr)$ satisfying conditions (a) – (d) of Theorem 9, and let $0 < T < \infty$. Then with probability 1, for each $0 \leq t \leq T$,*

$$\lim_{m \rightarrow \infty} \prod_{k=1}^{\lfloor t2^{2m} \rfloor} [1 + W_m(k2^{-2m}) - W_m((k-1)2^{-2m})] = \exp\{B(t) - t/2\}.$$

Proof. With probability 1, for all but finitely many m , let $\{\tau_m(k) : 0 \leq k \leq T2^{2m}\}$ be random partitions of $[0, T']$ with $T' > T$ satisfying condition (d) of Theorem 9. Then for any $0 < t \leq T$, and for all but finitely many m ,

$$\prod_{k=1}^{\lfloor t2^{2m} \rfloor} [1 + W_m(k2^{-2m}) - W_m((k-1)2^{-2m})] = \prod_{k=1}^{\lfloor t2^{2m} \rfloor} [1 + B(\tau_m(k)) - B(\tau_m(k-1))].$$

Since a Brownian motion is sample continuous, it is enough to prove that for any $0 < t \leq T$, with probability 1,

$$\lim_{m \rightarrow \infty} \prod_{k=1}^{T2^{2m}} [1 + B(\tau_m(k) \wedge t) - B(\tau_m(k-1) \wedge t)] = \exp\{B(t) - t/2\}.$$

Let $0 < t \leq T$. By Proposition 4.10 in [23, Part II], this is so once we show that with probability 1,

$$\lim_{m \rightarrow \infty} \max_k |B(\tau_m(k)) - B(\tau_m(k-1))| = 0$$

and

$$\lim_{m \rightarrow \infty} \sum_{k=1}^{T2^{2m}} [B(\tau_m(k) \wedge t) - B(\tau_m(k-1) \wedge t)]^2 = t.$$

Indeed, for each m , let $n = n_{m,t}$ be the integer such that $(n-1)2^{-2m} < t \leq n2^{-2m}$. By statements (d) and (a) of Theorem 9, we have for all but finitely many m ,

$$|B(\tau_m(k)) - B(\tau_m(k-1))| = |W_m(k2^{-2m}) - W_m((k-1)2^{-2m})| = 2^{-m}$$

for $1 \leq k \leq T2^{2m}$, and

$$\sum_{k=1}^{T2^{2m}} [B(\tau_m(k) \wedge t) - B(\tau_m(k-1) \wedge t)]^2 = (n-1)2^{-2m} + [B(t) - B(\tau_m(n-1))]^2.$$

Since $\tau_m(n-1) - t = [\tau_m(n-1) - (n-1)2^{-2m}] + [(n-1)2^{-2m} - t] \rightarrow 0$ as $m \rightarrow \infty$, the right side of the last display tends to t , proving the above two limiting conditions. The proof of Theorem 10 is complete. $\square$

5.3 Option pricing and hedging

Trading strategies. For $\lambda \in \Lambda[0, T]$ and $k = 0, \dots, n$, let (P_k, X_k) be a λ -evolutionary system (see Definition 5). Then $P = (P_0, \dots, P_n)$ is called a portfolio of $n+1$ stocks of the λ -evolutionary model during a time period $[0, T]$.

Definition 11 Let $P = (P_0, \dots, P_n)$ be a portfolio of $n+1$ stocks of the λ -evolutionary model for some $\lambda \in \Lambda[0, T]$, and let $\gamma_k, k = 0, \dots, n$, be real-valued functions on $[0, T]$. A vector function $\gamma = (\gamma_0, \dots, \gamma_n)$ is called a λ -trading strategy for P if for each $k = 0, \dots, n$, and a price scenario P_k , the Left Cauchy λ -integral $(LC) \int \gamma_k d_\lambda P_k$ is defined on $[0, T)$, $\Delta^- P_k(T) = 0$ whenever $\gamma_k(T-)$ is undefined, and there exists the limit

$$(LC) \int_0^{T-} \gamma_k d_\lambda P_k := \lim_{u \uparrow T} (LC) \int_0^u \gamma_k d_\lambda P_k. \quad (5.12)$$

Also, each γ_k is then called a λ -trading strategy for P_k , and for we let

$$(LC) \int_0^T \gamma_k d_\lambda P_k := \begin{cases} \gamma_k(T-) \Delta^- P_k(T) + (LC) \int_0^{T-} \gamma_k d_\lambda P_k & \text{if } \gamma_k(T-) \text{ exists,} \\ (LC) \int_0^{T-} \gamma_k d_\lambda P_k & \text{otherwise.} \end{cases} \quad (5.13)$$

It will be seen below (Proposition 14) that there is a price scenario P_k and a λ -trading strategy γ_k for P_k such that $\gamma_k(T-)$ does not exist but the limit (5.12) exists. By Definition 3.17, if the Left Cauchy λ -integral $(LC) \int \gamma_k d_\lambda P_k$ is defined on $[0, T]$ then its value on $[0, T]$ agree with (5.13).

Following our convention spelled out by Remark 6, the indefinite integrals for (5.12) and (5.13) may be viewed as families of functions corresponding to different strategies of a stock. Also a λ -trading strategy may denote a set of different strategies as well as a single strategy corresponding to a price scenario.

Definition 12 For $\lambda \in \Lambda[0, T]$, let $P = (P_0, \dots, P_n)$ be a portfolio of $n+1$ stocks of the λ -evolutionary model during a time period $[0, T]$, and let $\gamma = (\gamma_0, \dots, \gamma_n)$ be a λ -trading strategy for P .

(a) For each price scenario P define the functions $V = V(P)$ and $G = G(P)$ defined by

$$V(P; t) := \sum_{k=0}^n \gamma_k(t) P_k(t) \quad \text{and} \quad G(P; t) := \sum_{k=0}^n (LC) \int_0^t \gamma_k d_\lambda P_k$$

are called the *portfolio value function* and the *portfolio gain function*, respectively.

(b) The λ -trading strategy γ is called *self-financing* if $V(P; t) = V(P; 0) + G(P; t)$ for each price scenario P and $t \in [0, T]$.

Black-Scholes formula. Let $\lambda \in \Lambda[0, T]$, $1 \leq p < 2$ and $\sigma > 0$. Let $CQ_{\lambda, \sigma}[0, T]$ be the class of all continuous functions g on $[0, T]$ having the quadratic λ -variation with the bracket function $[g]_{\lambda}(t) = \sigma^2 t$, $0 \leq t \leq T$, and let $CW_p[0, T]$ be the class all continuous functions on $[0, T]$ having bounded p -variation. Consider the class of functions

$$X_{\lambda, p, \sigma}[0, T] := \left\{ w := g + h : g \in CQ_{\lambda, \sigma}[0, T], \quad h \in CW_p[0, T], \quad g(0) = h(0) = 0 \right\}. \quad (5.14)$$

By Corollary 3.16, each function $w \in X$ is continuous and has the quadratic λ -variation with the bracket function $[w]_{\lambda}(t) = \sigma^2 t$, $0 \leq t \leq T$, that is, $w \in CQ_{\lambda, \sigma}[0, T]$. Let X be a subset of $X_{\lambda, p, \sigma}[0, T]$. Thus $X \subset E_{\lambda}[0, T]$, and so by Proposition 7, $(\mathcal{E}_{\lambda}(X), X)$ is a λ -evolutionary system. By Theorem 3.48, for each $w \in X$, $P := \mathcal{E}_{\lambda}(w)$ and $0 \leq t \leq T$,

$$P(t) = 1 + (L) \int_0^t P dw = 1 + (RS) \int_0^t P dh + (LC) \int_0^t P d_{\lambda} g.$$

For example, X could be the set B_{λ} of all sample functions of a standard Brownian motion which are continuous and have the quadratic λ -variation (cf. Example 8), and so $B_{\lambda} \subset X_{\lambda, 1, 1}$. For another example, let B_H be a fractional Brownian motion with the Hurst index $1/2 < H < 1$ defined on the same complete probability space $(\Omega, \mathcal{F}, \Pr)$ as a standard Brownian motion B . Then there is $\Omega_0 \subset \Omega$ such that $\Pr(\Omega_0) = 1$ and $X := \{B(\cdot, \omega) + B_H(\cdot, \omega) : \omega \in \Omega_0\} \subset X_{\lambda, p, 1}$ for any $1/H < p < 2$.

Consider a portfolio (Q, P) with $P = \mathcal{E}_{\lambda}(X) = \{\mathcal{E}_{\lambda}(\omega) : \omega \in X\}$ for some $X \subset X_{\lambda, p, \sigma}[0, T]$, and a set Q consisting of a single price scenario $Q(t) := \exp\{rt\}$, $0 \leq t \leq T$, depending on $r > 0$. The set Q may be considered as a bond, while P is a risky asset. Given $K > 0$ and $r > 0$, we show that there is a self-financing λ -trading strategy (α, β) for the portfolio (Q, P) with the portfolio value on a date T :

$$V(T) = \alpha(T)Q(T) + \beta(T)P(T) = \max\{0, P(T) - K\}$$

for any price scenario (Q, P) . To define the desired λ -trading strategy (α, β) , for $t \in [0, T)$ and $x > 0$, let

$$d_1(t, x) = [\ln(x/K) + (r + \sigma^2/2)(T - t)]/(\sigma\sqrt{T - t}) \quad \text{and} \quad d_2(t, x) = d_1(t, x) - \sigma\sqrt{T - t}.$$

Denoting the standard normal distribution function by N , let $\phi \equiv \phi_{\sigma, K}$ be the function on $[0, T] \times [0, +\infty)$ defined by

$$\phi(t, x) := \begin{cases} xN(d_1(t, x)) - Ke^{-r(T-t)}N(d_2(t, x)) & \text{if } (t, x) \in [0, T) \times (0, +\infty), \\ \max\{0, x - K\} & \text{if } (t, x) \in \{T\} \times (0, +\infty), \\ 0 & \text{if } (t, x) \in [0, T] \times \{0\}. \end{cases} \quad (5.15)$$

Notice that $\phi(T, x) = \lim_{t \uparrow T} \phi(t, x)$ for each $x > 0$, and $\phi(t, 0) = \lim_{x \downarrow 0} \phi(t, x)$ for each $t \in [0, T]$. For a price scenario P , let $\Phi P(t) := \phi(t, P(t))$, $0 \leq t \leq T$. We use notation (3.79) and (3.80) for partial derivatives of ϕ . Then for a price scenario (Q, P) and $0 \leq s \leq T$, let

$$\beta(s) := \begin{cases} \Phi'_x P(s) = N(d_1(s, P(s))) & \text{if } 0 \leq s < T, \\ 0 & \text{if } s = T, \end{cases}$$

and

$$\alpha(s) := \{\Phi P(s) - P(s)\beta(s)\}/Q(s).$$

Therefore the portfolio value function V for a price scenario (Q, P) and $s \in [0, T]$ is given by

$$V(s) = \alpha(s)Q(s) + \beta(s)P(s) = \Phi P(s). \quad (5.16)$$

The values of α and β at $s = T$ do not affect the self-financing condition to be proved next, and they can be chosen arbitrarily so that (5.16) holds for $s = T$. An arbitrage argument assure that the portfolio value function V gives a fair price of a European call option on the stock $P = \mathcal{E}_\lambda(X)$. This price is called the *Black-Scholes formula*.

Theorem 13 For $\lambda \in \Lambda[0, T]$, $1 \leq p < 2$ and $\sigma > 0$, let X be a subset of $X_{\lambda, p, \sigma}[0, T]$, and let $P = \mathcal{E}_\lambda(X)$. Then (α, β) is a self-financing λ -trading strategy for the portfolio (Q, P) .

Proof. Let (Q, P) be a price scenario, that is, $Q(t) = \exp\{rt\}$, $0 \leq t \leq T$, and $P = \mathcal{E}_\lambda(w)$ for some $w \in X$. Therefore $w = g + h$ for some $g \in CQ_{\lambda, \sigma}[0, T]$ and $h \in CW_p[0, T]$. Thus by Corollary 3.16, w has the quadratic λ -variation with the bracket function $[w]_\lambda(t) = [g]_\lambda^\zeta(t) = \sigma^2 t$, $0 \leq t \leq T$. By Proposition 7, $P = \mathcal{E}_\lambda(w)$ has the quadratic λ -variation with the bracket function

$$[P]_\lambda(t) = (LY) \int_0^t \mathcal{E}_\lambda(w)^2 d[w]_\lambda = \sigma^2 \int_0^t P^2(s) ds, \quad 0 \leq t \leq T. \quad (5.17)$$

The function α is bounded on $[0, T]$ and continuous on $[0, T)$. The function Q is continuous and has bounded variation on $[0, T]$. Thus by Theorem II.13.16 in Hildebrandt [49], α is Riemann-Stieltjes integrable on $[0, T]$ with respect to Q , and for $0 \leq t \leq T$,

$$(RS) \int_0^t \alpha dQ = r \int_0^t \alpha(s)Q(s) ds = -r \int_0^t [P(s)\Phi'_x P(s) - \Phi P(s)] ds. \quad (5.18)$$

To prove that β is a λ -trading strategy for $\mathcal{E}_\lambda(w)$ and (α, β) is self-financing we use a chain rule for the composition ΦP . By definition (5.15), ϕ is continuous on $[0, T] \times (0, \infty)$, it is a C^2 class function on $[0, T) \times (0, \infty)$, and

$$\Phi'_t P(s) = \begin{cases} -\frac{\sigma P(s)}{2\sqrt{T-s}} N'(d_1(s, P(s)) - rKe^{-r(T-s)} N(d_2(s, P(s))) & \text{if } s \in [0, T), \\ 0 & \text{if } s = T, \end{cases} \quad (5.19)$$

$$\Phi''_{xx} P(s) = \begin{cases} \frac{1}{\sigma P(s)\sqrt{T-s}} N'(d_1(s, P(s))) & \text{if } s \in [0, T), \\ 0 & \text{if } s = T. \end{cases} \quad (5.20)$$

Notice that ϕ satisfies the partial differential equation

$$\begin{cases} \frac{\partial u}{\partial t} + Lu = 0 & \text{on } [0, T) \times (0, +\infty), \\ u(T, x) = \max\{0, x - K\} & \text{for } x \in (0, +\infty), \\ u(t, 0) = 0 & \text{for } t \in [0, T], \end{cases} \quad (5.21)$$

where

$$Lu(t, x) := \frac{\sigma^2}{2} x^2 \frac{\partial^2 u}{\partial x^2}(t, x) + rx \frac{\partial u}{\partial x}(t, x) - ru(t, x).$$

By the chain rule of Theorem 3.34(1) (cf. (3.82)), the Left Cauchy λ -integral $(LC) \int \Phi'_x P d_\lambda P$ is defined on $[0, T)$, and by (5.16), for each $0 \leq u < T$,

$$\begin{aligned} V(P; u) - V(P; 0) &= (R) \int_0^u \Phi'_t P + (LC) \int_0^u \Phi'_x P d_\lambda P + \frac{1}{2} (RS) \int_0^u \Phi''_{xx} P d[P]_\lambda \\ \text{by (5.17) and (5.18)} &= (RS) \int_0^u \alpha dQ + (LC) \int_0^u \beta d_\lambda P + (R) \int_0^u \left[\frac{\partial \phi}{\partial t} + L\phi \right](s, P(s)) ds \\ \text{by (5.21)} &= (RS) \int_0^u \alpha dQ + (LC) \int_0^u \beta d_\lambda P = G(P; u). \end{aligned} \quad (5.22)$$

The last equality holds because the Left Cauchy λ -integral extends the Riemann-Stieltjes integral.

To prove that the left and right sides of (5.22) are equal for $u = T$ we use the chain rule of Proposition 3.35. Letting $\chi(s) := 1/\sqrt{T-s}$ for $s \in [0, T)$ and $\chi(T) := 0$, there is a finite constant C such that for all $0 \leq s \leq T$,

$$|\Phi'_t P(s)| \leq C\chi(s) \quad \text{and} \quad |\Phi''_{xx} P(s)| \leq C\chi(s).$$

By the Hake theorem (Section 7.3 in [77]) and by the substitution theorem for the Henstock-Kurzweil integral (Section 7.2 in [77]), χ is Henstock-Kurzweil integrable on $[0, T]$ with respect to the identity function and with respect to the bracket function $[P]_\lambda$. By the dominated convergence theorem for the Henstock-Kurzweil integral (Section 7.8 in [77]), it then follows that the Henstock-Kurzweil integrals $(HK) \int_0^T \Phi'_t P$ and $(HK) \int_0^T \Phi''_{xx} P d[P]_\lambda$ are defined. By Proposition 3.35, we then have that

$$(LC) \int_0^{T-} \Phi'_x P d_\lambda P = V(P; T) - V(P; 0) - (HK) \int_0^T \Phi'_t P - \frac{1}{2} (HK) \int_0^T \Phi''_{xx} P d[P]_\lambda.$$

Since $\Delta^- P(T) = 0$, by (5.13), (α, β) is a λ -trading strategy for (Q, P) and $V(P; T) = V(P; 0) + G(P; T)$. The proof is complete. $\square$

The use of the Henstock-Kurzweil integrals in the preceding proof is essential because the functions $\Phi'_t P$ and $\Phi''_{xx} P$ defined by (5.19) and (5.20), respectively, can be unbounded near T . Before showing this, following Bick and Willinger [7], we notice that

$$\lim_{t \uparrow T} d_1(t, P(t)) = \lim_{t \uparrow T} \{ \ln(P(t)/K) / \sigma \sqrt{T-t} \} = \begin{cases} -\infty & \text{if } P(T) < K, \\ +\infty & \text{if } P(T) > K. \end{cases}$$

Thus in the case $P(T) \neq K$, $\Phi'_t P(T-) = \Phi''_{xx} P(T-) = 0$. However the case $P(T) = K$ is different. Bick and Willinger (see Remark following Proposition 1 in [7, p. 366]) conjectured that in this case (α, β) may not be trading strategies in their sense. In contrary the preceding theorem shows that (α, β) is always a self-financing λ -trading strategy. The following shows that $\Phi'_t P$ and $\Phi''_{xx} P$ can be unbounded and gives a positive answer to the second part of the problem of Bick and Willinger [7].

Proposition 14 *Let $\sigma = 1$ and $K = \sqrt{e}$. There exist $\lambda \in \Lambda[0, 1]$ and a price scenario $P = \mathcal{E}_\lambda(w)$ with $w \in X_{\lambda, 1, \sigma}[0, 1]$, $w(1) = 1$, such that the limit $\lim_{t \uparrow 1} \beta(t)$ does not exist, and the functions $\Phi'_t P$, $\Phi''_{xx} P$ are unbounded.*

Proof. To construct the desirable return w we choose a self-affine function of Example 3.6. By Theorem 1 of Kôno [58], a continuous self-affine function w on $[0, 1]$ with the scale parameter $H \in (0, 1)$ to base $r \geq 4$ satisfying $w(0) = 0$ and $w(1) = 1$ is completely determined by the sequence $x(k) \in \{1, -1\}$, $k = 0, \dots, r-1$, and such that $\sum_{k=0}^{r-1} x(k) = r^H$. Then for $0 \leq t \leq 1$,

$$w(t) = \sum_{n=1}^{\infty} y_{n-1}(t) s(\delta_n) r^{-nH},$$

where $t = \sum_{n=1}^{\infty} \delta_n r^{-n}$ with $\delta_n \in \{0, 1, \dots, r-1\}$, $s(0) = 0$, $s(j) = \sum_{i=0}^{j-1} x(i)$, $j = 1, \dots, r$, $y_0(t) = 1$ and

$$y_n(t) = \prod_{k=1}^n x(\delta_k). \tag{5.23}$$

For $m = 1, 2, \dots$,

$$0 \leq h = \sum_{n=m+1}^{\infty} \delta_n r^{-n} \leq r^{-mH} \quad \text{and} \quad t = \sum_{n=1}^m \delta_n r^{-n} = ir^{-m}, \quad (5.24)$$

letting $\epsilon_{i,m} := y_m(ir^{-m})$, we have

$$\begin{aligned} w(ir^{-m} + h) - w(ir^{-m}) &= \sum_{n=m+1}^{\infty} y_{n-1}(ir^{-m} + h)s(\delta_n)r^{-nH} \\ &= y_m(ir^{-m})r^{-mH} \sum_{n=1}^{\infty} y_{n-1}(r^m h)s(\delta_{m+n})r^{-nH} \\ &= \epsilon_{i,m}r^{-mH}w(r^m h). \end{aligned}$$

Now let w be such that $H = 1/2$, $r = 4$ and $\{x(k): k = 0, \dots, 3\} = \{1, 1, 1, -1\}$. For $h = r^{-m}$ with $m = 1, \dots$ and $t = 1 - h$ in (5.24), all $\delta_n = r - 1 = 3$ and $i = r^m - 1 = 3^m - 1$. Thus by (5.23), we have for $m = 1, 2, \dots$,

$$w(1) - w(1 - 4^{-m}) = (-1)^m 4^{-m/2} w(1). \quad (5.25)$$

Letting $\lambda_m := \{ir^{-m}: i = 0, \dots, r^m\}$, we have $\lambda = \{\lambda_m: m \geq 1\} \in \Lambda[0, 1]$, and so $w \in X_{\lambda, 1, \sigma}$. For the price scenario $P(t) = \exp\{w(t) - t/2\}$, $0 \leq t \leq 1$, we then have

$$\lim_{t \uparrow 1} d_1(t, P(t)) = \lim_{t \uparrow 1} \frac{\ln(P(t)/K)}{\sqrt{1-t}} = \lim_{h \downarrow 0} \frac{w(1-h) - w(1)}{\sqrt{h}}.$$

However by (5.25), the right side does not exist, proving the proposition. $\square$

5.4 Returns and the p -variation: examples

Let $0 < p < 2$ and let X be a stochastic process on a complete probability space $(\Omega, \mathcal{F}, \Pr)$ such that almost all its sample functions have bounded p -variation and are right-continuous. By Proposition 3.14 (see also Example 3.5), given $\lambda \in \Lambda[0, T]$, there is $\Omega_0 \in \mathcal{F}$ of full probability such that $X(\cdot, \omega)$ has the quadratic λ -variation for each $\omega \in \Omega_0$. Let τ be the first (random) moment in $(0, T]$ such that $X(\tau) - X(\tau-) \leq -1$, and let $X^{\tau-} := X(\cdot \wedge \tau-)$. Letting $X := \{X^{\tau-}(\cdot, \omega): \omega \in \Omega_0\}$ and $P_X := \{\mathcal{E}_\lambda(X^{\tau-}(\cdot, \omega)): \omega \in \Omega_0\}$, by Proposition 5, (P_X, X) is a λ -evolutionary system, that is, in the λ -evolutionary asset pricing model P_X is a set of price scenarios of a stock and X is a set of returns. So the quadratic λ -variation and the p -variation property are the main indicators of a stochastic process for its applicability in the λ -evolutionary asset pricing model.

The p -variation property has been investigated for several classes of stochastic processes. However, the results are scattered over many different journals. Below we provide some of these results related to financial models. More information can be found in the annotated reference list on p -variation in [23, Part IV].

Fractional Brownian motion. A fractional Brownian motion $B_H = \{B_H(t): t \geq 0\}$ with index $H \in (0, 1)$ is a mean zero Gaussian process with the covariance function

$$E\{B_H(t)B_H(s)\} = \frac{1}{2}\{t^{2H} + s^{2H} - |t - s|^{2H}\} \quad \text{for } t, s \geq 0 \quad (5.26)$$

and $B_H(0) = 0$ almost surely. Since the right side of (5.26) is equal to $t \wedge s$ for $H = 1/2$, B_H is a Brownian motion in this case. Many important properties of a fractional Brownian motion can be found in Kahane [53, Sect. 18]. In particular, from the a.s. bound $w_{B_H}(t) = O(t^H \sqrt{\log(1/t)})$ as $t \downarrow 0$ of the modulus of continuity on $[0, T]$ it follows that $v_p(B_H; [0, T]) < \infty$ a.s. whenever $p > 1/H$. The next statement provides a more precise description of the p -variation of a fractional Brownian motion. Let $s_\psi(f; \kappa) := \sum_{i=1}^n \psi(|f(x_i) - f(x_{i-1})|)$ for $\kappa = \{x_i: i = 0, \dots, n\}$ and let $\psi_H(u) := u^{1/H} / (2LLu)^{1/(2H)}$ for $u > 0$, where $LLu := \log_e |\log_e u|$, and $\psi_H(0) := 0$.

Proposition 15 *For $0 < T < \infty$, almost all sample functions of a fractional Brownian motion B_H with $H \in (0, 1)$ satisfy the relation*

$$\lim_{\delta \downarrow 0} \sup \{s_{\psi_H}(B_H; \kappa): \kappa \in \Xi[0, T] \text{ and the mesh } |\kappa| \leq \delta\} = T. \quad (5.27)$$

Remark 16 For the case $H = 1/2$, (5.27) was proved by Taylor [105, Theorem 1]. The general case of (5.27) with $\leq$ instead of $=$ follows from Theorem 3 of Kawada and Kôno [55] because almost all sample functions of B_H are continuous. The converse inequality does not follow from Theorem 4 of Kawada and Kôno [55] because their condition (v) fails to hold for $\sigma^2(u) = u^{2H}$, $u > 0$.

Proof. By the preceding remark, it is enough to prove (5.27) with $\geq$ instead of $=$. Let $\phi_H(u) := u^H \sqrt{2LLu}$ for $u > 0$. Oodaira [87] proved that $\eta_u := \{B_H(tu)/\phi_H(u): t \in [0, 1]\}$ satisfies the functional law of the iterated logarithm as $u \downarrow 0$. Taking a continuous functional $\Phi: C[0, 1] \rightarrow \mathbb{R}$ such that $\Phi(f) := f(1)$ for each $f \in C[0, 1]$, by the continuous mapping theorem, it then follows that, with probability 1,

$$\limsup_{u \downarrow 0} \frac{|B_H(u)|}{\phi_H(u)} = \sup \{f(1): f \in K_H\} \geq 1, \quad (5.28)$$

where K_H is the unit ball in the Reproducing Kernel Hilbert Space corresponding to the covariance of B_H . Denoting by ϕ_H^{-1} , the inverse function of ϕ_H , we have that $\psi_H(u)/\phi_H^{-1}(u) \rightarrow 1$ as $u \downarrow 0$. Thus, by (5.28), we have with probability 1

$$\limsup_{u \downarrow 0} \frac{\psi_H(|B_H(u)|)}{u} \geq 1.$$

This in conjunction with Fubini's theorem yields that the Lebesgue measure

$$|\{t \in [0, T]: \limsup_{u \downarrow 0} \frac{\psi_H(|B_H(t+u) - B_H(t)|)}{u} \geq 1\}| = T.$$

Now the proof can be completed using the Vitali covering theorem in the same way as for the case $H = 1/2$ in Taylor [105, Theorem 1]. $\square$

Statistical analysis of a financial data under the fractional Brownian motion hypothesis is discussed in Beran [3].

Ornstein-Uhlenbeck process. A mean zero Gaussian stochastic process $u = \{u(t): -\infty < t < +\infty\}$ with the covariance

$$E\{u(t)u(s)\} = \exp\{-\beta|t-s|\} \quad \text{for } t, s \in (-\infty, +\infty) \text{ and } \beta > 0$$

is called the OU (Ornstein-Uhlenbeck) process. It is the unique solution of the Langevin equation driven by a Brownian motion. Therefore u is also called the OU velocity process to distinguish it from the corresponding OU position process $x = \{x(t): t \geq 0\}$ defined by $x(t) - x(0) = \int_0^t u(s) ds$. The latter stochastic process models the x -coordinate of the free particle at the time t . The OU position process x is a mean zero Gaussian process with variance

$$E[x(t) - x(s)]^2 = 2\beta^{-2}[e^{-\beta|t-s|} - 1 + \beta|t-s|] \quad \text{for } t, s \geq 0.$$

When restricted to a bounded interval, almost all sample functions of x have bounded variation. This follows from the result of Ibragimov [51, Theorem 6]:

Theorem 17 *If $X = \{X(t): 0 \leq t \leq T\}$ is a separable measurable stochastic process such that $E|X(t) - X(s)| \leq K|t-s|$ for $0 \leq s, t \leq T$, then with probability 1, sample functions of X are of bounded variation.*

Goldenberg [45] argued that future prices in markets with frictions should be modelled by stochastic processes with smooth sample functions such as of an OU position process. Ross [99, Sect. 6.3.4] indicated a different financial application of an integrated Brownian motion having sample functions of bounded variation. Notice that the p -variation of an OU velocity process is the same as of a Brownian motion. The OU velocity process was scrutinized by Borell [10] as the return process on the basis of Danish stocks.

Homogeneous Lévy process. Let $X = \{X(t): t \geq 0\}$ be a separable, continuous in probability stochastic process with independent increments. Then X is called the *Lévy process* if almost all its sample functions are regulated, right continuous and $X(0) = 0$ almost surely. A Lévy process X is called *homogeneous* if a distribution of $X(t+s) - X(t)$ with $t, s \geq 0$ does not depend on t . Given a real number a , a positive number b , and a σ -finite measure L_X on $\mathbb{R} \setminus \{0\}$ such that $L_X(\{x \in \mathbb{R}: |x| > \delta\}) < \infty$ for $\delta > 0$ and $\int_{\mathbb{R} \setminus \{0\}} (1 \wedge x^2) L_X(dx) < \infty$, let

$$\Phi(u) := iau - bu^2 + \int_{\mathbb{R} \setminus \{0\}} (e^{iux} - 1 + iuh(x)) L_X(dx) \quad (5.29)$$

for $u \in \mathbb{R}$, where h is a bounded Borel function. The function Φ is called the *characteristic exponent* and the measure L_X is called the *Lévy measure*. Then the characteristic function of a homogeneous Lévy process X is given by $E \exp\{iuX(t)\} = \exp\{t\Phi(u)\}$ for each $t \geq 0$ and $u \in \mathbb{R}$.

It is well-known that sample functions of a homogeneous Lévy process X with the characteristic exponent (5.29) have bounded variation if and only if $b \equiv 0$ and

$$\int_{\mathbb{R} \setminus \{0\}} (1 \wedge |x|) L_X(dx) < \infty \quad (5.30)$$

(cf. e.g. Theorem 3 on p. 279 of Gikhman and Skorokhod [40]). The following result of Bretagnolle [12, Théorème III b] is less well-known:

Theorem 18 Let $1 < p < 2$ and let $X = \{X(t): t \geq 0\}$ be a mean zero homogeneous Lévy process with the characteristic exponent (5.29) such that $b \equiv 0$. Then $v_p(X; [0, 1]) < \infty$ with probability 1 if and only if

$$\int_{\mathbb{R} \setminus \{0\}} (1 \wedge |x|^p) L_X(dx) < \infty.$$

Less precise but more general result is a characterization of the p -variation index (see (4.6) for the definition) of sample functions of a Lévy process by the Blumenthal-Gettoor index. Let X be a homogeneous Lévy process. The *Blumenthal-Gettoor index* of X is defined by

$$\beta(X) := \inf \{ \alpha > 0: \int_{\mathbb{R} \setminus \{0\}} (1 \wedge |x|^\alpha) L_X(dx) < \infty \},$$

where L_X is the Lévy measure of X . Notice that $0 < \beta(X) \leq 2$. If X has no Gaussian part, then for any $0 < T < \infty$, the p -variation index $v(X; [0, T]) = \beta(X)$ almost surely. This follows from Theorems 4.1 and 4.2 of Blumenthal and Gettoor [9], and from Theorem 2 of Monroe [82].

Hyperbolic Lévy motion. A homogeneous Lévy process X is called a hyperbolic Lévy motion if in its characteristic exponent (5.29) $b \equiv 0$ and the Lévy measure $L_X(dx) = g(x; \alpha, \delta) dx$, where

$$g(x; \alpha, \delta) = \frac{1}{\pi^2 |x|} \int_0^\infty \frac{\exp\{-|x|\sqrt{2y + (\alpha)^2}\}}{y(J_1^2(\delta\sqrt{2y}) + Y_1^2(\delta\sqrt{2y}))} dy + \frac{\exp\{-|x|\}}{|x|}$$

for $\alpha > 0$ and $\delta \geq 0$. Here J_1 and Y_1 are the Bessel functions of the first and second kind, respectively. Eberlein and Keller [28] used a hyperbolic Lévy motion to model stock price returns and showed (ibid, p. 295) that $g(x; \alpha, \delta) \sim x^{-2}$ as $x \rightarrow 0$. Therefore, by Theorem 18, almost all sample functions of a hyperbolic Lévy motion have bounded p -variation on $[0, 1]$ for each $p > 1$, that is, the p -variation index $v(X; [0, 1]) = 1$.

Normal inverse Gaussian Lévy process. A homogeneous Lévy process X is called a normal inverse Gaussian Lévy process if in its characteristic exponent (5.29) $b \equiv 0$, $a = (2\alpha\delta/\pi) \int_0^1 \sinh(\beta x) K_1(\alpha x) dx$ and the Lévy measure $L_X(dx) = f(x; \alpha, \beta, \delta) dx$, where

$$f(x; \alpha, \beta, \delta) = \frac{\alpha\delta e^{-\beta x}}{\pi |x|} K_1(\alpha|x|) = \frac{\delta e^{\beta x}}{\pi\sqrt{2} |x|} \int_0^\infty y^{-1/2} e^{-|x|\sqrt{2y+\alpha^2}} dy$$

for $\alpha > 0$, $0 \leq |\beta| < \alpha$, $\mu \in \mathbb{R}$ and $\delta > 0$. Here K_1 denotes the modified Bessel function of third order and index 1. Barndorff-Nielsen [1] showed that $f(x; \alpha, \beta, \delta) \sim (\delta/\pi)x^{-2}$ as $x \rightarrow 0$. Therefore, by Theorem 18, almost all sample functions of a normal inverse Gaussian Lévy process have bounded p -variation on $[0, 1]$ for each $p > 1$, that is, the p -variation index $v(X; [0, 1]) = 1$.

The V. G. process. This process (variance gamma) is defined to be the composition $N(t) := B(G(t))$, $t \geq 0$, of a Brownian motion B and the process G which has independent gamma increments with mean t and variance νt . Then N is a homogeneous Lévy process with $a = b \equiv 0$ and with the Lévy measure $L_X(dx) = (\nu|x|)^{-1} \exp\{-|x|\sqrt{2}/\sigma\sqrt{\nu}\} dx$. Almost all sample functions of N have bounded variation because (5.30) holds. Madan and Seneta [72] introduced the V. G. process to replace a Brownian motion in models for stock price returns.

α -stable Lévy motion. A homogeneous Lévy process X is called an α -stable Lévy motion of index α if its characteristic exponent (5.29) is given by $b \equiv 0$ and the Lévy measure $L_X = L_{\alpha,r,q}$, where $L_{\alpha,r,q}(dx) := rx^{-1-\alpha}dx$ if $x > 0$ and $L_{\alpha,r,q}(dx) := q(-x)^{-1-\alpha}dx$ if $x < 0$ for $\alpha \in (0, 2)$ and $r, q \geq 0$ with $r + q > 0$. If $\alpha < 1$ then (5.30) holds and $\int h dL_X < \infty$. In this case, it is said that an α -stable Lévy motion has no drift if $a + \int h dL_X = 0$. The following result is due to Fristedt and Taylor [39, Theorem 2].

Theorem 19 *Let $X_\alpha = \{X_\alpha(t) : t \in [0, 1]\}$ be an α -stable Lévy motion with $\alpha \in (0, 2)$ having no drift if $\alpha < 1$ and with $r = q$ if $\alpha = 1$. For an increasing function $\psi : [0, \infty) \rightarrow [0, \infty)$, with probability 1*

$$\lim_{|\kappa| \rightarrow 0} s_\psi(X_\alpha; \kappa) \equiv \lim_{|\kappa| \rightarrow 0} \sum_{i=1}^n \psi(|X_\alpha(t_i) - X_\alpha(t_{i-1})|) = \sum_{(0,1]} \psi(|\Delta^- X_\alpha|). \quad (5.31)$$

To apply this result it is useful to recall that the right side of (5.31) is finite almost surely if and only if $\int_{-1}^1 \psi(u)u^{-1-\alpha} du < \infty$. Xu [111] established necessary and/or sufficient conditions for the boundedness of the p -variation of a symmetric α -stable processes with possibly dependent increments. More information about p -variation of stable processes can be found in Fristedt [38].

Multivariate stable distributions are often used as an alternative to the Gaussian hypothesis in financial models. Usually statistical analysis of financial data under the stable distribution hypothesis is based on the log-return. Therefore the conclusions of such analysis concern the exponential model. On the other hand, most theoretical work assuming the non-Gaussian stable distribution is based on the simple net return. This discord have been discussed by Elton et al. [29]. Financial applications of stable distributions are reviewed by McCulloch [76]. According to Campbell et al. [14, p. 18]: “Although stable distributions were popular in the 1960’s and early 1970’s, they are less commonly used today. They have fallen out of favor partly because they make theoretical modelling so difficult ...”.

5.5 Estimating the p -variation index

In the evolutionary asset pricing model a price scenario is represented by a function having the quadratic λ -variation with some $\lambda \in \Lambda[0, T]$. Therefore it is important to have a method to test this property when a function on $[0, T]$ is given by a finite set of values. This looks like a task comparable with the familiar problem of testing the hypothesis of finiteness of a second moment of a random variable having a finite sample. Instead, it seems simpler to start with a statistical analysis of whether a historical data with high probability correspond to a sample function of a stochastic process having bounded p -variation for some $p < 2$. This would be very usefull because a robustness of the evolutionary asset pricing model changes drastically if a price scenario follow a function having bounded p -variation for some $0 < p < 2$. Recall that the p -variation index of a function is the infimum of all p th such that the p -variation is finite (cf. (4.6)). In this section we discuss two methods of estimating the p -variation index of a function from a given finite set of its values.

Gladyshev class and the Orey index. A method of estimation of the p -variation index for a class of stochastic processes was suggested in [86]. It is based on the result of Gladyshev [43] concerning a rescaled quadratic variation property of a Gaussian stochastic process X with

stationary increments. Let $X = \{X(t): t \geq 0\}$ be a Gaussian stochastic process with stationary increments and continuous in quadratic mean. Let $\sigma = \sigma_X$ be the incremental variance of X given by

$$\sigma_X(u)^2 := E[(X(t+u) - X(t))^2], \quad t, u \geq 0.$$

Following Orey (1970), let

$$\gamma_* := \inf \{ \gamma > 0: u^\gamma / \sigma(u) \rightarrow 0, \text{ as } u \downarrow 0 \} = \limsup_{u \downarrow 0} \left(\frac{\log \sigma(u)}{\log u} \right)$$

and

$$\gamma^* := \sup \{ \gamma > 0: u^\gamma / \sigma(u) \rightarrow +\infty, \text{ as } u \downarrow 0 \} = \liminf_{u \downarrow 0} \left(\frac{\log \sigma(u)}{\log u} \right).$$

We always have that $0 \leq \gamma^* \leq \gamma_* \leq \infty$. If $\gamma_* = \gamma^*$ then we say that X has the *Orey index* $\gamma_X := \gamma_* = \gamma^*$. If X is a fractional Brownian motion B_H with the Hurst index H , then its Orey index $\gamma_X = H$. If X has the Orey index γ_X , then its p -variation index is $1/\gamma_X$, so that an estimation of the p -variation index reduces to an estimation of the Orey index.

For each integer $m \geq 1$, let $\lambda_m = \{i2^{-m}: i = 0, \dots, 2^m\}$ be a nested sequence of dyadic partitions of $[0, 1]$. Gladyshev [43] proved under mild conditions on X that if the Orey index γ_X exists then almost surely

$$\lim_{m \rightarrow \infty} \frac{\log \sqrt{s_2(X; \lambda_m)/2^m}}{\log(1/2^m)} = \gamma_X.$$

Motivated by Gladyshev's result, the following definition was suggested in [86].

Definition 20 Let $\{N_m: m \geq 1\}$ be a sequence of integers increasing to infinity and let $\lambda = \{\lambda_m: m \geq 1\}$ be a sequence of partitions $\lambda_m = \lambda(T/N_m) = \{iT/N_m: i = 0, \dots, N_m\}$ of $[0, T]$. We say that a stochastic process $X = \{X(t): t \geq 0\}$ belongs to the *Gladyshev class*, or *class $\mathcal{G}^\lambda$* whenever the limit

$$G_\lambda(X) := \lim_{m \rightarrow \infty} \frac{\log \sqrt{s_2(X; \lambda_m)/N_m}}{\log(1/N_m)} \quad (5.32)$$

exists for almost all sample functions of X .

We do not know how large Gladyshev's class is. If X is a Gaussian stochastic process, then some information can be derived from known results about the *strong limit theorem*: for some sequence $\{N_m: m \geq 1\}$ of integers increasing to infinity and for some constant C_1 , there exists the limit

$$\lim_{m \rightarrow \infty} \frac{s_2(X; \lambda_m)}{N_m \sigma_X(1/N_m)^2} = C_1^2, \quad \text{almost surely.} \quad (5.33)$$

Indeed for each $m \geq 1$ and any positive constant C_1 , we have the identity

$$\log \sqrt{\frac{s_2(X; \lambda_m)}{N_m}} = \log \sigma_X(1/N_m) + \log C_1 + \frac{1}{2} \log \left(1 + \frac{s_2(X; \lambda_m)}{C_1^2 N_m \sigma_X(1/N_m)^2} - 1 \right).$$

Then assuming that the Orey index γ_X exists, X belongs to the Gladyshev class $\mathcal{G}^\lambda$ and $\gamma_X = G_\lambda(X)$, almost surely. This relation was used in [86] to estimate the Orey index from a sample function of a stochastic process X given its values at finitely many points.

The rest contains a brief review of known results related to the strong limit theorem (5.33) for Gaussian processes. In addition to this theorem, there are several results in a literature concerning a restricted form of p -variation of stochastic processes. Corollary 22 below relates such results with the strong limit theorem used in the present paper. At the end of this section we formulate a part of Gladyshev's result concerning Gaussian stochastic processes with stationary increments.

For a Brownian motion B , Lévy [63], and independently Cameron and Martin [13], proved that the limit

$$\lim_{m \rightarrow \infty} s_2(B; \lambda(T/2^m)) = T \quad (5.34)$$

exists almost surely. Therefore the limit (5.32) with $X = B$ and $N_m = 2^m$ also exists almost surely and $G_\lambda(B) = 1/2$. Each Gaussian process X satisfying the strong limit theorem of Baxter [2] also belongs to Gladyshev's class and $G_\lambda(X) = 1/2$. Gladyshev [43] extended Baxter's result to a large class of Gaussian stochastic processes. In the case a Gaussian process X has stationary increments and has the Orey index $\gamma_X \in (0, 1)$, the limit

$$\lim_{m \rightarrow \infty} \frac{s_2(X; \lambda(T/2^m))}{Es_2(X; \lambda(T/2^m))} = T \quad (5.35)$$

exists almost surely provided the spectral density of X has a power behaviour (see Theorem 23 below for a precise statement). This yields (5.33) for $N_m = 2^m$. Yaglom [112] discussed several applications of this type strong limit theorems. He also suggested that relation (5.35), and its further modifications, should hold for a variety of stochastic processes.

Lévy [63] proved relation (5.34) for a sequence of dyadic partitions $\{kT2^{-n}: k = 0, \dots, 2^n\}$ of the time interval $[0, T]$. He also noticed that the same holds for any *nested* sequence $\{\kappa_n: n \geq 1\}$ of partitions κ_n such that $\cup_n \kappa_n$ is dense in $[0, T]$. Dudley [21] proved that the sequence of dyadic partitions in (5.34) can be replaced by an *arbitrarily* sequence of partitions $\kappa_n = \{0 = t_0^n < t_1^n < \dots < t_{N_n}^n = T\}$ as long as the mesh $|\kappa_n| := \max_i(t_i^n - t_{i-1}^n)$ tends to zero faster than $1/\log n$ as $n \rightarrow \infty$. The rate $o(1/\log n)$ cannot be improved as was shown by Fernández de la Vega [31]. Giné and Klein [42] extended the results of Baxter [2] and Gladyshev [43] to arbitrary sequences $\{\kappa_n: n \geq 1\}$ of partitions subject to a certain rate of convergence to zero of $|\kappa_n|$. For partitions $\kappa_n = \{kT/N_n: k = 0, \dots, N_n\}$, $n = 1, 2, \dots$, with equally spaced points we have $|\kappa_n| = 1/N_n$. It follows from Giné and Klein [42] that (5.35) holds provided $N_n^{-\gamma} = o(1/\log n)$ if $\gamma_X = \gamma \in (0, 1/2)$, $N_n^{-1} \log N_n = o(1/\log n)$ if $\gamma_X = 1/2$ and $N_n^{-1} = o(1/\log n)$ if $\gamma_X \in (1/2, 1)$. Recently Shao [102] proved that (5.35) does hold for Gaussian stochastic processes X with stationary increments such that σ_X is either convex or concave, and N_n satisfies conditions similar to the ones of Giné and Klein [42]. These conditions on σ_X in Shao's result are quite general and they do not seem to imply the existence of the Orey index γ_X .

Theorem 21 *Let $X = \{X(t): t \geq 0\}$ be a mean zero Gaussian stochastic process with stationary increments and continuous in quadratic mean. Suppose that for some $\gamma \in (0, 1)$, the Orey index $\gamma_X = \gamma$ and there exists the limit*

$$\lim_{m \rightarrow \infty} \frac{T/N_m}{\sigma_X(T/N_m)^{1/\gamma}} s_{1/\gamma}(X; \lambda(T/N_m)) = c \quad a.s. \quad (5.36)$$

for some sequence of integers $\{N_m: m \geq 1\}$ such that $N_m \rightarrow \infty$ as $m \rightarrow \infty$ and for some positive constants c, T . Then X belongs to the Gladyshev class $\mathcal{G}^\lambda$ and $G_\lambda(X) = \gamma$ almost surely.

Proof. For each integer $m \geq 1$, let $h_m := T/N_m$. First we prove that

$$\liminf_{m \rightarrow \infty} \frac{\log \sqrt{s_2(X; \lambda_m)/N_m}}{\log(1/N_m)} = \frac{1}{2} + \liminf_{m \rightarrow \infty} \frac{\log s_2(X; \lambda_m)}{2 \log h_m} \leq \gamma \quad \text{a.s..} \quad (5.37)$$

For strictly positive functions ϕ and ψ on $(0, 1]$ such that $\lim_{h \downarrow 0} \phi(h) = \lim_{h \downarrow 0} \psi(h) = 0$ the relation

$$\liminf_{h \downarrow 0} \frac{\log \phi(h)}{\log \psi(h)} = \sup \{ \alpha > 0 : \psi(h)^\alpha / \phi(h) \rightarrow +\infty \text{ as } h \downarrow 0 \}, \quad (5.38)$$

holds (see e.g. Annex A.4 in [106]). Therefore relation (5.37) holds if and only if

$$\sup \{ \alpha > 0 : h_m^{\alpha-1} s_2(X; \lambda_m) \rightarrow +\infty \text{ as } m \rightarrow \infty \} \geq 2 - 2\gamma \quad \text{a.s.,} \quad (5.39)$$

provided $s_2(X; \lambda_m)/h_m \rightarrow +\infty$ almost surely. Let $0 < \gamma'' < \gamma_X = \gamma < \gamma' \leq 1$. Since almost all sample functions of X satisfy Hölder's condition of order γ'' (see e.g. Section 9.4 in [15]), there is a positive random variable K such that

$$|X(ih_m) - X((i-1)h_m)| \leq Kh_m^{\gamma''} \quad (5.40)$$

for all $i = 1, \dots, N_m$ and $m \geq 1$. Denoting $\epsilon := \gamma''(1/\gamma - 1/\gamma') > 0$, we then have

$$s_{1/\gamma'}(X; \lambda_m) \geq K^{-\epsilon/\gamma''} \left[\frac{\sigma(h_m)}{h_m^{\gamma(1+\epsilon)}} \right]^{1/\gamma} \frac{h_m}{\sigma(h_m)^{1/\gamma}} s_{1/\gamma}(X; \lambda_m).$$

Since $\gamma(1+\epsilon) > \gamma = \gamma_*$, the quantity in square brackets above tends to infinity as $m \rightarrow \infty$. By assumption (5.36), it then follows that, for each $\gamma' \in (\gamma, 1]$,

$$\lim_{n \rightarrow \infty} s_{1/\gamma'}(X; \lambda_m) = +\infty \quad \text{a.s..}$$

To prove (5.39), first suppose that $1/2 \leq \gamma < 1$. For any $\gamma' \in (\gamma, 1]$, by Hölder's inequality, we have that

$$s_{1/\gamma'}(X; \lambda_m)^{2\gamma'} \leq N_m^{2\gamma'-1} s_2(X; \lambda_m)^2$$

for each $m \geq 1$. Thus for each $\gamma' \in (\gamma, 1]$,

$$\lim_{m \rightarrow \infty} h_m^{1-2\gamma'} s_2(X; \lambda_m) = +\infty \quad \text{a.s..} \quad (5.41)$$

This yields (5.39) in the case $1/2 \leq \gamma < 1$. Now suppose $0 < \gamma < 1/2$. For $\epsilon \in (0, 1/2 - \gamma)$, let $\gamma' := \gamma + \epsilon$ and $\gamma'' := \gamma - \epsilon$. Since almost all sample functions of X satisfy Hölder's condition (5.40), we have

$$s_{1/\gamma'}(X; \lambda_m) \leq K^{1/\gamma'-2} h_m^{\gamma''(1/\gamma'-2)} s_2(X; \lambda_m)$$

for each $m \geq 1$. Since $\gamma''(1/\gamma' - 2) = 1 - 2\gamma - 2\epsilon[1/(\gamma + \epsilon) - 1] < 1 - 2\gamma - \epsilon$, it follows that for an arbitrarily small $\epsilon > 0$,

$$\lim_{m \rightarrow \infty} h_m^{1-2\gamma-\epsilon} s_2(X; \lambda_m) = +\infty \quad \text{a.s..} \quad (5.42)$$

This yields (5.39) in the case $0 < \gamma < 1/2$. In both cases, by (5.41) and (5.42), we also have $s_2(X; \lambda_m)/h_m \rightarrow \infty$ almost surely. Therefore (5.39) implies (5.37).

To prove that $X \in \mathcal{G}^\lambda$ and $G_\lambda(X) = \gamma$ almost surely, by (5.37), it is enough to show that

$$\limsup_{m \rightarrow \infty} \frac{\log \sqrt{s_2(X; \lambda_m)/N_m}}{\log(1/N_m)} = \frac{1}{2} + \limsup_{m \rightarrow \infty} \frac{\log s_2(X; \lambda_m)}{2 \log h_m} \geq \gamma \quad \text{a.s.} \quad (5.43)$$

Under the same conditions as (5.38) we have the relation:

$$\limsup_{h \downarrow 0} \frac{\log \phi(h)}{\log \psi(h)} = \inf \{ \alpha > 0: \psi(h)^\alpha / \phi(h) \rightarrow 0 \text{ as } h \downarrow 0 \}. \quad (5.44)$$

Since $s_2(X; \lambda_m)/h_m \rightarrow \infty$ as $m \rightarrow \infty$ almost surely, (5.43) holds if and only if

$$\inf \{ \alpha > 0: h_m^{\alpha-1} s_2(X; \lambda_m) \rightarrow 0 \text{ as } m \rightarrow \infty \} \leq 2 - 2\gamma \quad \text{a.s.} \quad (5.45)$$

Let $0 < \gamma'' < \gamma$. Since almost all sample functions of X satisfy Hölder's condition (5.40), we have

$$s_{1/\gamma''}(X, \lambda_m) \leq K^{\epsilon/\gamma''} \left[\frac{\sigma(h_m)}{h_m^{\gamma(1-\epsilon)}} \right]^{1/\gamma} \frac{h_m}{\sigma(h_m)^{1/\gamma}} s_{1/\gamma}(X; \lambda_m)$$

for each $m \geq 1$, where $\epsilon := 1 - \gamma''/\gamma > 0$. Since $\gamma(1-\epsilon) < \gamma = \gamma^*$, the quantity in square brackets tends to zero as $m \rightarrow \infty$. By assumption (5.36), it then follows that for each $\gamma'' \in (0, \gamma)$,

$$\lim_{m \rightarrow \infty} s_{1/\gamma''}(X; \lambda_m) = 0 \quad \text{a.s.}$$

To prove (5.45), first suppose that $0 < \gamma \leq 1/2$. For any $\gamma'' \in (0, \gamma)$, by Hölder's inequality, we have that

$$s_2(X; \lambda_m) \leq N_m^{1-2\gamma''} s_{1/\gamma''}(X; \lambda_m)^{2\gamma''}$$

for each $m \geq 1$. Thus for each $\gamma'' \in (0, \gamma)$,

$$\lim_{m \rightarrow \infty} h_m^{1-2\gamma''} s_2(X; \lambda_m) = 0 \quad \text{a.s.} \quad (5.46)$$

This yields (5.45) in the case $0 < \gamma \leq 1/2$. Now suppose that $1/2 < \gamma < 1$. Let $\gamma'' \in (1/2, \gamma)$. Since almost all sample functions of X satisfy Hölder's condition (5.40), we have

$$s_2(X; \lambda_m) \leq K^{2-1/\gamma''} h_m^{2\gamma''-1} s_{1/\gamma''}(X; \lambda_m)$$

for each $m \geq 1$. Thus (5.46) holds for each $\gamma'' \in (1/2, \gamma)$. This yields (5.45) in the case $1/2 < \gamma < 1$. Since (5.45) implies (5.43), the proof of Theorem 21 is complete. $\square$

Corollary 22 *Let $X = \{X(t): t \geq 0\}$ be a mean zero Gaussian stochastic process with stationary increments and continuous in quadratic mean. Suppose that for some $\gamma \in (0, 1)$,*

- (a) *the incremental variance σ_X of X satisfies the condition $\sigma_X(h) = C_1 h^\gamma (1 + \epsilon(h))$ where C_1 is a positive constant and $\epsilon(h) \rightarrow 0$ as $h \downarrow 0$;*
- (b) *there exists the limit*

$$\lim_{m \rightarrow \infty} s_{1/\gamma}(X; \lambda(T/N_m)) = C_2 \quad \text{a.s.} \quad (5.47)$$

for some sequence of integers $\{N_m: m \geq 1\}$ such that $N_m \rightarrow \infty$ as $m \rightarrow \infty$ and some positive constants C_2, T .

Then the Orey index $\gamma_X = \gamma$, X belongs to the Gladyshev class $\mathcal{G}^\lambda$ and $G_\lambda(X) = \gamma$ almost surely, where $\lambda = \{\lambda(T/N_m): m \geq 1\}$.

Conditions for (5.47) to hold are given by Marcus and Rosen [74] in the case $0 < \gamma \leq 1/2$, and by Shao [102] in the case $0 < \gamma < 1$. Let X satisfy the assumptions of Corollary 22 except possibly (5.47) and let $0 < T < \infty$. Suppose that σ_X^2 is nondecreasing and concave on $[0, T]$ if $0 < \gamma \leq 1/2$, and that σ_X^2 is nondecreasing and convex on $[0, T + \epsilon]$ for some $\epsilon > 0$ if $1/2 < \gamma < 1$. Also, suppose that $1/N_m = o(1/(\log m)^a)$ as $m \rightarrow \infty$, where $a := 1/\{2[\gamma \wedge (1 - \gamma)]\}$. Then by Corollaries 1.1 and 1.2 of Shao [102], (5.47) holds with $C_2 = TC_1^{1/\gamma} E|\eta|^{1/\gamma}$, where η is a standard normal random variable and C_1 is the constant from (a).

Further conditions on existence of the Orey index will be given in terms of a spectral density of a Gaussian stochastic process with stationary increments. A nondecreasing function F on $\mathbb{R} \setminus \{0\}$ is called a Lévy-Khinchin function if

$$\int_{\mathbb{R} \setminus \{0\}} \min\{1, u^2\} dF(u) < \infty.$$

Let X be a real-valued mean zero Gaussian stochastic process with stationary increments, continuous in probability and $X(0) = 0$ almost surely. Then there is a unique Lévy-Khinchin function F_X on $\mathbb{R} \setminus \{0\}$ such that

$$E[X(t)X(s)] = \int_{\mathbb{R} \setminus \{0\}} (e^{it\lambda} - 1)(e^{-is\lambda} - 1) dF_X(\lambda).$$

If F_X is absolutely continuous, the derivative $f_X = F'_X$ is called the spectral density of X .

The spectral density of a fractional Brownian motion B_H with the Hurst exponent $0 < H < 1$ is given by the power function $f_H(\lambda) = c_H |\lambda|^{-1-2H}$, where $c_H = (2\pi)^{-1} \sin(\pi H) \Gamma(1+2H)$. This follows from the relation:

$$\sigma_{B_H}(u)^2 = \int_{-\infty}^{\infty} |e^{iu\lambda} - 1|^2 f_H(\lambda) d\lambda = 8c_H u^{2H} \int_0^{\infty} \left(\sin \frac{\lambda}{2}\right)^2 \frac{d\lambda}{\lambda^{1+2H}} = u^{2H}.$$

So that the Orey index γ_{B_H} exists and equals H .

A spectral density which differ from the spectral density f_H by a slowly varying function gives rise to a Gaussian stochastic process with the same Orey index H . Consider a quasi-monotone slowly varying (near infinity) function ℓ defined in [8, Section 2.7] and let $f_{\gamma, \ell}(\lambda) := \ell(|\lambda|) |\lambda|^{-1-2\gamma}$ for some $0 < \gamma < 1$. By Abelian Theorem 4.1.5 from [8], we have

$$\int_{-\infty}^{\infty} |e^{iu\lambda} - 1|^2 f_{\gamma, \ell}(\lambda) d\lambda = 8u^{2\gamma} \int_0^{\infty} \ell\left(\frac{\lambda}{u}\right) \left(\sin \frac{\lambda}{2}\right)^2 \frac{d\lambda}{\lambda^{1+2\gamma}} \sim \ell\left(\frac{1}{u}\right) u^{2\gamma} c_\gamma^{-1} \quad (5.48)$$

as $u \downarrow 0$, where $\phi(u) \sim \psi(u)$ as $u \downarrow 0$ means that $\lim_{u \downarrow 0} \phi(u)/\psi(u) = 1$. Let X be a Gaussian stochastic process with spectral density $f_{\gamma, \ell}$. By elementary properties of slowly varying functions (e.g. Proposition 1.3.6 in [8]), it follows that the Orey index γ_X exists and equals γ for any quasi-monotone slowly varying function ℓ .

The following statement contains a part of Theorem 3 of Gladyshev [43].

Theorem 23 *Let X be a mean zero Gaussian stochastic process with stationary increments having the spectral density f_X such that, for some $0 < \gamma < 1$,*

$$f_X(\lambda) = |\lambda|^{-1-2\gamma} [\ell(|\lambda|) + b(\lambda)],$$

where ℓ is a quasi-monotone slowly varying function, b is zero on a neighbourhood of zero, bounded elsewhere and $b(\lambda) = O(\lambda^{-2})$ as $\lambda \rightarrow \infty$. Then the Orey index $\gamma_X = \gamma$. If in addition $\ell(|\lambda|) = c$ for some $c > 0$, then

$$\frac{1}{2} - \lim_{m \rightarrow \infty} \frac{\log s_2(X, \lambda(2^{-m}))}{2m} = \gamma \quad a.s..$$

Proof. Suppose that b is zero on $[-1, 1]$. Then for some constant C depending on b , and for each $u > 0$, we have

$$\begin{aligned} \int_{-\infty}^{\infty} \left(\sin \frac{u\lambda}{2} \right)^2 \frac{b(\lambda) d\lambda}{|\lambda|^{1+2\gamma}} &\leq C u^{2+2\gamma} \int_u^{\infty} \left(\sin \frac{\lambda}{2} \right)^2 \frac{d\lambda}{\lambda^{3+2\gamma}} \\ &\leq C u^{2\gamma} \int_u^{u^{1/(2+2\gamma)}} \left(\sin \frac{\lambda}{2} \right)^2 \frac{d\lambda}{\lambda^{1+2\gamma}} + C u^{1+2\gamma} / (2 + 2\gamma). \end{aligned} \quad (5.49)$$

Since the right side of (5.49) is of order $o(u^{2\gamma})$ as $u \downarrow 0$, by (5.48), it follows that $\sigma_X(u) \sim \sqrt{c_\gamma^{-1} \ell(1/u)} u^\gamma$ as $u \downarrow 0$. Therefore the Orey index γ_X exists and equals γ , proving the first part of the theorem. The second part follows from Theorem 3 of Gladyshev [43] with $\alpha = 1 + 2\gamma$. $\square$

Appendix A

Convergence of directed functions

In this paper, the Moore-Smith limit theory as developed by McShane [80] (see also [81]) is used to define the convergence with respect to refinements of partitions. Namely, McShane's notion of the direction is used to define several Stieltjes type integrals, the unconditional convergence of sums and products, and the local p -variation. For readers convenience we state the main facts about directed functions in this appendix. A connection with the better known theories of limits based on nets and filters is discussed in the last section of [80].

Directed functions. We start with the notion of a directed function and its convergence which are basic to McShane's limit theory (see p. 10 in [80], or p. 33 in [81]).

Definition 1 Let D be a set.

- (a) A nonempty family $\mathfrak{U}$ of nonempty subsets of D is called a *direction in D* if it is directed downward by inclusion, that is, if A_1 and A_2 belong to $\mathfrak{U}$, then there is a set A_3 in $\mathfrak{U}$ such that $A_3 \subset A_1 \cap A_2$.
- (b) Let $\mathfrak{U}$ be a direction in D . If $P(x)$ is a property valid for $x \in D$ then we say that “*ultimately $P(x)$* ” (or “ $P(x)$ is ultimately true”) provided that there is a set A in $\mathfrak{U}$ such that $P(x)$ holds for each $x \in A$.
- (c) Let $\mathfrak{U}$ be a direction in D . If a real-valued function f is ultimately defined on D , that is ultimately “ $f(x)$, $x \in D$, is a real number”, then the ordered pair $(f, \mathfrak{U})$ is called a *directed function*. A directed function $(f, \mathfrak{U})$ converges to a real number r , in symbols

$$\lim_{\mathfrak{U}} f = \lim_{x, \mathfrak{U}} f(x) = r,$$

if for each neighborhood V of r , f is ultimately in V .

The convergence of directed functions satisfies all natural properties of limits including the uniqueness of the limit (p. 12 in [80]). Next is the Cauchy test for the convergence of directed functions (see p. 37 in [81] for the proof).

Theorem 2 Let f be a real-valued function defined on a set D , and let $\mathfrak{U}$ be a direction on D . The directed function $(f, \mathfrak{U})$ has a limit if and only if for each $\epsilon > 0$ there exists $A \in \mathfrak{U}$ such that $|f(x) - f(y)| < \epsilon$ for each $x, y \in A$.

Order convergence. Let $(f, \mathfrak{U})$ be a real-valued directed function. Define the *upper* and *lower limits* of $(f, \mathfrak{U})$, respectively, by

$$\limsup_{\mathfrak{U}} f := \inf_{A \in \mathfrak{U}} \sup_{x \in A} f(x) \quad \text{and} \quad \liminf_{\mathfrak{U}} f := \sup_{A \in \mathfrak{U}} \inf_{x \in A} f(x). \quad (\text{A.1})$$

Lemma 3 For a directed function $(f, \mathfrak{U})$, $\liminf_{\mathfrak{U}} f \leq \limsup_{\mathfrak{U}} f$.

A real-valued directed function $(f, \mathfrak{U})$ is *order convergent* to $b \in \mathbb{R}$ if

$$\liminf_{\mathfrak{U}} f = \limsup_{\mathfrak{U}} f = b$$

The following theorem is a special case of Theorem 11.4 of McShane and Botts [81, p. 55] proved for functions with values in the extended set of real numbers.

Theorem 4 Let $(f, \mathfrak{U})$ be a real-valued directed function and let $b \in \mathbb{R}$. Then $(f, \mathfrak{U})$ is order convergent to b if and only if it converges to b .

We use this theorem in Section 2.2 to characterize the Wiener class of functions having the local p -variation, and at the end of Section 2.3 to relate several definitions of extended Riemann-Stieltjes integrals.

Unconditionally convergent sums. Suppose that f is a real-valued function defined on an interval J . Let $\mathcal{F}(J)$ be the family of all *finite* sets of points of J . For each $\sigma \in \mathcal{F}(J)$, let

$$S(f; \sigma) := \sum_{x \in \sigma} f(x).$$

The *sum-function* $S(f) = \{S(f; \sigma) : \sigma \in \mathcal{F}(J)\}$ is then defined on $\mathcal{F}(J)$. For each finite subset σ of J , let $A(\sigma)$ be the set of all finite subsets σ' of J which contain σ . Then $A(\sigma)$ is a nonempty subset in the domain of the sum-function S . Let $\mathfrak{F}$ be the family of all sets $A(\sigma)$ for $\sigma \in \mathcal{F}(J)$, that is

$$\mathfrak{F} = \mathfrak{F}(J) := \{A(\sigma) : \sigma \in \mathcal{F}(J)\}. \quad (\text{A.2})$$

Let $A(\sigma_1)$ and $A(\sigma_2)$ be any two members of $\mathfrak{F}$. Then $A(\sigma_1 \cup \sigma_2)$ belongs to $\mathfrak{F}$, and every finite set σ of J which contains $\sigma_1 \cup \sigma_2$ also contains both σ_1 and σ_2 , that is, σ belongs to $A(\sigma_1) \cap A(\sigma_2)$. Hence $\mathfrak{F}$ is a direction in $\mathcal{F}(J)$.

Definition 5 Let f be a real-valued function defined on an interval J . We say that the *non-ordered sum* $\sum_J f$ *converges unconditionally* if the directed function $(S(f), \mathfrak{F})$ has a limit and let

$$\sum_J f := \lim_{\sigma, \mathfrak{F}} S(f; \sigma).$$

Also we say that the *non-ordered sum* $\sum_J f$ *converges absolutely* if there exists K such that $\sum_{x \in \sigma} |f(x)| \leq K$ for each $\sigma \in \mathcal{F}(J)$.

Theorem 6 For a real-valued function f on an interval J , the non-ordered sum $\sum_J f$ converges unconditionally if and only if it converges absolutely.

Proof. First, suppose that f is non-negative on J . In this case, the directed function $(S(f), \mathfrak{F})$ converges if and only if the set $\{S(f; \sigma) : \sigma \in \mathcal{F}(J)\}$ has an upper bound, and if both statements hold then the limit is the least upper bound. Now, suppose that f may have arbitrary real values. Let f^+ and f^- be positive and negative parts of f , respectively, so that $f = f^+ - f^-$ and $|f| = f^+ + f^-$. If $\{S(|f|; \sigma) : \sigma \in \mathcal{F}(J)\}$ is bounded then also the sets $\{S(f^+; \sigma) : \sigma \in \mathcal{F}(J)\}$ and $\{S(f^-; \sigma) : \sigma \in \mathcal{F}(J)\}$ are bounded. By the first part, the directed functions $(S(f^+), \mathfrak{F})$ and $(S(f^-), \mathfrak{F})$ converge. Hence by linearity of the limit, the sum $\sum_J f$ converges unconditionally. For the converse implication we use the Cauchy test (Theorem 2 above) to show that if the sum $\sum_J f$ converges unconditionally then so do the sums $\sum_J f^+$ and $\sum_J f^-$. Indeed, for $k = 1, 2$ and for any finite sets $\sigma_0 \subset \sigma_k$, let

$$\sigma_0 \subset \sigma'_k := \sigma_0 \cup \{x \in \sigma_k \setminus \sigma_0 : f(x) > 0\} \subset \sigma_k.$$

Then $S(f^+; \sigma_1) - S(f^+; \sigma_2) = S(f; \sigma'_1) - S(f; \sigma'_2)$. A similar relation holds between f and f^- . This implies the claim. By the above argument it follows that the set $\{S(|f|; \sigma) : \sigma \in \mathcal{F}(J)\}$ is bounded. The proof of Theorem 6 is complete. $\square$

For a function f with real values, by the preceding theorem, the sum $\sum_J f$ converges unconditionally if and only if it converges absolutely. This is still true for functions with values in any finite-dimensional Banach space. In an infinite-dimensional Banach space, absolute convergence implies unconditional convergence but unconditional convergence never implies absolute convergence (Dvoretzky and Rogers [26]).

Theorem 7 *Let J, J_1, J_2 be intervals such that $J = J_1 \cup J_2$ and $J_1 \cap J_2 = \emptyset$. Then the non-ordered sum $\sum_J f$ converges unconditionally if and only if both $\sum_{J_1} f$ and $\sum_{J_2} f$ converge unconditionally, and then*

$$\sum_J f = \sum_{J_1} f + \sum_{J_2} f. \quad (\text{A.3})$$

Proof. Suppose that the non-ordered sum $\sum_J f$ converges unconditionally. By the preceding theorem, the set $\{S(|f|; \sigma) : \sigma \in \mathcal{F}(J)\}$ is bounded. Therefore the same sets with $\mathcal{F}(J)$ replaced by $\mathcal{F}(J_1)$ or $\mathcal{F}(J_2)$ also are bounded, and hence the sums on the right side of (A.3) converge unconditionally. The converse implication follows similarly. The conclusion of the theorem now follows by linearity of the limit from equality (A.3) with J, J_1, J_2 replaced by finite sets $\sigma \subset J, \sigma_1 \subset J_1, \sigma_2 \subset J_2$ such that $\sigma = \sigma_1 \cup \sigma_2$. $\square$

Unconditionally convergent products. Here we extend the concept of unconditional convergence to non-ordered products. Again let f be a real-valued function defined on an interval J , and let $\mathcal{F}(J)$ be the family of all finite sets of points of J . For each $\sigma \in \mathcal{F}(J)$, let

$$P(f; \sigma) := \prod_{x \in \sigma} f(x).$$

The *product-function* $P(f) = \{P(f; \sigma) : \sigma \in \mathcal{F}(J)\}$ is then defined on $\mathcal{F}(J)$. Let $\mathfrak{F}$ be the direction on $\mathcal{F}(J)$ defined by (A.2).

Definition 8 Let f be a real-valued function defined on an interval J . We say that the *non-ordered product* $\prod_J f$ converges unconditionally if the directed function $(P(f), \mathfrak{F})$ has a *non-zero* limit, and then let

$$\prod_J f := \lim_{\sigma, \mathfrak{F}} P(f; \sigma).$$

If f is positive everywhere on J then $\log f(x)$ is defined for each $x \in J$. Also, if in this case the non-ordered product $\prod_J f$ converges then the directed function $(S(\log f), \mathfrak{F})$ has a limit $\sum_J \log f$ such that

$$\log \left(\prod_J f \right) = \sum_J \log f. \quad (\text{A.4})$$

We show below that the unconditional convergence of a non-ordered sum $\sum_J f$ is equivalent to the unconditional convergence of a non-ordered product $\prod_J f$. Thus the unconditional convergence of non-ordered products extends the absolute convergence of products of sequences.

Proposition 9 *For a real-valued function f on J , if the non-ordered product $\prod_J f$ converges unconditionally then for each $\epsilon > 0$, ultimately $|f - 1| < \epsilon$.*

Proof. Let $P = \prod_J f$, and let $\epsilon \in (0, |P|)$. Then there is $\sigma_0 \in \mathcal{F}(J)$ such that $|P(f; \sigma) - P| < \epsilon$ for each $\sigma \supset \sigma_0$. For $x \in J \setminus \sigma_0$, we then have

$$|f(x) - 1| = \frac{|P(f; \sigma_0 \cup \{x\}) - P(f; \sigma_0)|}{|P(f; \sigma_0)|} < \frac{2\epsilon}{|P| - \epsilon}.$$

Since ϵ is arbitrary, the conclusion follows. $\square$

It follows from the preceding proposition that f is equal 1 everywhere on J except on a countable set, and f is positive on J except on a finite set. In particular, the sum-function $S(\log f)$ is ultimately defined on $\mathcal{F}(J)$ with respect to the direction $\mathfrak{F}$, whenever the non-ordered product $\prod_J f$ converges unconditionally. Then the same arguments used to show (A.4) for a positive f , yield the following statement:

Theorem 10 *Let f be a real-valued function defined on an interval J . The non-ordered product $\prod_J f$ converges unconditionally if and only if the non-ordered sum $\sum_J \log f$ converges unconditionally.*

The following criterion for the unconditional convergence of non-ordered products is used in Lemma 3.37.

Theorem 11 *Let ξ be a real-valued function on J such that $\xi(x) \neq -1$ for each $x \in J$. Then the non-ordered product $\prod_J (1 + \xi)$ converges unconditionally if and only if the non-ordered sum $\sum_J \xi$ converges unconditionally.*

Proof. Suppose that the non-ordered product $\prod_J (1 + \xi)$ converges unconditionally. By Proposition 9, for each $\epsilon > 0$, ultimately $|\xi| < \epsilon$. Therefore ultimately $|1 - \xi^{-1} \log(1 + \xi)| < \epsilon$. By Theorem 10, it then follows that the directed function $(S(\xi), \mathfrak{F})$ has a limit proving the unconditional convergence of the non-ordered sum $\sum_J \xi$. The converse implication follows similarly because a limit of the directed function $(P(1 + \xi), \mathfrak{F})$ is non-zero. $\square$

Bibliography

- [1] O. E. BARNDORFF-NIELSEN, Normal inverse Gaussian distributions and stochastic volatility modelling. *Scand. J. Statist.*, **24** (1997), 1-14.
- [2] G. BAXTER, A strong limit theorem for Gaussian processes. *Proc. Amer. Math. Soc.*, **7** (1956), 522-527.
- [3] J. BERAN, *Statistics for long-memory processes*. New-York, Chapman & Hall, 1994.
- [4] S. K. BERBERIAN, The character space of the algebra of regulated functions. *Pacific J. Math.*, **74** (1978), 15-36.
- [5] P. L. BERNSTEIN, *Capital Ideas. The Improbable Origins of Modern Wall Street*. New York, The Free Press, 1992.
- [6] J. BERTOIN, Sur la mesure d'occupation d'une classe de fonctions self-affines. *Japan J. Appl. Math.*, **5** (1988), 431-439.
- [7] A. BICK AND W. WILLINGER, Dynamic spanning without probabilities. *Stoch. Proc. Appl.*, **50** (1994), 349-374.
- [8] N. H. BINGHAM, C. M. GOLDIE AND J. L. TEUGELS, *Regular variation*. Cambridge, Cambridge University Press, 1987.
- [9] R. M. BLUMENTHAL AND R. K. GETTOOR, Sample functions of stochastic processes with stationary independent increments. *J. Math. and Mech.*, **10** (1961), 493-516.
- [10] CH. BORELL, Analytic and empirical evidences of isoperimetric processes. In: *Probability in Banach Spaces 6*, U. Haagerup et. al. Eds., Progress in Probability **20**, 13-40. Boston, Birkhäuser, 1999.
- [11] N. BOULEAU AND D. LÉPINGLE, *Numerical Methods for Stochastic Processes*. New York, Wiley & Sons, 1994.
- [12] J. L. BRETAGNOLE, p -variation de fonctions aléatoires. *Lect. Notes Math.* (Springer), **258** (1972), 51-71.
- [13] R. H. CAMERON AND W. T. MARTIN, The behavior of measure and measurability under a change of scale in Wiener space. *Bull. Amer. Math. Soc.*, **53** (1947), 130-137.
- [14] J. Y. CAMPBELL, A. W. LO AND A. C. MACKINLAY, *The Econometrics of Financial Markets*. Princeton, Princeton University Press, 1997.

- [15] H. CRAMÉR AND M. R. LEADBETTER, *Stationary and Related Stochastic Processes*. New York, John Wiley & Sons, 1967.
- [16] C. DELLACHERIE AND P.-A. MEYER, *Probabilities and Potential B*. Transl. by J. P. Wilson. Amsterdam, North-Holland, 1982.
- [17] R. L. DOBRUSHIN, Generalization of Kolmogorov's equations for Markov processes with a finite number of possible states. *Mat. Sb.*, **33** (1953), 567-596 (in Russian).
- [18] C. DOLÉANS-DADE, Quelques applications de la formule de changement de variables pour les semimartingales. *Z. Wahrsch. verw. Geb.*, **16** (1970), 181-194.
- [19] J. D. DOLLARD AND C. N. FRIEDMAN, *Product integration with applications to differential equations*. Encyclopedia of Mathematics and Its Applications, **10**. London, Addison-Wesley, 1979.
- [20] J. L. DOOB, *Stochastic Processes*. New York, Wiley, 1953.
- [21] R. M. DUDLEY, Sample functions of the Gaussian process. *Ann. Probab.*, **1** (1973), 66-103.
- [22] R. M. DUDLEY, Fréchet differentiability, p -variation and uniform Donsker classes. *Ann. Probab.*, **20** (1992), 1968-1982.
- [23] R. M. DUDLEY AND R. NORVAIŠA, *Differentiability of Six Operators on Nonsmooth Functions and p -Variation*. Lect. Notes in Math., **1703**. Berlin, Springer, 1999.
- [24] R. M. DUDLEY AND R. NORVAIŠA, An introduction to p -variation and Young integrals. Lect. Notes, **1**. Aarhus Univ., 1998.
- [25] R. M. DUDLEY AND R. NORVAIŠA, *Concrete Functional Calculus*. Manuscript.
- [26] A. DVORETZKY AND C. A. ROGERS, Absolute and unconditional convergence in normed linear spaces. *Proc. Nat. Acad. Sci. U.S.A.*, **36** (1950), 192-197.
- [27] A. M. D'YAČKOV, Conditions for the existence of Stieltjes integral of bounded generalized variation. *Anal. Math.* (Budapest), **14** (1988), 295-313.
- [28] E. EBERLEIN AND U. KELLER, Hyperbolic distributions in finance. *Bernoulli*, **1** (1995), 281-299.
- [29] E. J. ELTON, M. J. GRUBER AND P. R. KLEINDORFER, A closer look at the implications of the stable paretian hypothesis. *Review of Economics and Statistics*, (1975), 231-235.
- [30] K.-J. ENGEL AND R. NAGEL, *One-Parameter Semigroups for Linear Evolution Equations*. New York, Springer, 2000.
- [31] W. FERNÁNDEZ DE LA VEGA, On almost sure convergence of quadratic Brownian variation. *Ann. Probab.*, **2** (1974), 551-552.
- [32] B. DE FINETTI AND M. JACOB, Sull' integrale di Stieltjes-Riemann. *Giorn. Istituto Ital. Attuari*, **6** (1935), 303-319.
- [33] D. L. FISK, Quasi-martingales. *Trans. Amer. Math. Soc.*, **120** (1965), 369-389.

- [34] H. FÖLLMER, Calcul d'Itô sans probabilités. *Séminaire de Probabilités XV*, Eds. J. Azéma and M. Yor; *Lect. Notes in Math.* (Springer) **850** (1981), 143-150.
- [35] H. FÖLLMER, Dirichlet processes. In: *Stochastic Integrals; Lect. Notes in Math.* (Springer) **851** (1981), 476-478.
- [36] H. FÖLLMER, CH.-T. WU AND M. YOR, On weak brownian motion of arbitrary order. *Ann. Inst. H. Poincaré, Probab. et Statist.*, **36** (2000), 447-487.
- [37] M. A. FREEDMAN, Operators of p -variation and the evolution representation problem. *Trans. Amer. Math. Soc.*, **279** (1983), 95-112.
- [38] B. FRISTEDT, Sample functions of stochastic processes with stationary, independent increments. In: *Advances in Probability and Related Topics*, Vol. 3, 241-396. New York, Dekker, 1974.
- [39] B. FRISTEDT AND J. TAYLOR, Strong variation for the sample functions of a stable process. *Duke Math. J.*, **40** (1973), 259-278.
- [40] I. I. GIHMAN AND A. V. SKOROHOD, *Introduction to the theory of random processes*. Philadelphia, W.B. Saunders, 1969.
- [41] I. I. GIHMAN AND A. V. SKOROHOD, *The Theory of Stochastic Processes*. Vol. I. New York, Springer, 1974.
- [42] E. GINÉ AND R. KLEIN, On quadratic variation of processes with Gaussian increments. *Ann. Probab.*, **3** (1975), 716-721.
- [43] E. G. GLADYSHEV, A new limit theorem for stochastic processes with Gaussian increments. *Theor. Probability Appl.*, **6** (1961), 52-61.
- [44] V. I. GLIVENKO, *The Stieltjes integral*. Leningrad, ONTI, 1936 (in Russian).
- [45] D. H. GOLDENBERG, Sample path properties of futures prices. *The Journal of Futures Markets*, **6** (1986), 127-140.
- [46] L. M. GRAVES, *The Theory of Functions of Real Variable*. New York, McGraw-Hill Book Company, 1946.
- [47] R. HENSTOCK, *Theory of integration*. London, Butterworths, 1963.
- [48] T. H. HILDEBRANDT, Definitions of Stieltjes integrals of the Riemann type. *Amer. Math. Monthly*, **45** (1938), 265-278.
- [49] T. H. HILDEBRANDT, *Introduction to the Theory of Integration*. New York, Academic Press, 1963.
- [50] G. A. HUNT, Random Fourier series. *Trans. Amer. Math. Soc.*, **71** (1951), 38-69.
- [51] I. A. IBRAGIMOV, Properties of sample functions for stochastic processes and embedding theorems. *Theory Probab. Appl.*, **18** (1973), 442-453.
- [52] N. C. JAIN AND D. MONRAD, Gaussian measures in B_p . *Ann. Probab.*, **11** (1983), 46-57.

- [53] J.-P. KAHANE, *Some random series of functions*. Second ed. Cambridge University Press, 1985.
- [54] T. KAMAE, A characterization of self-affine functions. *Japan J. Appl. Math.*, **3** (1986), 271-280.
- [55] T. KAWADA AND N. KÔNO, On the variation of Gaussian processes. In: *Proc. of the Second Japan-USSR Symposium on Probability Theory*, Lect. Notes Math. (Springer), **330** (1973), 176-192.
- [56] F. B. KNIGHT, On the random walk and Brownian motion. *Trans. Amer. Math. Soc.*, **103** (1962), 218-228.
- [57] N. KÔNO, On self-affine functions. *Japan J. Appl. Math.*, **3** (1986), 259-269.
- [58] N. KÔNO, On self-affine functions II. *Japan J. Appl. Math.*, **5** (1988), 441-454.
- [59] H. KUNITA AND S. WATANABE, On square integrable martingales. *Nagoya Math. J.*, **30** (1967), 209-245.
- [60] J. KURZWEIL, Generalized ordinary differential equations and continuous dependence on a parameter. *Czechoslovak Math. J.*, **7** (1957), 568-583 (in Russian).
- [61] S. KWAPIEŃ AND W. A. WOYCZYŃSKI, *Random Series and Stochastic Integrals: Single and Multiple*. Boston, Birkhäuser, 1992.
- [62] D. LÉPINGLE, La variation d'ordre p des semi-martingales. *Z. Wahrsch. verw. Geb.*, **36** (1976), 295-316.
- [63] P. LÉVY, Le mouvement brownien plan. *Amer. J. Math.*, **62** (1940), 487-550.
- [64] P. LÉVY, *Random functions: general theory with special reference to Laplacian random functions*. University California Press, Berkeley, 1953.
- [65] E. R. LOVE, The refinement-Ross-Riemann-Stieltjes (R^3S) integral. In: *Analysis, Geometry and Groups: A Riemann Legacy Volume*, (H. M. Srivastava and Th. M. Rassias, eds.), Part I, pp. 289-312; Hadronic Press, Palm Harbor, Florida, 1993.
- [66] E. R. LOVE, Integration by parts and other theorems for R^3S -integrals. *Real Analysis Exchange*, **24** (1998/9), 315-336.
- [67] E. R. LOVE AND L. C. YOUNG, Sur une classe de fonctionelles linéaires. *Fund. Math.*, **28** (1937), 243-257.
- [68] E. R. LOVE AND L. C. YOUNG, On fractional integration by parts. *Proc. London Math. Soc.*, Ser. 2, **44** (1938), 1-35.
- [69] T. LYONS, On the nonexistence of path integrals. *Proc. Roy. Soc. London Ser. A*, **432** (1991), 281-290.
- [70] T. LYONS, Differential equations driven by rough signals (I): an extension of an inequality of L. C. Young. *Math. Res. Lett.*, **1** (1994), 451-464.

- [71] J. S. MAC NERNEY, Integral equations and semigroups. *Illinois J. Math.*, **7** (1963), 148-173.
- [72] D. B. MADAN AND E. SENETA, The V. G. model for share market returns. *The Journal of Business*, **63** (1990), 511-524.
- [73] B. B. MANDELBROT, *Fractals and Scaling in Finance. Discontinuity, Concentration, Risk* Selecta volume E. New York, Springer, 1997.
- [74] M. B. MARCUS AND J. ROSEN, p -variation of the local times of symmetric stable processes and of Gaussian processes with stationary increments. *Ann. Probab.*, **20** (1992), 1685-1713.
- [75] P. R. MASANI, The place of multiplicative integration in modern analysis. Appendix II in [19].
- [76] J. H. MCCULLOCH, Financial applications of stable distributions. In: *Handbook of Statistics*, Vol. 14, G. S. Maddala and C. R. Rao (Eds.), 393-425. Amsterdam, Elsevier Science B.B., 1996.
- [77] R. M. MCLEOD, *The Generalized Riemann Integral*. The Carus Mathematical Monographs, No. 20. U.S.A., The Mathematical Association of America, 1980.
- [78] P. A. MEYER, A decomposition theorem for supermartingales. *Illinois J. Math.*, **6** (1962), 193-205.
- [79] P. A. MEYER, Decompositions of supermartingales; The uniqueness theorem. *Illinois J. Math.*, **7** (1963), 1-17.
- [80] E. J. MCSHANE, A theorey of limits. In: *Studies in Modern Analysis*, Vol. 1, Ed. R.C. Buck, pp. 7-29. U.S.A., The Mathematical Association of America, 1962.
- [81] E. J. MCSHANE AND T. A. BOTTS, *Real analysis*. Princeton, D. van Nostrand Company, 1959.
- [82] I. MONROE, On the γ -variation of processes with stationary independent increments. *Ann. Math. Statist.*, **43** (1972), 1213-1220.
- [83] R. NORVAIŠA, p -variation and integration of sample functions of stochastic processes. In: *Probab. Theory and Math. Statist.*, B. Grigelionis et. al. (Eds.), Proc. of the Seventh Vilnius Conference (1998), 521-540. TEV, Vilnius, 1999.
- [84] R. NORVAIŠA, Modelling of stock price changes: A real analysis approach. *Finance and Stochastics*, **4** (2000), 343-369.
- [85] R. NORVAIŠA, Chain rules and p -variation. Preprint.
- [86] R. NORVAIŠA AND D.M. SALOPEK, Estimating the Orey index of a Gaussian stochastic process with stationary increments: An application to financial data set. In: *Stochastic Models. A Conference in Honor of Professor Donald A. Dawson*, Eds. L. G. Gorostiza and B. G. Ivanoff; Conference Proceedings, **26**, pp. 353-374. Amer. Math. Soc. 2000.
- [87] H. OODAIRA, On Strassen's version of the law of the iterated logarithm for Gaussian processes. *Z. Wahrsch. verw. Geb.*, **21** (1972), 289-299.

- [88] S. OREY, F-processes. In: *Proc. Fifth Berkeley Symp. on Math. Stat. and Prob.*, II 1 (1965), 301-313.
- [89] M. F. M. OSBORNE, Brownian motion in the stock market. *Operations Research*, **7** (1959), 145-173.
- [90] R.E.A.C. PALEY AND N. WIENER, *Fourier transforms in the complex domain*. New York, AMS Coll. Publ. **19**, 1934.
- [91] R.E.A.C. PALEY, N. WIENER AND A. ZYGMUND, Notes on random functions. *Math. Z.*, **37** (1933), 647-668.
- [92] G. PISIER AND Q. XU, The strong p -variation of martingales and orthogonal series. *Probab. Th. Rel. Fields*, **77** (1988), 497-514.
- [93] S. R. PLISKA, *Introduction to Mathematical Finance. Discrete time models*. U.S.A., Blackwell Publishers, 1997.
- [94] S. POLLARD, The Stieltjes' integral and its generalizations. *Quart. J. Pure Appl. Math.*, **49** (1923), 73-138.
- [95] P. PROTTER, *Stochastic Integration and Differential Equations. A New Approach*. Berlin, Springer, 1990.
- [96] P. RÉVÉSZ, *Random Walk in Random and Non-Random Environments*. Singapore, World Scientific, 1990.
- [97] K. A. ROSS, Another approach to Riemann–Stieltjes integrals. *Amer. Math. Monthly*, **87** (1980), 660-662.
- [98] K. A. ROSS, *Elementary Analysis: The Theory of Calculus*. New York, Springer, 1980.
- [99] S. M. ROSS, *Stochastic Processes*. New York, John Wiley & Sons, 1983.
- [100] D.M. SALOPEK, Tolerance to arbitrage. *Stoch. Proc. Appl.*, **76** (1998), 217-230.
- [101] P. A. SAMUELSON, Mathematics of speculative price. *SIAM Review*, **15** (1973), 1-42.
- [102] Q.-M. SHAO, p -variation of Gaussian processes with stationary increments. *Studia Sci. Math. Hungar.*, **31** (1996), 237-247.
- [103] G. SOROS, *The Alchemy of Finance. Reading the Mind of the Market*. New York, John Wiley & Sons, 1994.
- [104] T. SZABADOS, An elementary introduction to the Wiener process and stochastic integrals. *Studia Sci. Math. Hungar.*, **31** (1996), 249-297.
- [105] S. J. TAYLOR, Exact asymptotic estimates of Brownian path variation. *Duke Math. J.*, **39** (1972), 219-241.
- [106] C. TRICOT, *Curves and fractal dimension*. New York, Springer, 1995.
- [107] B. Z. VULIKH, *Introduction to the Theory of Partially Ordered Spaces*. Groningen, Wolters-Noordhoff Scientific, 1967.

- [108] A. J. WARD, The Perron-Stieltjes integral. *Math. Z.*, **41** (1936), 578-604.
- [109] N. WIENER, The quadratic variation of a function and its Fourier coefficients. *J. Math. and Phys.*, **3** (1924), 72-94.
- [110] W. WILLINGER AND M. S. TAQQU, Pathwise stochastic integration and applications to the theory of continuoud trading. *Stoch. Proc. and their Appl.*, **32** (1989), 253-280.
- [111] Q. H. XU, Espaces d'interpolation réels entre les espaces V_p : propriétés géométriques et applications probabilistes. In: Séminaire d'Analyse Fonctionnelle 1985/1986/1987, Paris VI-VII, 77-123. Paris, Publ. math. Univ. Paris VII 28, 1988.
- [112] A. M. YAGLOM, Strong limit theorems for stochastic processes and orthogonality conditions for probability measures. In: *Bernoulli 1713, Bayes 1763, Laplace 1813*, Anniversary volume; eds. J. Neyman and L. M. LeCam; pp. 253-262. New York, Springer, 1965.
- [113] L. C. YOUNG, An inequality of the Hölder type, connected with Stieltjes integration. *Acta Math.* (Sweden), **67** (1936), 251-282.
- [114] L. C. YOUNG, Inequalities connected with bounded p -th power variation in the Wiener sense and with integrated Lipschitz conditions. *Proc. London Math. Soc.*, (Ser 2), **43** (1937), 449-467.
- [115] L. C. YOUNG, General inequalities for Stieltjes integrals and the convergence of Fourier series. *Math. Ann.*, **115** (1938), 581-612.